

2001 Project Abstract

For the Period Ending June 30, 2004

AUG - 9 2004

TITLE: Big Rivers Partnership: Helping Communities to Restore Habitat

PROJECT MANAGER: Deborah Karasov

ORGANIZATION: Great River Greening

ADDRESS: 35 West Water Street, Suite 201, Saint Paul, MN 55107-2106

WEB SITE ADDRESS: www.greatrivergreening.org

FUND: Environment and Natural Resources Trust Fund (TF)

LEGAL CITATION: ML 2001, 1st Special Session, Ch. 2, Sec. 14, Subd. 4(i)

FINAL REPORT

4i

APPROPRIATION AMOUNT: \$910,000

Overall Project Outcome and Results

Led by Great River Greening, the Big Rivers Partnership is one of the first important restoration collaborations in the state, bringing together nonprofit, government and private landowners to restore river valley habitat in the Twin Cities. Guided by ecological and resource criteria, projects were located within the important and beautiful Mississippi River Gorge running through Minneapolis and Saint Paul; the Pine Bend Bluff Natural Area, a regionally significant ecological resource on the urban Mississippi; the Minnesota River Valley, a critical and unmatched urban corridor of wetland and associated upland habitat; and numerous native plant community remnants. Projects consisted of plant and animal surveys and restoration activities that regularly engaged volunteers. More than 3,600 volunteers participated in habitat projects, triple the goal. The partnership also leveraged over \$1.3 million in non-state funds, almost double the goal, and implemented restoration on over 1,500 acres, 150% of the goal.

Project Results Use and Dissemination

Great River Greening lists projects and surveys on its website, promoting them to partners, cooperators, and landowners. Where possible, we work with cooperators to continue stewardship beyond state funding with volunteers or other community members. The Partnership also completed an ecological ranking of sites within the river valleys to complement the regionally significant areas identified by the Minnesota Department of Natural Resources and disseminated the ranking information through various conservation forums. The success of the project proves that multiple organizations can work together to achieve conservation goals.

Date of Report: July 13, 2004
Title: LCMR Final Work Program Report

Date of Next Status Report:
Date Work Program Approved: December 3, 2001
Project Completion Date: June 30, 2004
#CH34

I. PROJECT TITLE: Big Rivers Partnership: Helping Communities to Restore Habitat

Project Manager: Deborah Karasov
Affiliation: Great River Greening
Mailing Address: 35 West Water Street, Suite 201, Saint Paul, MN 55107-2016
Telephone Number: 651-665-9500 **e-mail:** dkarasov@greatrivergreening.org **Fax:** 651-665-9409

Web Address: www.greatrivergreening.org

Total Biennial Project Budget:
\$910,000 LCMR Appropriation - \$ 909,140 Amount Spent = \$ 860 Balance

Legal Citation: ML 2001, 1st Special Session, Ch. 2, Sec. 14, Subd. 4(i)

Appropriation Language: (i) Big Rivers Partnership: Helping Communities to Restore Habitat \$455,000 the first year and \$455,000 the second year are from the trust fund to the commissioner of natural resources for an agreement with Great River Greening to implement private and public habitat projects on cost-share basis in the Mississippi and Minnesota river valleys. This appropriation is available until June 30, 2004, at which time the project must be completed and final products delivered, unless an earlier date is specified in the work program.

Status of Cost-Share Requirement: Cost-share of \$691,771 will be provided for this project. Cost-share is defined as an expense that is directly related to the approved activities of the project (Big Rivers Partnership - BRP), that is not paid for with state funds, and is not an expense that is ineligible for LCMR reimbursement. Cost-share may be incurred by the project's recipient (GRG), partners (see Section VI-C), or landowners. Cost-share may include volunteer work, which would be valued using a rate of \$11.00/hr.

Documentation of cost-share is the following:

- Brief itemized description of the costs incurred (or the in-kind services provided) and their value
- Statement that certifies that the cost-share expenses reported are directly related to the approved activities of the Big Rivers Partnership.
- Statement that certifies that these expenses are NOT paid for with state funds.
- Signature by private individual landowner, or person authorized to represent the organization.

II. and III. FINAL PROJECT SUMMARY:

Overall Project Outcome and Results

Led by Great River Greening, the Big Rivers Partnership is one of the first important restoration collaborations in the state, bringing together nonprofit, government and private landowners to restore river valley habitat in the Twin Cities. Guided by ecological and resource criteria, projects were located within the important and beautiful Mississippi River Gorge running through Minneapolis and Saint Paul; the Pine Bend Bluff Natural Area, a regionally significant ecological resource on the urban Mississippi; the Minnesota River Valley, a critical and unmatched urban corridor of wetland and associated upland habitat; and numerous native plant community remnants. Projects consisted of plant and animal surveys and restoration activities that regularly engaged volunteers. More than 3,600 volunteers participated in habitat projects, triple the goal. The partnership also leveraged over \$1.3 million in non-state funds, almost double the goal, and implemented restoration on over 1,500 acres, 150% of the goal.

Project Results Use and Dissemination

Great River Greening lists projects and surveys on its website, promoting them to partners, cooperators, and landowners. Where possible, we work with cooperators to continue stewardship beyond state funding with volunteers or other community members. The Partnership also completed an ecological ranking of sites within the river valleys to complement the regionally significant areas identified by the Minnesota Department of Natural Resources and disseminated the ranking information through various conservation forums. The success of the project proves that multiple organizations can work together to achieve conservation goals.

IV. OUTLINE OF PROJECT RESULTS:

- **Result 1.** **LCMR Budget:** \$ 700,508 **Cost-Share Requirement:** \$691,771
 Balance: \$ 774 **Cost-Share Spent:** \$ 1,339,100

Implement habitat projects on 1000 acres with community volunteers

This project will build on the already burgeoning partnership for community restoration in the river corridors, spearheaded by Great River Greening. The partnership will implement habitat projects evaluated through GRG's ecological inventory and analysis, using GRG's successful collaborative process of soliciting landowners, recruiting and training volunteers and volunteer supervisors, training property owners, and providing restoration and management plans. Habitat projects will include activities such as plantings of native trees, shrubs, grasses and wildflowers, removal of exotic species and prescribed burning. Project sites that we intend to implement beginning in the spring of 2002 are Koch Refinery's Pine Bend Bluff property in Dakota County, the Minneapolis Longfellow River Gorge, and Belle Plaine Prairie. A nomination process to be completed early in 2002 will identify additional projects for 2002 and 2003.

Great River Greening has distinguished itself by the success and quality of its volunteer restorations and plantings largely because of its volunteer training program. Specialized training provides volunteer supervisors with opportunities to learn many skills, including native plant identification, principles of ecological restoration, and how to organize workers in carrying out activities for habitat restoration projects. Other volunteer field workers learn about ecological restoration by participating in habitat restoration activities. This project will recruit and train 1000 field workers and 75 supervisors, and will strengthen this training by focusing on two key areas: working with diverse groups of volunteers, and teaching principles and methods for restoring native plant

communities. This project will also strengthen our collaborations with community partners and landowners to develop restoration and adaptive management plans.

Budget:	Budgeted	Current	Balance
Personnel	\$ 390,756	390,489	267
Contractual Fees	139,442	139,060	382
Supplies/Equipment	160,748	160,623	125
Operations	5,737	5,737	0
Other	3,825	3,825	0
Result 1 - LCMR Subtotal	\$ 700,508	699,734	774

See attached worksheet for detail of expenses.

Completion Date:

- a) Implement habitat projects on 1000 acres
 - 1) 500 acres; December 31, 2002
 - 2) 500 acres; June 30, 2004
- b) Recruit and train 1000 volunteer field workers
 - 1) 500 volunteers; Dec. 31, 2002
 - 2) 500 volunteers; June 30, 2004
- c) Recruit and train 75 volunteer supervisors
 - 1) 45 supervisors; Dec. 31, 2002
 - 2) 30 supervisors; June 30, 2004

Result 1 Status:

June 30, 2004 Final Report: The Partnership completed all of its quantitative goals in this result. We implemented restoration on more than 1,500 acres. More than 3,600 volunteers participated in habitat projects, including over 300 volunteer supervisors. The partnership also leveraged over \$1.3 million in non-state funds.

Geographically, our work stretched from Belle Plaine and the west end of the Minnesota Valley National Wildlife Refuge to the Mississippi River Gorge in Minneapolis to Pine Bend Bluffs near Hastings. Site selection considerations encompassed both ecological and social factors. Ecological factors included the physical characteristics of site, the condition of the surrounding landscape, the plant community goal, and long-term maintenance and monitoring needs. Social considerations included screening/visibility, aesthetic needs, environmental needs, and sufficient management resources. Project sites included: East Mississippi Bluff-Desnoyer unit, Pine Bend Bluffs in Dakota County, Hastings Riverfront, Mississippi River Gorge-South Park in Minneapolis, West Side Bluff-Cherokee Park in Saint Paul, Minnesota Valley National Wildlife Refuge, River Park-Brooklyn Park, South Saint Paul Levee, and Prospect Crest in Saint Paul, to name a few.

The involvement of thousands of volunteers in our habitat projects helps to ensure that restoration will continue to occur far beyond the end of the BRP program funding period. For example, the Belle Plaine Elementary School Prairie is an integral part of the school. Our recent volunteer planting event at River Park, Brooklyn Park drew a tremendous response from the community, tapping into their volunteer spirit and providing educational opportunities for the volunteers, as well as direct habitat improvement on the Mississippi River bank.

As noted above, the Partnership leveraged over \$1.3 million in non-state funds. These dollars demonstrate the commitment from the community for such important work. Large contributors of cost-share included the Mississippi National River and Recreation Area (MNRRA), the National Fish and Wildlife Foundation (NFWF), and the City of Saint Paul.

- **Result 2. LCMR Budget: \$ 128,762**
Balance: \$ 56

Prioritize sites for ecological restoration and management

In partnership with the Minnesota Department of Natural Resources (DNR), National Park Service (NPS), and others, Great River Greening (GRG) will evaluate ecological inventory data of portions of the urban/suburban stretches of the Mississippi and Minnesota River valleys and habitat connections not covered by the Minnesota County Biological Survey (MCBS). MCBS looks only at the highest quality natural areas, not at those areas that could or should be restored. Using this GIS-based, landscape-scale ecological inventory data and social and economic criteria (e.g. landowner participation, BRP partner involvement, local community planning and interest), high-priority restoration project areas will be identified and mapped. Potential users of this mapped information will be municipal parks programs, DNR, NPS – Mississippi National River and Recreation Area, Trust for Public Lands, Friends of the Mississippi River, Friends of the Minnesota Valley and others.

In addition, we will conduct detailed site surveys at high-priority sites to collect information necessary to guide ecological restoration. These surveys will characterize infestations of exotic species, abundance of native species, social uses of sites, and other important factors.

Budget:	Budgeted	Current	Balance
Personnel	\$ 117,839	117,839	0
Contractual Fees	5,000	4,980	20
Supplies/Equipment	3,911	3,875	36
Operations	2,012	2,012	0
Other	0	0	0
Result 2 - LCMR Subtotal	128,762	128,706	56

See attached worksheet for detail of expenses.

Completion Date:

- a) Combine ecological inventory data with social and economic criteria to establish a prioritized map of restoration project areas
 - 1) Identify and map high-priority restoration project areas along the Mississippi and Minnesota River corridors; December 31, 2002 (mailed under separate cover 12/31/03)
- b) Conduct at least 4 detailed site surveys
 - 1) Conduct 2 surveys; December 31, 2002 (mailed under separate cover 12/31/03)
 - 2) Conduct 2 surveys; June 30, 2004 (attached)

Result 2 Status:

June 30, 2004 Final Report:

Great River Greening completed a land cover inventory using the Minnesota Land Cover Classification System for the portion of the Mississippi National River and Recreation Area (MNRRA) upstream of the Lower Saint Anthony Lock and Dam. In addition, Greening developed an ecological ranking protocol for MNRRA. The National Park Service, in conjunction with Greening, produced a CD-ROM which contains information on the ecological value of undeveloped land within MNRRA. The information provided on the CD, when used in combination with other regional or local data, can help identify open space protection opportunities (OSPO) within the river corridor.

Greening, in conjunction with the City of Saint Paul, also completed an ecological inventory (included in the larger restoration management plan) at Cherokee Park Prairie in Saint Paul. The plan makes recommendations, based upon the inventory, for the ongoing management of the vegetation to meet ecological goals and social needs. Recommendations include plantings of native plant species, actions to reduce/re-direct trails and prevent erosion, and the removal and control of invasive plant species that are degrading the ecological health of the bluff.

Great River Greening ecologists conducted surveys for four Mississippi River islands owned by the National Park Service in the BRP prioritized areas. Following an initial assessment by a Department of Natural Resources ecologist, Greening focused on these islands due to the immediate threat to their ecological quality from the invasion of exotic species, severe erosion, and/or human use. Two islands were located adjacent to Inver Grove Heights and two were adjacent to Anoka. For each island, Greening completed species lists and documented concentrations of invasive species, hydrologic conditions, soils, and structural conditions.

Greening ecologists also conducted a survey of plant communities in 68 acres of Cherokee Park (aka West Side Bluff Sector 1) that will guide restoration and management activities in the area adjacent to the oak forest and prairie remnant. This will further assist in the restoration, management and protection of these critical bluffland remnants.

In conjunction with Saint Paul Parks and Recreation, Great River Greening conducted a site survey of current trail conditions at the Bluff Trail at Crosby Park and created a design plan to guide improvements. The plan will help the City of Saint Paul manage the problematic Bluff Trail and will also act as a model for similar projects in the Twin Cities area.

- **Result 3. LCMR Budget: \$ 80,730
Balance: \$ 29**

Create a landowner stewardship support program.

Stewardship support and information sharing are critical to the efficiency and effectiveness of restoration. Technical information, on-going encouragement, and key stewardship services are essential to sustain current and future restoration efforts. Through a GRG River Steward Program, we will interpret and distribute relevant information through an information network (e.g. GRG website) to participating landowners, project cooperators, and natural resource managers. The River Steward Program will also provide additional services to selected participating landowners, including monitoring the condition of project sites and recommendations for volunteer stewardship activities.

Budget:	Budgeted	Current	Balance
Personnel	\$ 72,573	72,573	0
Contractual Fees	6,000	5,971	29
Supplies/Equipment	1,021	1,021	0
Operations	1,136	1,136	0
Other	0	0	0
Result 3 - LCMR Subtotal	\$ 80,730	80,701	29

See attached worksheet for detail of expenses.

Completion Date:

a) Compile information for distribution and install on website.

- 1) Compile technical information and install on website; June 30, 2002
- 2) Update information and website; December 31, 2002
- 3) Update information and website; June 30, 2003
- 4) Update information and website; June 30, 2004

b) Select and monitor at least 4 sites, and provide recommendations for volunteer stewardship activities.

- 1) Complete for 2 sites; Dec. 31, 2002. (mailed under separate cover 12/31/03)
- 2) Complete for 2 sites; June 30, 2004 (attached)

Result 3 Status:

June 30, 2004 Final Report:

We have updated and added information to our website. We developed a new and updated interactive project map with all our projects categorized by type of work. We added three management plans to the resources page: Bloomington Bluffs, Eagle Creek and West Side Bluff Action Plan Phase I, as well as fact sheets on storm water management. We also added the East-Central Minnesota Species list and a news page with the past year's project news releases and monthly e-postcards to update our constituents on Greening activity. We also made sign-up for Greening volunteer events easier with on-line registration. Technical information on Greening's website is now comprehensive and we have established a process for updating and expanding it. We have already notified our partners and other interested organizations of the website resource.

In 2002, Greening ecologists completed recommendations for stewardship activities for two important sites. Belle Plaine prairie is one of eleven remaining small native prairie remnants identified by the Minnesota County Biological Survey's 1995 inventory of Scott County. Pine Bend Bluff Natural Area is a unique natural area in the metropolitan area because of its size and quality, and a large portion is owned by Flint Hills Resources (FHR). For both of these sites, specific volunteer activities were identified within a larger stewardship plan, and volunteer groups were identified (school and company employees, respectively). Tasks for volunteers included cutting invasive species, replanting with native species, periodic weeding, and girdling trees.

In 2003, Greening finished an inventory and survey, included within a larger management plan, for the remnant savanna and forest in Cherokee Park (aka West Side Bluff Sector 1). This document provides guidance on how to incorporate volunteers in future planting/restoration events with West Side Citizens Organization and Saint Paul Parks and Recreation.

Working with employee volunteers at Flint Hills Resources, a Greening ecologist outlined and recommended 2004 activities for monitoring exotic plant population numbers and flowering, prairie seed

ripening, and acorn mast ripening, collection and storage. All of the Greening ecologist's recommendations have been accepted and are an integral part of the on-going restoration.

V. TOTAL PROJECT BUDGET:

All Results: Personnel	\$581,168
All Results: Contractual Fees	150,442
All Results: Supplies/Equipment	165,680
All Results: Operations	8,885
All Results: Other	3,825
Total Project Budget	\$910,000

See attached worksheet for detail of expenses.

In order to meet the cash flow needs of GRG, LCMR will advance \$60,000 to GRG at the beginning of the Big Rivers Partnership contract with the State of Minnesota. This amount is the most that GRG estimates it will need in any given month to cover expenses that are eligible for reimbursement by the State (see attached sheet detailing GRG's estimate of cash flow needs). In order to mitigate the interest lost by the State through this advance, GRG will hold any unused portions of the advance in an interest-bearing account that is transferable on demand to their checking account, and will deduct any interest earned on the balance each month from their requests for reimbursement. As the contract comes to a close, GRG will perform reconciliation in order to use the \$60,000 to pay the final reimbursements of the \$910,000 contract.

Final Report Status: We are submitting the final bill for \$130,003.06, of which \$60,000 has been advanced to us, and we received \$5.43 in interest on the advance in the month of June, 2004. Therefore we are requesting a final payment of \$130,003.06, less \$60,000, less \$5.43, which comes to \$69,997.63.

VI. PAST, PRESENT, AND FUTURE SPENDING:

A. Past Spending

GRG will serve as project manager for the proposed collaboration, which builds on the past work and present organization of the GRG project and its many partnerships. The budget numbers below relate to LCMR funding of the GRG project since its inception. In particular, this project builds on the successful Big Rivers Partnership, funded by an \$800,000 appropriation from the LCMR last biennium and matched by at least \$374,000 through public and private sources. The LCMR appropriation also leveraged \$50,000 of funds from the National Park Service for inventory work. In addition to the work sponsored by LCMR, this partnership of eleven government and community collaborators has stimulated hundreds of thousands of dollars of future projects. (LCMR History: \$1.1 million RIM Critical Habitat Match & LCMR appropriation; Non-LCMR History: \$2.2 million foundation, corporate, and private contributions.)

B. Current and Future Spending

In terms of future spending, this project will result in a database of prioritized future projects for restoration and management, which could not be included in the project period. The private landowners will provide matching resources. As well, the National Park service, which is one of our partners, awards approximately \$150,000 in planning grants to communities within the Mississippi Natural Resources and Recreational Area corridor; communities may leverage these funds with GRG resources. The National Park service also works with the MN DNR to award nearly half a million dollars in land use planning funds for the Mississippi Critical Areas Program; once communities complete planning they will be in a position to implement perhaps with GRG's assistance.

This proposal is also a complementary effort to the nearly \$6 million of proposed money for the Metro Greenways program. While that program focuses on planning, acquisition, and land protection, this project will stimulate on-the-ground stewardship expertise and work. For example, while the Greenways program may provide grants to local communities to complete resource inventories, this project may assist communities in implementing habitat improvements.

C. Project Partners

Great River Greening

MN Department of Natural Resources, Metro Region (in-kind contribution of ecological services; also technical support and housing for GIS ecological inventory)

Nelson French, Lori Nelson, Holly Buchanan, Kevin Bigalke, Friends of the Minnesota Valley

Whitney Clark, Tom Lewanski, Friends of the Mississippi River

Arne Stefferud, Metropolitan Council (in-kind contribution to coordinate implementing agencies)

Kate Hanson, Nancy Duncan, National Park Service (in-kind contribution of technical support for GIS)

Judy Barr, Rebecca Stenberg, City of Saint Paul (recipient of restoration services)

Cordelia Pierson, Trust for Public Land (in-kind contribution of landowner outreach)

Greg Mack, John Moriarty, Ramsey County Parks and Recreation (in-kind contribution of ecological services; also recipient of restoration services)

Rick Schultz, U. S. Fish and Wildlife Service (in-kind contribution for partnership development and planning; also recipient of restoration services)

Private landowners

D. Time

Restoration and planning require a minimum of two entire field seasons. We will begin the project January 1, 2002 and end it June 30, 2004.

VII. DISSEMINATION:

Volunteer events and supervisor training sessions are announced and reported on through the GRG newsletter and website. The map of the project area with high-priority project areas identified will be provided to the members of the Big Rivers Partnership. The Great River Greening website will be the primary tool for dissemination of landowner stewardship support and other information resulting from this grant.

VIII. LOCATION:

The Mississippi-Minnesota river corridors and habitat connections in the seven-county metropolitan region.

IX. REPORTING REQUIREMENTS:

Periodic work program reports will be submitted not later than June 30 and December 31, 2002, and June 30 and December 31, 2003. A final work program report and associated products will be submitted by June 30, 2004, or by the completion date as set in the appropriation.

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2002

EVENT	DATE (if event)	VOLUNTEERS	SUPERVISORS	ACRES	COST-SHARE	COMMENTS
PRING-SUMMER 2002						
Spr 02 Invasive Species Training	3/30/2002		28		\$ 924.00	GRG vol time @ \$11/hr
Spr 02 Basic Training	4/20/2002		35		\$ 1,155.00	GRG vol time @ \$11/hr
East Mississippi Bluff-Desnoyer unit	5/4/2002	30	6	20	\$ 3,806.00	City of St. Paul
					\$ 1,254.00	GRG vol time @ \$11/hr
Westside Blufflands-Smith Avenue	5/11/2002	125	21	8	\$ 40,805.00	City of St. Paul
					\$ 5,049.00	GRG vol time @ \$11/hr
Native Shrub Identif. Training	6/5/2002		32		\$ 1,056.00	GRG vol time @ \$11/hr
Barge Terminal No. 1, phase 1	6/8/2002	35	3	1	\$ 8,140.00	Wildlife Habitat Council cash
					\$ 1,287.00	GRG vol time @ \$11/hr
Carpenter Nature Center (FMR)		108		40	\$ 7,224.00	FMR
Pine Bend Bluffs (FMR)		15		54	\$ 50,000.00	FHR
					\$ 418.00	FHR volunteers
Hastings Riverfront plan (FMR)					\$ 7,300.00	FMR
SUBTOTAL Spring-Summer 2002		313	125	123	\$ 128,418.00	
FALL 2002						
Harriet Island entrance planting	10/10/2002	134	5	5	\$ 4,686.00	GRG vol time @ \$11/hr
Training for public parks	8/20/2002				\$ 528.00	GRG vol time @ \$11/hr
Mississippi River Gorge--South Park	9/21/2002	150	15	12	\$ 2,883.00	LCC
					\$ 5,610.00	GRG vol time @ \$11/hr
Mississippi River Gorge--other					\$ 5,170.00	MNRRA volunteers
					\$ 18,152.00	MNRRA inkind
					\$ 1,294.00	MNRRA tool purchases
Ford property crew buckthorn-cutting				20	\$ 89,466.00	Ford
Westside Blufflands-Cherokee Park		31	0	35	\$ 28,400.00	NFWF, federal
					\$ 6,868.00	WSCO inkind
					\$ 6,210.00	City of St. Paul inkind
Ordway Nature Area (FMR)	10/12/2002	68		7	\$ 17,074.00	Macalester inkind
					\$ 494.00	FMR
					\$ 2,244.00	FMR volunteers
Battle Creek north				35	\$ 45,000.00	MNRRA
Prospect Blvd.					\$ 4,709.00	City of St. Paul staff time
River Corrditor					\$ 2,000.00	survey staff/City of St. Paul
FMV 2002						
event totals:		1140		536	\$ 52,002.50	
non-state employee contributions:					\$ 24,979.02	
incurred expenses:					\$ 10,343.84	
mileage:					\$ 166.90	
FMV Projects:						
MN Valley Refuge--Louisville Swamp						
MN Valley Refuge--Long Meadow						
Belle Plaine						
Black Dog						
Visitor Center						
Pond Dakota Mission Park						
Minnegasco Dakota Station						
Fort Snelling						
SUBTOTAL Fall 2002		1523	20	650	\$ 328,280.26	
YEARLY 2002 TOTAL		1836	145	773	\$ 456,698.26	
REQUIRED, Result 1 2002		500	45	500	\$ 345,886.00	

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2003

EVENT	DATE(if event)	VOLUNTEER	SUPERVISOR	ACRES	COST-SHARE	COMMENTS
SPRING-SUMMER 2003						
Advanced Training	2/22/2003		19		\$ 627.00	GRG vol time @ \$11/hr
Advanced Training	4/12/2003		27		\$ 1,188.00	GRG vol time @ \$11/hr
Ford	4/26/2003	82		8.1	\$ 3,157.00	GRG vol time @ \$11/hr
					\$ 163,967.00	Ford
					\$ 7,954.59	Ford inkind
Minnesota National Wildlife Refuge:						
Louisville Oak Savanna Restoration	3/1/2003				\$3,653	USFWS Staff time
Ordway (FMR)	5/3/2003	62	4	1	\$ 12,670.00	Macalester inkind
					\$ 2,079.00	FMR vol time @ \$11/hr
Cherokee Prairie Implem	6/7/2003	10		13.3	\$ 330.00	GRG vol time @ \$11/hr
Mounds Park East Overlook	4/3/2003	55	6	0.125	\$ 6,433.00	City of St. Paul
					\$ 825.00	City vol time @ \$11/hr
					\$ 2,035.00	GRG vol time @ \$11/hr
Westside Blufflands-Cherokee Park	4/3/2003	33		35	\$ 1,408.00	GRG vol time @ \$11/hr
					\$ 17,000.00	NFWF 2003 Grant
Eagle Creek Invasive removal	5/17/2003	103	10	6	\$ 3,839.00	GRG vol time @ \$11/hr
Brooklyn Park River Park				12.8	\$46,000	MNRRA
35 West Water Street	6/10/2003			0.1	\$6,732	Mitigation
Crosby Exotic Removal					\$40,000	MNRRA
FMV 2003						
event totals:		448		314	\$ 15,697.00	
non-state employee contributions:					\$ 28,566.26	
incurred expenses:					\$ 2,363.34	
mileage:					\$ 83.96	
FMV Projects:						
Savage Fen						
Pond Dakota Mission Park						
Minnegasco Dakota Station						
Fort Snelling/Pike Island						
Long Meadow						
SUBTOTAL Spring-Summer 2003						
		793	66	390.425	\$ 366,608.15	
FALL 2003						
Pine Bend Bluffs (FMR) event	10/25/2003	100	10	7	\$ 50,000.00	FHR cash
					\$3,630	vol time @ \$11/hr
		8			\$88	FHR vol time @ \$11/hr
						FHR food, t-shirts, table tent & chair
					\$3,456	rental, biffs
South St. Paul Levee				11	\$7,000	City cash
					\$49,000	MNRRA cost share
Hastings Riverfront Park (FMR)				250	\$ 24,531.57	NAWCA
					\$ 1,168.00	Hastings city staff time
					\$ 1,800.00	Wetland plants
		35			\$ 990.00	vol time @ \$11/hr
3M (FMR)					\$ 9,460.00	3M cash (FMR)
Sand Coulee (FMR)				4.26	\$ 3,000.00	Landowner contrib.
		10			\$ 275.00	vol time @ \$11/hr
Minnesota National Wildlife Refuge	10/11/2003	71	6	30	\$ 2,673.00	GRG vol time @ \$11/hr
					\$ 2,000.00	plant material Refuge
					\$ 3,087.00	MN Refuge staff time
Land O' Lakes	9/26/2003	50	5	1	\$ 167,000.00	DCSWCD
					\$ 1,870.00	GRG vol time @ \$11/hr
Battle Creek	9/2/2003	165	2	19	\$ 5,533.00	cash and inkind
Rexam	8/19/2003	10		0.5	\$14,250	Mitigation/Cash
					\$330	GRG vol time @ \$11/hr
Training St Paul Parks & Rec	9/9/2003	20			\$660	GRG vol time @ \$11/hr
Advanced Training	11/1/2003		16		\$704	GRG vol time @ \$11/hr
Saint Paul Park & Rec. Projects:						
Cherokee					\$5,038	Staff time
Crosby					\$948	Staff time
Desnoyer					\$1,566	Staff time
Mounds					\$1,385	Staff time
Prospect Crest					\$1,484	Staff time
Lilydale Wetland					\$210	Staff time
SUBTOTAL Fall 2003						
		469	39	322.76	\$ 363,136.32	
YEARLY 2003 TOTAL						
		1262	105	713.185	\$ 729,744.47	
REQUIRED, Result 1 2003						
		500	30	500	\$ 345,886.00	

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2004

EVENT	DATE(if event)	VOLUNTEER	SUPERVISO	ACRES	COST-SHARE	COMMENTS
2004						
Crosby	2/21/2004				\$ 10,000	Carolyn Foundation
		46	3	4.5	\$ 1,650	GRG vol time @ \$11/hr
Advanced Training	4/3/2004		13		\$ 286	GRG vol time @ \$11/hr
Hastings River Flats	3/20/2004	87	8	11	\$ 3,223	GRG vol time @ \$11/hr
					\$ 1,459	FMR
Brooklyn Park River Park	5/1/2004	216	20	3	\$ 8,008	GRG vol time @ \$11/hr
					\$9,848	MNRRRA staff time
					\$81,000	MNRRRA
Prospect Crest	5/22/2004	70	5	0.25	\$ 2,530.00	GRG vol time @ \$11/hr
South St. Paul Levee	6/12/2004	126	18	0.25	\$ 4,950.00	GRG vol time @ \$11/hr
Flint Hills	5/1/2004			38	\$ 21,514	FHR
Dayton Install	5/1/2004			3.9	\$ 8,190	City of Dayton
SUBTOTAL, 2004		545	67	60.9	\$ 152,657	
YEARLY 2004 TOTAL		545	67	60.9	\$ 152,657	

Great River Greening
 Big Rivers Partnership
 LCMR Costshare Summary - Totals

EVENT	VOLUNTEERS	SUPERVISORS	ACRES	COST-SHARE
YEARLY 2002 TOTAL	1836	145	773	\$ 456,698.26
YEARLY 2003 TOTAL	1262	105	713.19	\$ 729,744.47
YEARLY 2004 TOTAL	545	67	60.9	\$ 152,657.41
Grand Total:	3,643	317	1,547	\$1,339,100.14
REQUIRED, Result 1 total	1,000	75	1,000	\$ 691,772.00

Fred Harris
Lead Ecologist
Great River Greening
24 February, 2004


Proposed Tasks for Flint Hills Volunteers in 2004

1. **Prairie forb seed broadcasting.** Very early May. 1 day (or split into 2 half days?) to hand broadcast prairie forb seed and work it into the ground with a hand pushed roller. Approximately 10 acre area in savanna restoration areas designated as SV-1b, SV-1c, SV-2. The purpose of this is to enhance the native species diversity in the savanna restoration area. Some site preparation with broadleaf herbicide will be done by GRG 2-3 weeks in advance.
2. **Weed control in native sand-gravel prairies.** Throughout the growing season in the native sand-gravel prairies. Mow sweet clover as it flowers; pull knapweed as it bolts and flowers. Repeat after 2 weeks. Remove all plant material from site. If there is time, pull garlic mustard at east end on ridge top.
3. **Collect and process acorns for fall event.** Mid to late August. Collect 200-300 bur oak acorns as they ripen and begin to fall from trees. Soak acorns in water for 24 hours. Store soaked acorns in burlap bags in refrigerators (GRG has some refrigerator space) until we plant them in the September 25 event. The timing of acorn collection is important: acorns have to be collected as soon as they fall due to competition from deer, turkeys and insects. Bur oaks generally ripen about August 20. In the September event, we will plant the acorns in the field where we have planted prairie grasses (SV-1a). In the following years we can thin out the trees to achieve the desired density for the savanna. This is the most ecologically appropriate way to restore savanna trees to the area as it utilizes the local genetic stock.
4. **Collect native prairie seed on sand-gravel prairies.** Late September – October. Collect native grass and forb seed and store in an unheated space over the winter. In early 2005, spread seed onto bare spots in sand-gravel prairie, such as in cheat grass control areas.

Exhibit A
Flint Hills Resources Bluffland Restoration Project- 2004
revised 3-19-2004

priority	1	5	2	4	3	
Work areas:	SGP-all	MP-1, SV-1a,	SV-1b, 1c, 2, 3, 4a, 4b	MP-1, SV-1a	SV-3	Total
Acres	7.26	12.135	19.287	24.022	6.6	69,304
Project Description:	SGP cheatgrass management. Continue cheat grass herb. & cutting in units set up in 2003, monitor cheat in 'I' that was burned in 2003, herb. in area of 'I' that was not burned in 2003, cut back brush 1-2 ft beyond where there are no grasses, stack brush off site.	Prescribed burn. Mow breaks, construct break through ravine in s. SV-1a to allow burn into shrubs.	Forb Enhancement and Establishment Mowing. Non-selective herbicide in late April, not including SV-3. FHR volunteers hand broadcast and rake forb seed into ground in early May. GRG to coordinate & monitor mowing contractor 2-4 times depending on weather & need. Stop mowing when natives are 4-6" tall or higher than weeds.	Fall Event (50 vols). FHR volunteers collect acorns in late August, soak for 24 hrs and refrigerate. During event, plant acorns in SV1a; broadcast and work in forb seed in SV1-a and MP-1.	Site Prep without restoration. GRG crew does 2 herbicide applications, plant oat cover crop (mowing covered in Forb Enhancement and Establishment Mowing column).	
Time Line:	Growing Season	Early October	April, May, June	October 23, 2004	Spring, Summer 2004	Total
Project management - FMR	\$1,000.00	\$1,000.00	\$1,000.00	\$1,000.00	\$1,000.00	\$5,000.00
FMR Ecologist*	\$2,000.00					\$2,000.00
Volunteer events staff & material:	\$0.00	\$0.00	\$0.00	\$6,325.00	\$0.00	\$6,325.00
Field work staff & material:	\$7,600.00	\$6,921.00	\$7,857.00	\$6,709.00	\$8,602.00	\$37,689.00
Subcontractors:	\$0.00	\$0.00	\$5,000.00	\$0.00	\$0.00	\$5,000.00
Cost Total:	\$10,600.00	\$7,921.00	\$13,857.00	\$14,034.00	\$9,602.00	\$56,014.00
BRP-GRG Funding			\$3,000.00		\$3,000.00	\$6,000.00
FHR funding:	\$10,600.00	\$7,921.00	\$10,857.00	\$14,034.00	\$6,602.00	\$50,000.00
Total Funding:	\$10,600.00	\$7,921.00	\$13,857.00	\$14,034.00	\$9,602.00	\$56,014.00
GRG 2004 funding	\$7,600.00	\$6,921.00	\$9,857.00	\$13,034.00	\$5,602.00	\$43,014.00
FMR 2004 funding	\$2,986.00	\$1,000.00	\$1,000.00	\$1,000.00	\$1,000.00	\$6,986.00
(Note: Cost estimate for 50 person volunteer event assumes FHR will provide lunch/soda and materials such as tents, tables, chairs, blfs etc.)						
	*GRG field staff will conduct management activities as described in project description above, at the appropriate times in 2004. FMR's Ecologist, Karen Schik, will monitor the managed areas throughout the field season, record pertinent data, and make management recommendations for the 2005 field season.		\$2040.00 estimated for plant material	\$2500.00 estimated for plant material		

Cherokee Park Management Plan



*One of several eroded paths cutting down the bluff that rises
above the Mississippi River in St. Paul*



July 31, 2003

**Prepared by: Great River Greening
35 West Water St., Suite 201
St. Paul, Minnesota 55107-2016**

Cherokee Park Restoration Management Plan

Compiled by
Ellen Fuge, Dan Shaw, and Shannon Farrell
Great River Greening

July 2003

Great River Greening (GRG), a nonprofit organization, helps communities coordinate cost-effective and sustained efforts to manage ecosystems of the Mississippi, Minnesota and St. Croix River valleys in the Twin Cities metropolitan area. We are primarily an implementing organization, providing on-the-ground ecological restoration and management of both public and private land. We engage thousands of volunteers in the planting of native vegetation, removal of exotic weeds, native seed collection and stewardship—work that results in an informed and involved citizenry. GRG also acts as a catalyst, creating effective partnerships among agencies, municipalities, and private landowners responsible for managing river valleys and their natural resources. Restoration ecologists and other scientists provide technical expertise. (See page 28 for more information about Great River Greening.)

Ellen L. Fuge has an M.S. in Botany from the University of Minnesota and currently works as the Lead Ecologist with Great River Greening. She conducts ecological inventories and analysis, writes restoration and management plans, and acts as a burn boss on the burn crew. Previously, she worked for many years with the Minnesota Department of Natural Resources in several different capacities; as the Management Supervisor for Minnesota's Scientific and Natural Areas (SNA) Program, and as a plant ecologist with the Minnesota County Biological Survey (CBS).

Dan Shaw is involved in restoration work, stormwater design and landscape planning at Great River Greening. He has worked as an ecologist for several years in both the public and private sector and is author of the publications *Plants for Stormwater Design* and *Native Vegetation in Restored and Created Wetlands*. He also teaches as an adjunct assistant professor in the Landscape Architecture Department at the University of Minnesota.

Shannon Farrell has a M.S. in Environmental Studies from the University of Strathclyde in Glasgow, Scotland and is working as a volunteer with Great River Greening assisting with writing management plans and associated field work. Previously, she earned a B.S. degree in Ecology, Evolution and Behavior from the University of Minnesota doing research projects and field studies on ecological issues concerning birds and insects. She is also involved in an internship with Friends of the Boundary Waters Wilderness working on regional wilderness conservation proposals.

Summary

The Cherokee Park Restoration Management Plan is an attachment to the West Side Bluff Ecological Inventory and Vegetation Management Plan (“West Side Bluff Management Plan”, 2001, GRG). The latter plan focuses on the inventory and management of Sectors 2, 3, and 4 (see Map 1) as designated by the West Side Bluff Task Force (West Side Bluff Management Plan, p. 22). The Cherokee Park Restoration Management Plan specifically addresses management of Cherokee Park (West Side Bluff – Sector 1) and presents recommendations for the ongoing management of the vegetation in Cherokee Park to meet ecological goals and social needs. The Cherokee Park Inventory Results (2002, GRG) is included in this plan as Appendix C.

The goals of the management recommendations are to identify ways to improve the ecological health of the bluff vegetation while also allowing for viewing areas and other uses. Recommendations include plantings of native plant species, actions to reduce bluff erosion, and removal and control of invasive plant species that are degrading the ecological health of the bluff. The management plan also identifies those tasks that can be conducted by volunteers and those that are more appropriate for trained professionals.

Appendices to the management plan provide technical information to supplement the recommendations, including detailed plant species lists of current and target native plant communities, information about controlling exotic species, and techniques and methods for many types of vegetation management.

Table of Contents	Page
Summary	03
List of Maps	05
List of Figures	05
List of Tables	06
Appendix Table of Contents	06
Acknowledgements	08
Introduction	09
Purpose of This Plan	09
Principles Guiding This Plan	09
Description of the Project Area	12
Bedrock, Soils and Erosion	12
Climate	15
Current Land Cover and Management Recommendations	17
Land Cover	17
Target Plant Communities	17
Management Units and Recommendations	18
Lawn	19
Side Slopes	21
Bottom Lands	24
Summary of Management Approaches	25
Bibliography and References	26
Great River Greening	28
List of Maps	
Map 1. Cherokee Park Location in West Side Bluff Project Area	31
Map 2. Cherokee Park Soils, Slope and Erosion	33
Map 3. Cherokee Park Existing Land Cover	35
Map 4. Cherokee Park Management Units and Target Plant Communities	37
Map 5. Cherokee Park Cultural Features	39
List of Figures	
Figure 1. Cherokee Park West, section A' – A with the three management units delineated	19
Figure 2. Conceptual Oak Savanna restoration along the edge of the slope that would reduce mowed lawn area, restore native vegetation, and protect slope from erosion.	20

List of Tables

Table 1: Monthly Station Normals of Temperature and Precipitation 1971 – 2000	15
Table 2: Median Frost Dates Derived from 1971-2000 Averages	15
Table 3: Mean Number of Snow Cover Days for Indicated Depths and the First and Last Dates of 1-inch Snow Cover October 1959 – May 1979	16
Table 4: Cherokee Park Management Unit Summary	18
Table 5: Management Unit Priority Status, Management Actions and Volunteer Involvement	25

Appendix Table of Contents

Appendix A. Fact Sheets for Exotic and Invasive Plants

Forbs

Canada thistle	A-3
Purple loosestrife	A-5
Sweet clover	A-7

Grasses

Bluegrass	A-9
Reed canary grass	A-11
Smooth brome grass	A-13

Appendix B. Species Lists for Proposed Restoration Target Communities

Table 1. Dry Prairie Species List	B-1
Table 2. Oak Savanna Species List	B-3
Table 3. Maple-basswood Forest Species List	B-5
Table 4. Dry Oak Forest Species List	B-8
Table 5. Lowland Hardwood Forest Species List	B-10
Table 6. Floodplain Forest Species List	B-12
Table 7. Mixed Emergent Marsh Species List	B-14
Table 8. Willow Swamp Species List	B-17

Appendix C. Cherokee Park Inventory Results

Introduction	C-1
Maple-basswood forest	C-1
Short grasses and mixed trees with 26-50% impervious cover	C-3
Short grasses and sparse tree cover on upland soils	C-5
Boxelder-green ash disturbed native forest (Northwest portion)	C-6
51-75% Impervious cover with deciduous trees (Northern portion)	C-8
51-75% Impervious cover with deciduous trees (Southern portion)	C-9
Disturbed deciduous woodland (Northern portion)	C-10
Disturbed deciduous woodland (Southern portion)	C-12
Boxelder-green ash disturbed native forest (Southeast portion)	C-13
Lowland hardwood forest	C-15

Mixed emergent marsh (Northern portion)	C-16
Mixed emergent marsh (Southern portion)	C-17
Willow swamp	C-19
Floodplain forest	C-20
Dry prairie/savanna-sand gravel subtype (including prairie edges)	C-21
Oak forest	C-23

Appendix D. Earthworms

Without earth worms	D-1
With earth worms	D-7
Sampling methods for earthworms	D-12

Appendix E. Resources

Contacts	E-1
Equipment	E-1
Native plant community restoration	E-2

Acknowledgements

The Cherokee Park Restoration Management Plan is the product of a collaborative project between Great River Greening, the West Side Bluff Task Force of the West Side Citizens Organization, and the City of Saint Paul Division of Parks and Recreation. This project is one of several funded by the Big Rivers Partnership, a large partnership led by Great River Greening. Funding for the Big Rivers Partnership is provided by the Minnesota Environment and Natural Resources Trust Fund, as recommended by the Legislative Commission on Minnesota Resources.

Introduction

Purpose of This Plan

This plan provides recommendations for managing the City of St. Paul's Cherokee Park vegetation in a way that meets the various needs of bluff residents and visitors while also being cost-effective and ecologically sustainable. Specifically, the plan recommends ways to maintain viewing areas from the bluff, while also increasing biological diversity, reducing bluff erosion, increasing landscape aesthetics, improving wildlife habitat and reducing ongoing maintenance costs.

Community interest in Cherokee Park is high. Active community members in the West Side neighborhood and city agency staff have agreed that a sustainable natural resource management approach is crucial to ensuring successful management of the bluff. Vegetation management is best done with an adaptive approach, which recognizes that as management recommendations are implemented, monitoring will be needed to evaluate the results and continue to refine and develop the management plan.

Principles Guiding the Plan

The fundamental principle of the Cherokee Park Restoration Management Plan is that management must be based on ecological and social goals. The Cherokee Park ecosystem is the interacting group of plants, animals (including humans), and physical elements (slope, soil, climate, water) at the site. These interactions need to be considered in developing goals for managing the vegetation in the park. An ecosystem approach to management acknowledges that people are included in the natural system and that maintaining a healthy and diverse ecosystem is the best way to meet the needs of park users, bluff residents and all the other organisms living on the site. Ecosystem management integrates current scientific knowledge and human values with the underlying goal of protecting the health of the ecosystem for the long term. Following are principles used to guide the development of this management plan.

1. Our management efforts should protect or enhance the health of the bluffland ecosystem and the native biological diversity of its habitats. Protecting and restoring high-quality natural areas and the ecological processes that sustain them are high priorities, because these sites are much more biologically diverse than other areas. The native plants and animals of Cherokee Park have evolved together for thousands of years and are particularly suited to surviving and thriving in the natural communities present at this site. Over the long term, native communities will maintain a level of health that can adapt to disease, drought, flood, fire, wind and other natural disturbances, and therefore should require less effort to maintain and manage than degraded or exotic communities. These native communities offer a varied and interesting environment, providing people with many opportunities for recreation and chances to learn about our natural heritage. Finally, high-quality natural areas are increasingly rare worldwide. They are worth protecting because they are rare and difficult and, in some cases, impossible to restore. The longer they are protected, the more valuable they will become. Once high-

quality areas are secure, attention should be given to buffering and/or connecting these areas and restoring lower-quality areas.

2. Planning should recognize that species are interdependent. Individual plant and animal species in a community depend on one another for survival. Relationships and interactions among species are complex and still poorly understood. Saving plant communities where these species and their interactions occur should be a management priority.

3. Planning should acknowledge that people are part of nature. Human actions have been influencing Cherokee Park for hundreds of years. Plans for managing the park's vegetation should recognize that people are a part of this landscape. Appropriate recreational and viewing opportunities as well as cultural and historical resources should be a part of plans for ecological management.

4. Planning should be based on ecological boundaries, not simply political boundaries, and should be based on extended time frames. The natural communities and systems that make up Cherokee Park have existed for thousands of years and extend across the property ownership boundaries. Developing common management goals with surrounding landowners will ensure the long-term health of the park's important natural resources.

5. Management planning should be based on an adaptive approach. Adaptive management refers to an ongoing approach to effectively manage projects. Adaptive management starts with project planning, a thorough site inventory and development of the adaptive management plan. The plan involves discussion of how vegetation establishment should be conducted at a site as well as clear and measurable goals for how monitoring should be conducted to detect changes/responses at the site. Every site is unique and constantly changing and, as a result, monitoring is a key component of any project. Monitoring provides information about the changes that are occurring at a project and about the success of management efforts. Monitoring involves evaluating the development of a project and provides information about how the adaptive management plan should be changed to modifying future practices to increase the effectiveness of management efforts. Ultimately, the adaptive management approach allows for unique responses of a site to management efforts and provides an effective method to adjust management strategies

6. Exotic species should be excluded or carefully controlled. Introduction of exotic species (see examples in Appendix A) can reduce native diversity, the quality of habitat and the general health of the bluff's natural resources. Therefore, exotics should be excluded or carefully controlled.

7. Management should be based on cooperative efforts. In addition to the City of Saint Paul Division of Parks and Recreation and the West Side Bluff Organization, many other individuals, organizations and agencies affect Cherokee Park's resources. Decisions made by surrounding landowners and agencies can alter the

blufflands. For example, undesirable invasive, exotic species planted by neighbors could invade and degrade adjacent city-owned natural areas. Because so many groups and individuals influence the ecology of the park, the Management Plan should not be developed in isolation.

Description of the Project Area

The surrounding landscape, soils, geology and current land cover and of a site all provide clues about an area's ecological condition and how the site should be managed. This section of the management plan looks at the climate of the area in addition to the geological, soil and land cover conditions of Cherokee Park. The larger landscape around Cherokee Park is discussed in the West Side Bluff Management Plan.

Bedrock, Soils and Erosion

Soil type is a major factor controlling a site's hydrologic characteristics, the likelihood of erosion, and the vegetation of the site. Three major soil types are present on three different portions of the bluff: Dorerton soils are present on the slope, Udorthents on the bluff base and Kingsley or Copaston soils on blufftop. Map 4 (page 90) describes the locations of these soil types in Cherokee Park.

The bedrock of Cherokee Park is primarily composed of sandstone, shale and limestone. The bedrock on the site affects root growth and influences the park's hydrology.

Factors contributing to erosion

Soil type: Dorerton-rock outcrop soils are prone to erosion because of their steep slopes, small particle sizes and low organic content. Walking trails and areas where water is directed down the bluff are located on these soils and are currently sources of erosion on the entire West Side Bluff. At Cherokee Park, heavy rains in June 2003 washed out a large hole at the head of a deeply eroded ravine in the middle of the park (see Map 2). This damage extends the erosion another 20 feet up-slope toward the top of the bluff.

In addition to changing runoff patterns from streets, parking lots and lawns, replacing mowed lawn with a band of taller native woodland edge species would help to slow the flow of water to the bluff edge and alleviate this kind of damage.

Significance of organic matter and soil organisms: Erosion is common on bare slopes that lack vegetation or organic matter. Organic matter plays an important role in controlling erosion by slowing water as it moves over a slope, absorbing moisture and providing nutrients for ground-layer woodland plant species. The organic layer of a healthy forest floor is generally composed of accumulated leaves and twigs as well as roots, bulbs, seed and fungi. Soil organisms including bacteria and fungi slowly decompose accumulated organic material, but new leaves and twigs continually regenerate the forest floor. The high productivity and slow decomposition of the forest results in the development of a thick organic layer. The accumulated plant material is generally loose and spongy providing ideal conditions for root growth of woodland plants and cool, moist conditions for seed germination. The organic layer also provides a good insulating layer during the winter.

Mycorrhizae are particularly important to the health of woodland plants. These specialized structures are formed through a symbiotic (mutually beneficial) relationship between a plant's roots and specialized mycorrhizal fungi. The fungal symbiont adds an

extensive network of root-like filaments to the plants roots. The expanded root system provides more nutrients and water for plants and, in turn, plants supply carbohydrates to the fungi.

Because of the crucial roles of both organic matter and mycorrhizal fungi, increasing the organic matter on the bluff is important and should be conducted in combination with tree and shrub plantings by applying thick layers of wood chips or, preferably, shredded bark. Highly degraded areas will benefit from the reintroduction of mycorrhizal fungi. One method for reintroducing mycorrhizae is to broadcast a site with wood chips collected from trails or plantings within healthy forests (Sauer 1998).

One of the greatest threats to the structure of hardwood forests is the presence of earthworms (see a more detailed discussion in Appendix D). All earthworms found in this area are non-native and damage forests by quickly consuming the organic layer. Earthworms consume leaves and other organic material on the forest floor, exposing the roots of woodland plants and preventing their growth. Soil exposed after native plants disappear is often colonized by weedy or invasive species that thrive on disturbed sites. Earthworms also consume bacteria and fungi that are essential to the normal functions of the forest floor.

Soil type on the slope – Dorerton-Rock Outcrop Complex

The most common and ecologically significant soil on the project site is the Dorerton-rock outcrop complex soil found on the slope of the bluff. Dorerton-rock outcrop complex soils are generally found on 25 to 65 percent slopes and are common in stream valleys. The soil complex is composed of 50 to 75 percent Dorerton soils and 15 to 20 percent outcrop. The Dorerton soil generally has a surface layer of dark gray sandy loam about 4 inches thick over about a 6 inch-thick subsurface layer. The subsoil is about 12 inches thick and is dark brown gravelly clay loam in the upper part and dark brown flaggy clay loam in the lower part. Underlying material is pale brown flaggy loamy sand.

Permeability of the Dorerton soil is moderate to moderately rapid, indicating that the bluff slopes are generally dry. This soil type favors plant communities that can survive low soil moisture, such as oak forest, oak savanna, and brushland. However, in ravines and the base of cliffs, where soil moisture is higher, plant communities requiring higher moisture, such as maple-basswood forest, are favored.

Soil type at the base of the bluff

Sediment and rock eroded from the steep bluffs collect at the base of the bluff. Beneath these eroded sediments are wet substratum soils called udorthents. Udorthents consist of fill material and industrial waste that has been placed on poorly drained and very poorly drained mineral or organic soils. This fill material provides sites for buildings, roads, recreation areas and other uses. Permeability and available water capacity of urban fill soils varies. Many areas are highly compacted, and in these locations, water collects on the surface after heavy rainfall. Runoff and internal drainage are also variable, and the depth to the seasonal high-water table varies from 1 foot to more than 6 feet. In most

parts of the project area where udorthents are present, they are covered by the accumulated soil and rock that have eroded from the bluff above.

Soil type on the blufftop

Soils on the blufftop consist of a variety of urban land complexes. Urban land-Kingsley-complex and urban land-Copaston complex are the most common. The urban land portions of these map units are covered by roads, parking lots, buildings and other structures. Changes to the soils have obscured or altered the original appearance so significantly that identification is not feasible. However, original soils may still be found in yards that have been established for many years. Savanna vegetation helped form the dark soils of the Kingsley and Copaston soils. Decomposition of the deep root systems of the prairie grasses and forbs of the savanna added organic material to the soil, contributing to the soil's dark color.

Kingsley or Copaston soils that remain both consist of dark surface layers from 6 to 8 inches deep. The surface layer of Kingsley soils consists of sandy loams, while Copaston soils consist of loam. The Kingsley subsoil is sandy loam and about 26 inches deep, and the Copaston subsoil is about 9 inches deep. Kingsley and Copaston soils differ in their permeability. Kingsley soils are moderate in the surface layer and moderately slow in the subsurface. Available water capacity is moderate and runoff is rapid. Copaston soils have moderate permeability, water capacity is high because of underlying bedrock, and runoff is moderate.

Bedrock of the site

Limestone bedrock underlies the soil at a depth of 45 to 70 inches. In Cherokee Park, bedrock is primarily a combination of sandstone, shale and limestone. The sandstone layer, known as St. Peter sandstone, was formed as large inland seas slowly filled with sand that eroded from surrounding uplands. The sand compressed over time, binding the sand grains into stone. St. Peter sandstone is found at the base of the bluff where caves have formed from natural processes and human excavation. Above the sandstone is a layer of shale that formed from mud deposition on top of the sandstone. Fossils such as brachiopods, gastropods and trilobites are common in the shale. Water levels rose after the mud layer developed. The higher water level allowed many organisms with calcium shells to thrive. A layer of limestone was formed from the chemical precipitation of calcite and the remains of animal life. The limestone at Cherokee Park, known as Platteville limestone, is about 30 feet thick. Above the limestone, another layer of mud formed as the sea receded. This mud layer formed into what is known as Decorah shale. A layer of glacial deposits or drift, consisting primarily of rock, gravel and sand deposited by glacial action, overlays the bedrock.

Bedrock and glacial deposits on the site can influence plant growth in a number of ways. Both bedrock and drift can influence the pH and the hydrology of soils depending on the composition and structure of the materials. Exposed bedrock lacking soil supports scant vegetation. Trees that do grow on bedrock may have a higher chance of being blown over during storms, especially if erosion is exposing their roots. Where bedrock is cracked or composed of soft sandstone, water can travel and seep out of the bluff. Seeps will

generally have species that require high amounts of moisture and are often prone to erosion. Eroding bedrock affects vegetation by smothering some species and creating new areas of growth that favor rapidly establishing species.

Climate

The climate of a site is an extremely important component of the resources and determines what species can grow and sustain themselves. Temperature and moisture are particularly important. This site is located in a typical continental climate with moderate precipitation and wide ranges in temperature from summer to winter. The west-facing aspect of the bluff

The climatological information relevant to Cherokee Park is based on the data from the weather data collecting station at St. Paul, Minnesota. The monthly normals for temperature range from a minimum of 6.2 °F in January to a high of 83.2 °F in July. Precipitation ranged from .76 inches in February to 4.98 inches in August with an annual average precipitation of 32.59 inches.

Table 1. Monthly Station Normals of Temperature and Precipitation 1971 – 2000

Temperature Normals (degrees Fahrenheit)														
Station Name	Element	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
St Paul	Max	22.8	29.7	41.7	58.2	71.2	79.1	83.2	80.8	71.8	59.4	40.5	26.7	55.4
	Mean	14.5	21.4	32.8	47.2	59.9	68.4	73.0	70.8	61.8	49.8	33.3	19.5	46.0
	Min	6.2	13.0	23.9	36.2	48.5	57.6	62.7	60.7	51.7	40.1	26.1	12.3	36.6

Precipitation Normals (Total in inches)														
Station Name		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Annual
St. Paul		1.02	.78	1.92	2.54	3.73	4.98	4.41	4.37	3.20	2.51	2.09	1.04	32.59

From: Climatology of the United States No. 81. National Oceanic and Atmospheric Administration. United States Dept. of Commerce. National Climatic Data Center, Ashville, NC. December 1, 2001.

The average date of the last freezing temperature (32 °F) in spring occurs on May 2 and on an average, the first freezing temperature in fall occurs on September 19 resulting in an average of 159 days free of frost in between.

Table 2. Median Frost Dates Derived from 1971-2000 Averages

Station Name	Median Date of Last Min. Temp. in Spring			Median Date of First Min. Temp. in Fall			Days in Between Median Dates		
	24°F	28°F	32°F	32°F	28°F	24°F	32°F	28°F	24°F
St. Paul	04/08	04/19	05/02	10/04	10/18	10/31	159	186	210

From: The Midwest Regional Climate Center, Champaign, IL. Available: http://mcc.sws.uiuc.edu/html/Mwclimate_data_summaries.htm (Accessed: June 26, 2003)

The average date of the first 1-inch snowfall in the fall is November 22 and the average date of the last 1-inch snow cover in the spring is April 2. The average number of days when there is a snow cover greater than 1 inch is 16.3. The annual average snowfall is 52.4 inches.

Table 3. Mean Number of Snow Cover Days for Indicated Depths and the First and Last Dates of 1-inch Snow Cover, October 1959 – May 1979.

Station	Average seasonal snow cover days					Average date of last 1" snow cover in the spring	Average date of first 1" snow cover in the fall
	1"	3"	6"	12"	24"		
St. Paul	100	79	54	24	1	November 22	April 2

From: Climate of Minnesota, Part XIII – Duration and Depth of Snow Cover. Kuenhast, E. L., D. G. Baker and J. A. Zandlo. Tech. Bull. 333-1982. Agricultural Experiment Station, University of Minnesota.

Current Land Cover and Management Recommendations

Land Cover

Land cover is defined as the physical cover, including vegetation (natural or planted) and human constructions (buildings, roads, etc.) present on the landscape. Information about existing land cover can help guide decisions about what human uses are appropriate at a site, where restoration efforts should be focused and what plant communities should be connected. Map 3 presents the land cover for the Cherokee Park project area.

The land cover map for the Cherokee Park project is based on work completed under a cooperative agreement between the National Park Service, the Minnesota Department of Natural Resources and Great River Greening. The principal objective of this agreement was to complete a land-cover inventory using the Minnesota Land-Cover Classification System (MLCCS) (Leete et al. 2000, Map 2) for the entire Mississippi National River and Recreation Area and additional areas.

The land cover data collected for Cherokee Park served as a framework for more detailed surveys conducted at the site. Additional background information was gleaned from the Minnesota Natural Heritage Database, Minnesota County Biological Survey data and the Ramsey County Soil Survey.

Land cover types in the Cherokee Park project area include:

- Dry oak savanna sand-gravel subtype
- Oak forest
- Maple-Basswood forest
- Lowland hardwood forest
- Floodplain forest
- Willow swamp
- Mixed-emergent marsh
- Disturbed deciduous woodland
- Boxelder - green ash disturbed native forest
- Short grasses with sparse tree cover on upland soils
- Short grasses and mixed trees with 26-50% impervious cover
- 51 to 75% impervious cover with deciduous trees
- Buildings and pavement with 76-90% impervious cover

Target Plant Communities

The determination of target plant communities was made by considering the historic vegetation of the area and the current land cover. According to the land surveyors of the 1850's, most of the uplands were covered with oak openings or savanna. The slopes of the bluff supported oak forest and maple-basswood forest and the bottomlands were floodplain forest and wetlands. Today, remnants of the original vegetation are still apparent although highly disturbed. Oak savanna, oak forest and maple-basswood forest

Management Units and Recommendations

The Cherokee Park site is divided into management units based on variations in the land use, terrain, and vegetation (Map 4). On this basis, three general management units are identified: Lawn, Side Slopes, and Bottom Lands. The following sections present management recommendations for each management unit. Priority status is based on current ecological condition, management history, future management and social considerations.

The following table describes the current conditions of each management unit, the land cover types it encompasses, and the recommended target plant communities.

Table 4: Cherokee Park Management Unit Summary


Management Unit	Description	Land-cover types	Target Plant Communities
Lawn	<i>Activity areas including picnic areas, playgrounds, parking lots, and sidewalks are prominent features in this unit which occupies the generally level, top part of the bluff. The large oaks scattered throughout this unit are reminiscent of the native oak savannas found here in presettlement times. The principal vegetation management is mowing.</i>	<ul style="list-style-type: none"> • Short grasses and mixed trees with 26-50% impervious cover • Short grasses and sparse tree cover on upland soils • Buildings and Pavement with 76-90% Impervious Cover 	<ul style="list-style-type: none"> • Oak savanna
Side Slopes	<i>The steep slopes of the bluff face are criss-crossed with paths used by people and animals, especially deer. This activity, water runoff from the lawns and pavements above, and invasive earthworms have removed much of the ground cover throughout the unit. Never the less, some of the most intact native plant communities are found in this unit.</i>	<ul style="list-style-type: none"> • Maple-Basswood Forest • Oak Forest • Dry Oak Savanna Sand-gravel Subtype • Disturbed Deciduous Forest • Boxelder-Green Ash Disturbed Native Forest 	<ul style="list-style-type: none"> • Oak Forest • Maple-basswood Forest
Bottom Lands	<i>In addition to trails, roads and past use of adjacent areas as "brick yards", frequent floods periodically disturb a large part of the Bottom Lands unit. These activities and exotic invasive plants, such as reed canary grass and buckthorn have extensively disturbed the plant communities of this unit. Debris is common throughout the floodplain forest floor.</i>	<ul style="list-style-type: none"> • Floodplain Forest • Willow Swamp • Mixed Emergent Marsh • Lowland Hardwood Forest • 51-75% Impervious Cover with Deciduous Trees 	<ul style="list-style-type: none"> • Lowland Hardwood Forest • Floodplain Forest

As in Sectors 2, 3 and 4 (refer to West Side Bluff Management Plan, pp. 22-27), mowed lawn areas are maintained on the top of the bluff in Cherokee Park along the streets and in activity areas around parking lots and picnic grounds. The early vegetation of this level upland was typically oak savanna, which was developed for housing and city streets early in the history of the city.

Some patches of oak forest, maple basswood forest and dry-prairie along the side slopes or bluff face contain moderate native plant diversity with lower degrees of disturbance

and invasive exotic plants. These areas are among the most intact native plant communities of the West Side Bluff and are collectively cited by the Minnesota County Biological Survey as “a site with high biodiversity significance”. The bluffs of Cherokee Park are predominantly west facing and receive more sun exposure than the rest of the West Side Bluff, which generally faces north. Increased sun exposure creates hotter and drier conditions favoring prairie, savanna and oak forest. Remnants of maple-basswood forest are found in areas that are cooler, shadier and moister, such as on north-facing slopes, in ravines and at the base of slopes.

The bottom lands are heavily disturbed with only very tolerant native species remaining in the Floodplain Forest, Lowland Hardwood Forest, Willow Swamp and Mixed Emergent Marsh native plant communities found there.



The Cherokee Park Inventory Results, completed in 2002, can be found in Appendix C.

Management recommendations for the Dry Oak Savanna Sand-gravel Subtype remnant in the Side Slopes Management Unit are not included within this document. In December 2002, the Cherokee Park Prairie - Ecological Inventory and Restoration Management Plan was completed by Daniel B. Shaw to guide the protection, restoration, and management of this unique area.


Lawn

Target Plant Community: Oak Savanna, Oak Woodland Brushland

Priority Status: 1

Description: This management unit has two parts: the first following Cherokee Heights Boulevard and Cherokee Avenue, and the second being the southern-most recreational portion of Cherokee Park (with picnic tables, a playground, and restrooms) (Map 5). A vegetation inventory of the lawn management unit can be found in Appendix C, pp. C-5 – C-7, in the sections titled “Short grasses and mixed trees with 26-50% impervious cover” and “Short grasses and sparse tree cover on upland soils.”

Management Approach: Currently, mowed lawn with various scattered landscape trees are maintained up to the edge of the bluff, contributing to increased erosion from water runoff and inviting foot traffic along sensitive areas of the bluff’s edge. To improve water infiltration and slow erosion on the bluff, convert a minimum of 5 to 10 feet of existing lawn to native plants typical of a woodland edge or oak savanna. Erosion would be abated and weedy species would be replaced by prairie and savanna trees, shrubs, grasses and flowers. A broader band of tall vegetation along the edge of the bluff would serve to protect this highly erodible edge from excessive pedestrian traffic as well. The large grove of mature oak trees shown on Map 4 and referenced on page C-5 of Appendix C, is an example of a larger area for potential oak savanna restoration by re-introducing native ground layer grasses and forbs (flowering plants). Some of the planted non-native trees, especially Norway maple, are spreading to near-by natural areas. Replacing these with native tree species (e.g. bur oak) and converting other potentially invasive non-native plantings to natives in the future should be considered.



The first phase of restoration involves the application of herbicide to the areas of lawn to be converted to woodland edge or savanna. A herbicide with a combination of a broad-spectrum herbicide (such as Rodeo) and a broad-leaf weed killer (such as Garlon) is used to ensure that all current vegetation is eliminated. Two applications are recommended to ensure that hard-to-eliminate species such as quack grass, smooth brome, burdock and Canada thistle are removed from the site. If a large enough area is to be planted, a seed drill can be used to plant native grass seed directly through the dead turf. If a small strip

were planted, removing the dead grass by burning would be advisable. Then the seed could be hand broadcast and raked into the exposed soil.

The final step in the process is conducted after prairie grasses are established and involves the planting of forb species, either as potted plants or plugs. After forbs are added to the planting, fire will be the most useful management technique for eliminating weeds from the planting. However, if perennial weeds persist or if burning is not possible, targeted mowing or herbicide application may still be necessary.

Long-term management of the site would involve a combination of mowing and burning. If weeds become common, they must be cut with a weed cutter before they reach one foot in height, and definitely before they flower and go to seed. The planting should be monitored for invasion of problem weed species such as Canada thistle, quack grass, smooth brome or burdock. After 3 or 4 years, the savanna planting can be managed through burning.

Many of the invasive and weedy species at the site will persist into the future. Long-term management involves monitoring of the site by a trained ecologist and adjustment of the management plan as the site changes.

Side Slopes

Target Plant Communities: Maple-Basswood Forest, Oak Forest

Priority Status: 1

Maple-Basswood Forest

Inventory Results: A detailed inventory of the Maple-Basswood Forest land cover type and the areas targeted for Maple-Basswood Forest restoration can be found in Appendix C in the sections titled: "Maple-basswood forest," "51-75% Impervious cover with deciduous trees (Northern portion)," "Disturbed deciduous woodland (Northern portion)," "Disturbed deciduous woodland (Southern portion)," "Oak forest," and "Boxelder-green ash disturbed native forest (Southeast portion)."

Management Approach: Vegetative diversity is relatively good in the Maple-basswood forest unit despite on-going disturbance due to human activity, deer, and non-native earthworms.

In many areas, the forest is devoid of groundcover and root crowns are exposed, indications of earthworm infestations. A review of storm water drainage should be conducted to mitigate erosion in the major ravines. Soil-rooted species of shrubs, forbs, grasses and sedges could help revegetate and stabilize the slopes. Ferns, such as interrupted, ostrich, or lady fern, should be planted in the ravine area. Although this is a typical species present in healthy ravine habitats, very few were spotted and most were isolated plants.

Human activity in the area needs to be curtailed. A trail plan is recommended to formalize the trail system by determining which trails to eliminate, re-route, or stabilize. Reducing and re-routing the trails will reduce continued damage and erosion of the bluff.

As with the other management units, management in this area should include common buckthorn and Tartarian honeysuckle removal.

The area of disturbed deciduous woodland in the northernmost portion of the project area is targeted for Maple-basswood forest restoration. Young sugar-maple and ironwood trees are common, indicating it may be developing into maple-basswood forest. It appears that a combination of factors, including bridge construction, erosion, the presence of invasive species and tree cutting have contributed to the disturbed nature of the woodland. To control erosion, water bars should be placed in some of the deeper ravines. Both common buckthorn and Tartarian honeysuckle are common on the slope. Buckthorn is locally abundant in areas of more intense disturbance. The exotic species should be removed and replaced with native trees and shrubs. The buckthorn should be cut and treated and tree species such as sugar maple, hackberry and ironwood, shrubs like alternate-leaved dog wood, arrow wood, and red-berried sumac should be planted in its place.

Human activity is quite evident in this area. A large amount of garbage has been dumped into the woodland from the top of the slope. Removing this trash at some point in the future would benefit the area's vegetation.

From the map of target plant communities (Map 4), it can be illustrated that some areas within the current oak forest land cover type are targeted for maple-basswood forest. The two principal locations where this occurs are on the cooler, north facing slopes of large ravines. While the north-facing slope is targeted for maple-basswood forest, the hotter and drier south-facing slope is best suited for oak forest.

Recommendations for management of the northern ravine include working to keep the buckthorn out of the area. Currently, there is not much buckthorn present. Monitoring the area will allow any new buckthorn and other invasives to be eliminated before they become established. More ferns and horsetails can be found in this location than in the ravines farther north, probably because the area is wetter and suffers from less foot traffic.

The north-facing slope of the southern most ravine, which is targeted for maple-basswood forest, has a relatively good ground cover composition. Very little honeysuckle and buckthorn are present. Management of this slope will involve removing these exotics and monitoring their regrowth.

The patch of land cover identified as 51-75% impervious cover with deciduous trees in this management unit can be planted to Maple-basswood Forest species. Any exotic species found here must be controlled before planting to natives takes place.

Oak Forest

Inventory Results: A detailed inventory of the Oak Forest land cover type and the areas targeted for Oak Forest restoration can be found in Appendix C in the sections titled "Oak forest," "Boxelder – green ash disturbed native forest (Northwest portion)," "Disturbed deciduous woodland (Southern portion)," and "Boxelder – green ash disturbed native forest (Southeast portion)."

Management Approach: Erosion is a serious problem in several portions of the oak forest unit. One large ravine is experiencing extensive erosion. Other smaller ravines are eroding to a lesser degree. A large slump is also present. The Ramsey Soil and Water Conservation District is aware of the ravine with severe erosion and has an interest in future stabilization efforts.

Very few sedges or ferns were observed near the ravines; these native species should be reintroduced and planted to help combat erosion and replace native ground layer species.

Human activity and trails contribute to erosion in many areas and also cause soil compaction and habitat fragmentation. This is particularly true with reference to the large ravine present near the Baker St. and Chippewa Ave. intersection. Closing or rerouting the trail leading to the ravine and the one that passes by the head of the ravine may help to curtail the damage in this area. A trail plan should be encouraged for the park to decide which trails can be closed, stabilized or re-routed.

Buckthorn removal has been taking place in the oak forest unit. The cut stump treatment appears to be very effective. However, crews will need to return to remove the occasional plants that were missed and include honeysuckle in the removal. Re-cutting and treating buckthorn and honeysuckle every couple of years will be necessary. This should occur when the buckthorn is either getting dense enough to shade out natives or starting to produce seed.

There are a couple of seeps present. These seeps occur just north of a large ravine. There is buckthorn present in the area around the seeps, but it appears to have been both cut and treated with herbicide. The seeps themselves are relatively free of buckthorn. Management in this area should thus focus on keeping the buckthorn out.

Some locations that are targeted for oak forest have much less buckthorn present than others, such as around the area following a major trail in the disturbed deciduous woodland (southern portion). These areas will need to be monitored to ensure that the invasive shrubs are removed at the early stages of colonization to prevent them from becoming established.

The current boxelder - green ash disturbed native forest in the south-eastern portion of the management area that contains a major ravine and is targeted for oak forest on the south-facing slope has a fair amount of buckthorn that needs to be cut. This especially needs to occur along the rim of the ravine. Some honeysuckle is also present and needs to be

removed. The south-facing slope is in much worse shape than the north-facing slope. It is bare and eroded; engineers should be contacted to figure out how to repair this before any activity can occur to re-vegetate the location.

Bottom Lands

Target Plant Communities: Lowland Hardwood Forest, Floodplain Forest, Willow Swamp, Mixed Emergent Marsh

Priority Status: 2

Floodplain

Inventory Results: A detailed inventory of the floodplain vegetation land cover can be found in Appendix C in the sections titled “Floodplain Forest,” “Willow Swamp” and “Mixed Emergent Swamp.”

Management Approach: These vegetation communities are regularly flooded. Consequently, vegetation management is difficult since soil and water conditions cannot be readily controlled. However, some management can be carried out including periodic debris removal, exotic brush removal, and planting tree and shrub species to augment and restore plant diversity. Timing plantings for periods after flooding in the spring will give the saplings a growing season to become established. Use as large a sapling as financially possible to increase the survival through floods. Debris will inevitably be deposited after each flood. Monitor the status of the area after flooding to assess the need for any clean up activities and survival of plantings. What little buckthorn is present can be removed by cutting and treating in the spring or fall with an aquatically approved herbicide such as Rodeo.

Lowland Hardwood Forest

Inventory Results: A detailed inventory of the Lowland Hardwood Forest land cover type and the areas targeted for Lowland Hardwood Forest restoration can be found in Appendix C in the sections titled “Lowland hardwood forest,” “Boxelder – green ash disturbed native forest (Northwest portion).” and “51-75% Impervious Cover with Deciduous Trees.”

Management Approach: The southwest end of the area targeted for lowland hardwood forest, dense stands of buckthorn have become established. A large amount of buckthorn was cut in the fall of the previous year, but the floor is dense with seedlings and resprouting stems. This area will need to be cut within the next two years. In order to remove the dense cover of buckthorn, a foliar spray may need to be used before native plants can be re-introduced into the area. In the spring or fall when desirable native vegetation is dormant, a broad spectrum herbicide (such as Roundup) or one that targets broad leaved plants (such as Garlon) may be used as a foliar spray on the still-green leaves of buckthorn.

Planting of tree, shrub and ground layer species in the spring of the year will augment existing native vegetation and increase plant diversity in the Lowland Hardwood Forest target areas.

Summary of management approaches

The table below summarizes the priorities for managing the units in Cherokee Park and shows where volunteers could be involved.

Table 5: Management Unit Priority Status, Management Actions and Volunteer Involvement

Cherokee Park Management Units	Priority Status	Management Action	Possible Volunteer Involvement
Lawn	1	Reconstruction of Oak Savanna along edge of bluff:	
		• Site Prep	
		• Seeding and planting of plugs	X
Side Slopes	1	Restoration of Oak and Maple-Basswood Forests	
		• Exotic brush removal	X
		• Planting plugs of grasses, forbs, and shrubs	X
		• Collecting seed of selected forest trees and shrubs and planting in specified areas	X
Bottom Lands	2	Restoration of Floodplain and Lowland Hardwood Forests	
		• Exotic brush removal	X
		• Planting plugs and seedlings of grasses, forbs, shrubs and trees	X
		• Collecting seed of selected forest trees and shrubs and planting in specified areas	X

Bibliography and References

Site Preparation, Seeding and Planting

- Armstrong, P.K. 1990. "Three No-Till Methods of Establishing Prairie on Small Sites (Illinois)." *Restoration and Management Notes* 8:33.
- Bronny, C. 1992. "Successional Restoration of an Oak Woodland (Illinois)." *Restoration and Management Notes* 10:77-78.
- Curtis, John T. 1959. *The Vegetation of Wisconsin: An Ordination of Plant Communities*. The University of Wisconsin Press: Madison.
- Kilde, Rebecca. *Going Native: A Prairie Restoration Handbook for Minnesota Landowners*. Minnesota Department of Natural Resources Section of Ecological Services Scientific and Natural Areas Program.
- Packard, Stephen and Cornelia F. Mutel, Society for Ecological Restoration. 1997. *The Tallgrass Restoration Handbook: for Prairies, Savannas, and Woodlands*. Washington DC: Island Press.
- Packard, Steve. 1993. "Restoring Oak Woodlands." *Restoration and Management Notes* 11:5-16.
- Sauer, L.J. 1998. *The Once and Future Forest*. Andropogon Associates, Ltd. Washington D.C.
- Shaw, Daniel. Native Vegetation in Restored and Created Wetlands, Its Establishment and Management in Minnesota and the Upper Midwest. 2000. Minnesota Board of Water and Soil Resources.
- Shirley, Shirley. 1994. *Restoring the Tallgrass Prairie: An Illustrated Manual for Iowa and the Upper Midwest*. Iowa City: University of Iowa Press.
- Thompson, Janette R. 1992. *Prairies, Forests, and Wetlands, the Restoration of Natural Landscape Communities in Iowa*. Iowa City: University of Iowa Press.

Long-term Management

- Bedker, P.J., J.G. O'Grien, and M.E. Mielke. *How to Prune Trees*. USDA Forest Service Pub. #NA-FR-01-95.
- Collins, S.L., and D.J. Gibson. 1990. *Fire in North American Tallgrass Prairies*. Norman: University of Oklahoma Press.
- Davis, Kenneth P. 1959. *Forest Fire: Control and Use*. McGraw-Hill Book Company, Inc.: New York.

- Howe, H.F. 1994. *Managing Species Diversity in Tallgrass Prairie: Assumptions and Implications*. Conservation Biology: 8(3):691-704.
- Ross, Laurel M., and Tom Vanderpoel. 1994. "Mowing Encourages Establishment of Prairie Species." *Restoration and Management Notes* 9(1):34-35.
- Tester, J.R.. 1989. *Effects of Fire Frequency on Oak Savanna in East-Central Minnesota*. Bulley of the Torrey Botanical Club. 116(7):134-144.
- Wisconsin Department of Natural Resources. 2000. *Ecologically Invasive Species*. www.dnr.state.wi.us/. Department of Natural Resources.
- Zins, M. and D. Brown. 1997. *Pruning Trees and Shrubs*. University of Minnesota Extension Service Fact Sheet FO-0628-B.

Invasive Species Control

- Henderson, Richard. 1982. *Vegetation - Fire Ecology of Tallgrass Prairie*. Natural Areas Journal 2(3)17-26.
- Kline, V. 1981. "Control of Honeysuckle and Buckthorn in Oak Forests. (Wisconsin)." *Restoration and Management Notes* 1:18.
- Minnesota Department of Agriculture, Agronomy and Plant Protection Division. 1998. *An MDA Pest Alert: Buckthorn and its Control*.
- Minnesota Department of Natural Resources Region V State Parks. 1995. *Fact Sheet #1: European (Common) Buckthorn*.
- Minnesota Department of Natural Resources Region V State Parks. 1997. *Problem Species Fact Sheet #2: Exotic Honeysuckles*.
- Packard, Stephen and Cornelia F. Mutel (editors). Society for Ecological Restoration. 1997. *The Tallgrass Restoration Handbook: For Prairies, Savannas, and Woodlands*. Washington DC: Island Press.
- Shirley, Shirley. 1994. *Restoring the Tallgrass Prairie: An Illustrated Manual for Iowa and the Upper Midwest*. Iowa City: University of Iowa Press.

The Ecological Classification System can be found at the Minnesota DNR website:
www.dnr.state.mn.us/ebm/ecs.

Great River Greening

Helping communities restore, manage and learn about their natural environment through volunteer involvement.




Figure 1.

The Challenge

Erosion, trash, and the invasion of exotic and invasive plant species are degrading our urban river valleys, reducing ecological diversity destroying wildlife habitat. Many public and private organizations are working to protect the river valleys, but these programs often lack long-term community involvement and stewardship.

These problems are especially pressing in the Twin Cities metropolitan region, home to more than 2 million people. The river valleys in this area:

- ☐ Hold some of the region's last intact native landscapes
- ☐ Serve as vital wildlife corridors for hundreds of migratory bird species
- ☐ Provide a water source for millions of the region's residents
- ☐ Contain some of the region's most scenic sites and vistas

Great River Greening's response

Great River Greening, a nonprofit organization, helps coordinate a cost-effective and sustained effort to manage ecosystems of the three great river valleys of the metropolitan area: the Mississippi, Minnesota and St. Croix. We are primarily an implementing organization, providing on-the-ground ecological restoration and management of both public and private land. We engage thousands of volunteers in the planting of native

vegetation, removal of exotic and invasive weeds, native-seed collection, and stewardship—work that cultivates an informed and involved citizenry. We also act as a catalyst, creating effective partnerships among agencies, municipalities, and private landowners responsible for managing river valleys and their natural resources. Restoration ecologists and other scientists provide technical expertise.

Key values

Great River Greening bases its work on these values:

1. Native trees and other vegetation have ecological and sociological value: They contribute to the health and biodiversity of ecosystems; they beautify surroundings; and they enhance a community's natural heritage and sense of place.
2. People want opportunities for direct involvement in natural resource protection and management, which help them feel connected and committed to their local natural areas.
3. Volunteer involvement in restoration and planning is one of the most effective methods of environmental education. When people work side by side to improve their environment, their communities become stronger and more vital.
4. Environmental restoration and stewardship require collaboration and inclusiveness.

We are committed to:

- ☐ Citizen-based restoration, stewardship and education
- ☐ Ecologically sound implementation and evaluation
- ☐ Collaboration to help advance ecosystem-based management
- ☐ Long-term stewardship.

Accomplishments—highlights

Since 1995, Great River Greening has involved more than 10,700 volunteers in the planting of 35,000 trees and shrubs and 16,000 wildflowers and grasses, as well as exotic-species removal, prairie-seed collection and broadcasting, plant inventories, training programs, and ongoing stewardship. In 2000 alone, we organized 30 events attended by nearly 1,500 volunteers!


We've also provided design and ecological consulting for numerous groups, including the city of Saint Paul Parks and Recreation Division, the Saint Paul Port Authority, the Science Museum of Minnesota, River Center, and the Greater Minnesota Housing Fund.

Great River Greening's major partners

City of Saint Paul • Friends of the Minnesota Valley • Friends of the Mississippi River • Metropolitan Council • Minneapolis Park and Recreation Board • Minnesota Department of Natural Resources • National Park Service • Ramsey County Parks and Recreation • Saint Paul Audubon Society • Trust for Public Land • U.S. Fish and Wildlife Service • Private landowners

To Contact Us

Great River Greening, 35 West Water Street, Suite 201, Saint Paul, MN 55107
651-665-9500 <http://www.greatrivergreening.org>













Cherokee Park Location in West Side Bluff Project Area


**Partners: West Side Citizen's Organization - Bluff Task Force and the City of St. Paul
Division of Parks and Recreation**

Map Created June 10, 2003, E. Fuge, Great River Greening
With Funding as Recommended by the Legislative Commission on Minnesota Resources

Legend

-  Project Boundary
-  Areas of Erosion
- Soil Type
 -  Dorerton-Rock Outcrop complex, 25 to 65 percent slopes
 -  Grays silt loam, 2 to 6 percent slopes
 -  Kerston muck, frequently flooded
 -  Pits, quarry
 -  Santiago silt loam, 2 to 6 percent slopes
 -  Udorthents, wet substratum
 -  Urban Land-Kingsley complex, 3 to 15 percent slopes
-  Contours

200 0 200 400 600 Feet


















Cherokee Park Soils, Slope and Erosion

Partners: West Side Citizens' Organization - Bluff Task Force
and the City of St. Paul Division of Parks and Recreation

Map created June 10, 2003, E. Fuge, Great River Greening

With Funding as Recommended by the Legislative Commission on Minnesota Resources

Legend

-  Management Units
-  Seep
- Existing Land Cover
 -  Buildings and pavement with 76-90% impervious cover
 -  51% to 75% impervious cover with deciduous trees
 -  Short grasses and mixed trees with 26-50% impervious cover
 -  Short grasses with sparse tree cover on upland soils
 -  Dry oak savanna sand-gravel subtype
 -  Oak forest
 -  Maple-basswood forest
 -  Boxelder - green ash disturbed native forest
 -  Disturbed deciduous woodland
 -  Lowland hardwood forest
 -  Floodplain forest
 -  Willow swamp
 -  Mixed emergent marsh

200 0 200 400 600 Feet











Cherokee Park Existing Land Cover

Partners: West Side Citizens' Organization - Bluff Task Force
and the City of St. Paul Division of Parks and Recreation


Map created June 10, 2003, E. Fuge, Great River Greening

With Funding as Recommended by the Legislative Commission on Minnesota Resources

Legend

-  Management Units
-  Target Plant Communities
-  Oak Savanna
-  Oak Forest
-  Maple-Basswood Forest
-  Lowland Hardwood Forest
-  Floodplain Forest
-  Lawn or Developed Activity Areas

200 0 200 400 600 Feet



Cherokee Park Management Units and Target Plant Communities

Partners: West Side Citizens' Organization - Bluff Task Force and the City of St. Paul Division of Parks and Recreation


Map created June 10, 2003, E. Fuge, Great River Greening

With Funding as Recommended by the Legislative Commission on Minnesota Resources

Legend

- Project Boundary
- Baseball diamond
- Benches
- Fire hydrants
- Fire Pit
- Gates
- Lamp posts
- Manholes
- Power poles
- Signs
- Picnic and restroom area
- Tennis courts
- Fences
- Sidewalks
- Trails

200 0 200 400 600 Feet



Cherokee Park Cultural Features

Partners: West Side Citizens' Organization - Bluff Task Force and the City of St. Paul
Division of Parks and Recreation

Map created June 10, 2003, E. Fuge, Great River Greening

With Funding as Recommended by the Legislative Commission on Minnesota Resources

Appendix A: Fact Sheets for Exotic and Invasive Plants

The following pages contain information on the habitat, phenology, and niche of exotic and invasive plants found or potentially found in the Cherokee Park plant communities. These fact sheets pertain to troublesome plants that compete with the native plants typical of undisturbed native communities and threaten the integrity, structure and function of those communities. Active management to control invasive plant species is essential to restoring the health of plant communities and the habitats they provide for a diverse group of native animals. Additional fact sheets for common buckthorn, box elder, Tartarian honeysuckle, Siberian elm, staghorn sumac, burdock, leafy spurge, garlic mustard, and poison ivy can be found in Appendix B of the West Side Bluff Ecological Inventory and Vegetation Management Plan.

Forbs:

Canada thistle
Purple loosestrife
Sweet clovers

Cirsium arvense
Lithrum salicaria
Melilotus officinalis
M. alba

Grasses:

Bluegrass
Reed canary grass
Smooth brome grass

Poa pratensis, *P. compressa*
Phalaris arundinacea
Bromus inermis

Effective management of these species, which are present or potential problems in Cherokee Park, is described in the following fact sheets. Except for the reed canary grass and purple loosestrife, wetland plants, most of the invasive plants are found in and threaten the woodlands.

Buckthorn is generally established throughout the woodlands of the park with some areas of heavier concentrations. Much of the buckthorn was cut and treated in the fall of 2002. Seedlings are still prevalent and resprouts will occur. Consequently, continued treatment is recommended initially. In the future, periodic surveys and localized cutting may only be required every two years or so.

Garlic mustard was not found in Cherokee Park in 2003, but is a potential invader to watch for, especially in the wooded areas. Invasive exotics such as burdock, bluegrass, European brome, Canada thistle and the sweet clovers are nuisances in young prairie and savanna restorations.

Canada Thistle (*Cicium arvense*)



Photo by Merel R. Black

Effects of Invasion:

Canada thistle is an alien species capable of crowding out and replacing native grasses and forbs. It is detrimental to natural areas where it occurs, particularly non-forested communities, and it can change the natural structure and species composition where it becomes well established. Prairies, barrens, savannas, and glades are susceptible, particularly those sites that have been disturbed as well as those undergoing manipulative restoration management. It is important to control this species prior to restoration work.

The plant grows in clonal patches of all female or male plants. As a result, some patches produce seeds and others do not. Seeds mature quickly and are capable of germinating within 8 to 10 days after the flowers open, even if the plants are cut when flowering. Most seeds germinate within one year, but may remain viable in the soil for up to 20 years. Seeds are mostly dispersed by wind and sometimes by water runoff. Small sections of broken roots are capable of producing new plants.

Canada thistle is considered a noxious weed under Minnesota law and should not be allowed to go to seed.

Size: Canada thistle is a 2 to 5 foot (0.6 to 1.5 meters) tall herbaceous plant with deep, wide spreading, horizontal roots. The root system is usually within a foot of the surface, but may extend 6 feet deep or more in loose soil. The horizontal roots stemming from the fibrous taproot of a single plant can spread 10 to 12 feet in one season, resulting in a circular infestation 20 feet across. Aerial shoots are sent up in 2 to 6 inch intervals, and generally produce basal leaves the first year and flowering stems the next year.

Habit: Canada thistle is a clone-forming perennial. The grooved, slender stems branch only at the top and are slightly hairy when young; becoming covered with hair as the plant grows.

Leaves: The oblong, tapering, sessile leaves are deeply divided, with prickly margins. Leaves are green on both sides with a smooth or slightly downy lower surface.

Fruit: Seeds are small (3/16 inch or 0.5 cm long), light brown, smooth and slightly tapered, with a tuft of tan hair loosely attached to the tip.

Flowers: Numerous small, compact (3/4 inch or 1.9 cm. diameter), rose-purple or white flowers appear on upper stems from June to September.

Origin: Canada thistle is native to Europe, not Canada, as its name suggests. Its current range encompasses the northern portion of the United States east of the Rocky Mountains.

Mechanical Control:

Repeated pulling, routine mowing or selective cutting will eventually starve underground stems and effectively reduce an infestation within 3 or 4 years. The ideal time to cut is in the very early bud stage when food reserves are at their lowest point. Plants cut 8 days or more after flowers have opened should be removed from the site because seeds mature quickly. Cutting should be completed prior to flowering and seed set. If seeds are ripe, cut flower heads must be removed from the site immediately to avoid further seed dispersal. Plants should be pulled or cut at least three times during the growing season -- for example, in June, August, and September. Some persons have had success killing individual plants by cutting the top and putting table salt down the hollow stem.

Prescribed fire can be effective in controlling this species and is a preferred treatment. Late spring burns between May and June, effectively discourage this species, whereas early spring burns can increase sprouting and reproduction. During the first 3 years of control efforts, burns should be conducted annually. Healthy, dense prairie vegetation can produce enough competition to reduce the abundance of Canada thistle.

On severely disturbed sites with heavy infestations, such as cropland or abandoned cropland, the site could be plowed and sowed to a cover crop (wheat, alfalfa, and rye), if practical and desirable. The following May, the cover crop should be plowed under and desired native species should be seeded. Tillage disturbance of soil may provide ideal conditions for reinvasion and for introduction of other exotics. Grazing is not an effective control measure as the prickles prevent livestock from grazing near Canada thistle.

Chemical Control:

Control of this species with herbicides in natural areas is not recommended, as the herbicide can damage native vegetation more than the damage caused by the thistle. However, spot application of the amine formulation of 2,4-D using a wick applicator or hand sprayer can control individual stems if necessary. Infested lands that are not considered high quality natural areas may be controlled using a foliar application of a 1-2% active ingredient solution of glyphosate in spring when plants are 6-10 inches tall.

Spot application of Transline (a formulation of clopyralid), according to label instructions can control this plant. Individual plants of Canada thistle should be treated with a wick applicator or hand sprayer. The herbicide Transline is selective for broadleaf plants. To reduce vapor drift and improve plant up-take of the chemical, a surfactant may be added to the spray solution. Precautions should be taken to avoid contacting nontarget plants with the solution.


A foliar application of a 1-2% solution of Roundup (a formulation of glyphosate) applied in spring when plants are 6-10 inches (15.2 -25.4 cm) tall is an effective herbicide treatment. Individual plants should be spot-treated with a wick applicator. Roundup normally kills the entire plant, including the roots, when applied in this manner. Roundup is a nonselective herbicide and precautions should be taken to avoid contacting nontarget plants with the solution.

Sources:

Wisconsin Department of Natural Resources, 2002

Vegetation Management Manual, Vol. 1, No. 2. Illinois Nature Preserves Commission, approved 02/06/90

Purple Loosestrife (*Lithrum Salicaria*)



Effects of Invasion

Purple loosestrife spreads mainly by seed, but it can also spread from roots or stems. A single stalk can produce 100,000–300,000 seeds per year. Sunny and partly shaded wetland is susceptible to invasion. Purple loosestrife generally builds up a large seed bank in the soil for several years before becoming dominant. After disturbance, loosestrife can spread rapidly, eventually taking over entire wetlands. Purple loosestrife degrades wetlands by displacing native wetland vegetation and decreasing habitat for wildlife species.

Habit: Purple loosestrife is a perennial herb 3–7 feet tall with a dense bushy growth of 1–50 stems.

Size: 3–7 feet tall.

Leaves: Leaves are opposite, nearly linear, and attached to 4-sided stems without stalks.

Stem: Stems range from green to purple.

Flower: Flowers vary from purple to magenta, have 5–6 petals and are aggregated into numerous long spikes. Flowering occurs from July to September.

Origin: Europe.

Mechanical Control

Small young plants can be hand pulled while older plants can be removed with a shovel. If possible, entire root systems should be removed to prevent re-sprouting. Soil disturbance should be minimized to prevent seedling establishment. Plants should be controlled before the onset of seeds around the first week of August or seeds should be cut and bagged. Plant parts should be dried and disposed of accordingly. Follow-up treatments are recommended for at least 3 years after removal. Mowing and burning have not been effective with purple loosestrife. However, water-level manipulation has been successful. Water levels are reduced until loosestrife has sprouted, then levels are increased until stems are drowned.

Biological Control

Biocontrol is currently considered the most viable option for purple loosestrife control. Several natural insect enemies of purple loosestrife from Europe have been introduced. A species of weevil (*Hylobius transversovittatus*) lays eggs in the stem and upper root system of the plant and its larvae eat root tissue. In

addition, two species of leaf-eating beetles (*Galerucella californiensis* and *G. pusilla*) and a weevil that feeds on flowers (*Nanophyes marmoratus*) are being used. These insects almost exclusively feed on *Lythrum salicaria* and not native plants. The insects generally do not eradicate loosestrife but reduce the population to a state where it does not dominate native habitats.

Chemical Control

Glyphosate is the most common chemical used for killing purple loosestrife. The formula designed for use on wet or standing water sites should be applied in late July or August. A 1% active ingredient (a.i.) solution should be used, and only 25% of the foliage of each plant needs to be covered. Glyphosate mixed to 3%–10% solution can also be used on freshly cut stems (this is effective on larger plants in areas of low loosestrife densities). Cut stems should be removed from the site and disposed of appropriately. Triclopyr formulated for water dilution is an effective herbicide for loosestrife. This broadleaf herbicide does not harm sedges or monocots. Foliar application should cover nearly all of the foliage.

Source: Wisconsin Department of Natural Resources, 1997.

Yellow Sweet Clover (*Melilotus officinalis*)
White Sweet Clover (*Melilotus alba*)



Photo by John M. Randall, TNC

Effects of Invasion:

Sweet clovers are fire-influenced, aggressive, weedy plants that produce populations with high rates of fluctuation. Both species degrade native grasslands by overtopping and shading native sun-loving species. Sweet clovers are members of the legume family.

Both white and yellow sweet clovers are biennials. After germination in late spring or summer, the plants put their energy into developing a healthy root system. Plants are strictly vegetative in the first year and have a small, branched stem with clover-like leaves. First-year plants can be found in late summer. In the second year, plants may be seen in late April or early May. By that time, individuals have a strong taproot and a root crown from which new shoots appear. Plant height is dependent on root development and growing conditions; healthier plants are taller. Sweet clovers flower from late May through September, set seed, and die. Both plants produce small, hardy seeds that remain viable in the soil for as many as thirty years.

Burning produces excellent growing conditions for clover by scarifying seeds and stimulating germination. During the next year following a burn, many flowering plants generally emerge.

Size: In the second year, plants may appear bushy, and grow from three to six feet in height.

Habit: First year seedlings are leafy, green, few-stemmed and around a foot tall. Second year plants generally have three main stout stems arising from the root crown. The 3 – 6 foot plants are conical and bushy.

Leaves: Leaves are alternate, divided into three finely toothed leaflets, with the middle leaflet occurring on a distinct stalk.

Fruit: The legume is ovoid, leathery and wrinkled, longer than the calyx, and scarcely dehiscent, with one or two small seeds.

Flower: Yellow and white sweet clovers appear very similar except for the distinguishing yellow or white flowers. Yellow sweet clover is usually smaller than white sweet clover and blooms earlier. The flowers are packed densely on the top four inches of an elongated stem. Each small flower is attached to the stem by a minute stalk.

Origin: Sweet clovers are native to Europe and Asia. They were brought to North America in the late 1600's as an agricultural crop for forage and honey production. These clovers also fix nitrogen, and thus became popular as soil enhancers. The chemical used in the production of the blood thinner Warfarin was first discovered in sweet clover. Due to the economic values of white and yellow sweet clover, these species will continue to be planted despite the problems they pose for land managers. Both species are found in all fifty states, although they are most frequently found in the states of the Upper Midwest and Great Plains. Sweet clovers grow well in direct sunlight or in partial shade. Neither species can tolerate complete shade. Sweet clovers seem to prefer calcareous or loamy soils, and are most frequently found in open, disturbed, upland habitats such as prairies, savannas, and dunes.

Mechanical Control

On grasslands managed with prescribed burning, it is possible to greatly reduce sweet clover by burning two years in a row. Burning should be done early the first year (before green-up--usually in early to mid-April) to stimulate germination. The burned area should be checked in late summer for first year plants. If plants are found, another burn should be conducted the next year in early to mid May. If burning is conducted before the buds are developed, the plants will resprout. Heavily infested areas may need this burning sequence repeated after a few years. The fire may be of low intensity--just enough to touch the stems. Damaged plants wither quickly if they are not completely destroyed by fire. For small patches or those areas not completely burned, a flame gun (torch) may be used when the vegetation is damp to avoid burning surrounding prairie. Another burning strategy is to mow later in the summer, allow the cut plants to dry, and then burn. This can be stressful to the native vegetation and should not be done annually.

Small amounts of sweet clover can be controlled by hand-pulling in late fall after first-year plant root-crown buds have developed, or in May or June before second-year plants flower. Pulling is easier when the soil is wet. Plants can also be cut at ground level with brush loppers. If pulling is tried too early, many plants may be missed, and those with succulent stems may break off and resprout. But pulling must be done before seeds are set; otherwise cut plants will have to be removed from the natural area. It is necessary to inspect the area a couple of times in summer for late flowering plants.

For very dense small patches, cutting with a power brush-cutter using a heavy duty saw blade is effective. The stand should be cut just before flowering, and checked a week later for individuals missed or partly cut.

It is necessary to conduct annual inspections to remove scattered individual plants. Disturbed areas such as fox dens provide habitats that can allow sweet clover to greatly increase over time if not controlled. Habitats adjacent to managed areas should also be inspected to reduce sweet clover invasion on managed sites. Due to the long viability of sweet clover seeds (up to 30 years) and continued agricultural use, these plants generally must be managed on a continuous basis.

Chemical Control

Sweet clover can be managed using mechanical controls, and should not require chemical use.

Source: Wisconsin Department of Natural Resources, 2002; The Nature Conservancy, 2002

Kentucky Bluegrass (*Poa pratensis*)
Canada Bluegrass (*Poa compressa*)



(c) John M. Randall/The Nature Conservancy

Effects of invasion:

Because bluegrass grows early in the season (when most other species are still dormant), it can spread very quickly. However, its shallow root system makes it susceptible to high soil temperatures and low soil moisture. Bluegrass has successfully invaded both remnant and restored prairies, savannas, and barrens. Establishment can be attributed to intentional introduction, past mowing, grazing, or cessation of fire. If left unattended, bluegrass can out-compete native prairie grasses and forbs, and will dominate shaded areas resulting from woody species invasions.

Description: Most of the cool season grasses that begin growing early are not native to Wisconsin prairies. Bluegrass can be distinguished vegetatively from other early grasses by its narrow blade, which is V-shaped in cross section, and by the leaf tip, which is shaped like the bow of a boat. Kentucky bluegrass is distinguished from Canada bluegrass by the shape of the stem. In Kentucky bluegrass the stem is round; Canada bluegrass has a flat stem. Their effects on the natural systems are equivalent and therefore should be treated as one problem. Many of the other cool-season European grasses (brome, timothy, orchard grass, quack grass, etc.) have similar growth habits and can be controlled using the techniques discussed below.

Distribution and habitat: Kentucky bluegrass was introduced as a cultivar from Europe, and has been bred into multiple cultivars since its introduction. Because of its extensive use for lawns and in pastures, it is common in most grasslands, even those managed for native species. Canada bluegrass is also naturalized from Europe. Kentucky bluegrass is a common lawn and pasture grass. Canada bluegrass is often mistaken for Kentucky bluegrass, but is distinguished by forming extensive sods in dry, sterile soils (especially acidic soils) that cannot sustain the more common Kentucky bluegrass. Kentucky bluegrass is usually found on more mesic and fertile soils, although it will grow on dry neutral or alkaline soils.

Mechanical Control

A controlled fire can dramatically reduce bluegrass in a native or planted prairie, savanna, or barrens. Fire will also set back the woody species whose shade encourages the proliferation of cool-season grasses. In southern Wisconsin, a late April or early May burn will destroy three to eight inches of new growth. Timing of burns may change on a year-to-year basis depending on weather conditions. Observing bluegrass growth is essential for effective control by burning. Fire is most effective when bluegrass is three to eight inches high. Burning at this time kills new growth and removes accumulated leaf litter. Burning off the moisture-retaining blanket of leaf litter increases stress on the shallow-rooted bluegrass by exposing the

darkened surface to the sun. This helps reduce the competitive ability of bluegrass by encouraging summer dormancy and decreasing the chance of flowering and seed production. The effect is most pronounced on dry prairies and barrens. Burning can reduce bluegrass by more than 90%, but it is rarely 100% effective. Burning at the right time also improves the competitive advantage of native, warm-season grasses and forbs. Native species emerge later and benefit from the elimination of duff and a darkened soil surface.

When converting areas dominated by cool-season grasses into prairie, it is helpful to reduce the grass cover and seed bank before planting native seeds. This can be accomplished by any combination of tilling, smothering the grass, or applying herbicide. Till several times a year for at least one season to expose the seed bank and prevent further growth of the grass sod. Herbicide use followed by a season of tilling is also effective. On small sites, grasses can be killed by covering with black plastic or layers of newspapers during the growing season.

Chemical Control

Herbicide use is not recommended to control bluegrass on grasslands or savannas where there are native prairie plants. However, herbicide may be required on severely degraded areas or where prairie restoration is beginning. In such cases, the herbicide glyphosate has proven effective when used according to label applications.

Source: Wisconsin Department of Natural Resources, 2002

Reed Canary Grass (*Phalaris arundinacea*)



Effects of Invasion:

Reed canary grass reproduces by seed or creeping rhizomes and spreads aggressively. It prefers disturbed areas but can easily move into native wetlands. In less than 12 years, reed canary grass can form large, monotypic stands that harbor few other plant species and therefore are of little use to wildlife. Reed canary grass dominates an area by building up a tremendous seed bank that can eventually erupt, germinate, and recolonize treated areas. Reed canary grass is difficult to eradicate; no single control method is universally applicable.

Size: 2–9 feet in height.

Habit: A large, coarse, cool-season, sod-forming, perennial wetland grass. Sprouts early in spring, forming a thick rhizome system that dominates the subsurface soil.

Blades: Erect, hairless stem with gradually tapering leaf blades 3.5–10 inches long and .25–.75 inches wide. The ligule is highly transparent.

Panicles: Compact, erect or slightly spreading (depending on the plant's reproductive stage), ranging from 3–16 inches long with branches .5–1.5 inches long.

Flowers: Single flowers occur in dense clusters in May to mid-June. They are green to purple, changing to beige over time.

Seeds: Shiny brown.

Origin: Eurasia and North America.

Mechanical Control

- Small, discrete patches may be covered by black plastic for at least one growing season then seeded with native species. This method is not always effective and must be monitored because rhizomes can spread beyond the edge of the plastic.

- Prescribed burns in late spring or late fall may help reduce the population if repeated annually for 5–6 years. The application of 1.5% glyphosate solution will “brown off” reed canary grass enough to conduct burns. A late spring burn followed by mowing or wick application of glyphosate to the emerging flowering shoots will eliminate seed production for that year. Burning is ineffective in eliminating dense stands of reed canary grass that lack competition from native, fire-adapted species in the seed bank.
- Mowing twice yearly (early to mid-June and early October) may help control reed canary grass by removing seed heads before the seed matures and by exposing the ground to light, which promotes the growth of native wetland species. Discing the soil in combination with a mowing or burning regimen may help by opening the soil to other species.
- Hand-pulling or digging may work on small stands in the early stages of invasion.
- A bulldozer can be used to remove reed canary grass and rhizomes (12–18 inches deep), after which native species should be seeded. Discing or plowing can also be used in this way.
- Repeated cultivation for one full growing season followed by dormant seeding near the first-frost date. Combine with spot herbicide application in sections too wet for early or late cultivation.

Chemical Control

Cut and spray

- Tie the stems of small clones together just before they flower, then cut them and apply glyphosate in a 33% solution to the cut stems.
- Perform foliar application of a 5% glyphosate solution designed for use in wetlands in early spring when most native species are dormant to the foliage. Remove the dead leaves from the previous year before applying herbicide. Two herbicidal applications may be necessary to ensure complete coverage. Mow in mid-September then apply herbicide in October (after big bluestem is dormant).
- Perform wick application of a 5% glyphosate solution designed for use in wetlands in the first to third weeks of June, followed by a late June to mid-July burn. This technique reduces reed canary grass cover, depletes the seed bank, and stimulates native seed banks.
- In non-aquatic environments, apply Dalpon and trichloroacetic in late fall or early winter at a rate of 20lbs.–40 lbs./acre on dried foliage.

Source: Wisconsin Department of Natural Resources, 1997.
Minnesota Department of Natural Resources, 1995.

Smooth (Awnless) Brome (*Bromus inermis*)



Seed head



Field of brome

Photos: Minnesota DNR-Angela Anderson

Effects of Invasion: Smooth brome is a cool season exotic that is especially troublesome in disturbed portions of native plant communities and restorations in the tallgrass and mixed prairie regions. Although less invasive than Kentucky bluegrass, with which it often occurs and is managed, it is also less responsive to management. Smooth brome has been widely planted as a forage and cover crop. Although perhaps not as invasive as *Poa pratensis*, with which it often grows, it is highly persistent. It forms a dense sod that often appears to exclude other species, thus contributing to the reduction of species diversity in natural areas.

Size: *Bromus inermis* is a perennial cool season grass that grows 2 - 3' high with a hairless erect stem. Brome roots have been known to reach a depth of 4.7 feet.

Habit: *Bromus inermis* is a deeply rooting, rhizomatous, sod-forming perennial grass. The drought resistance of smooth brome is probably accounted for in part by its deeply penetrating root system. The heavy concentration of total root mass near the surface is the result of smooth brome's creeping rhizomatous habit. Old brome fields develop a "sod bound" condition in which shoot density is reduced and symptoms of nitrogen deficiency are exhibited. Because of its fairly distinctive foliage and habit of growing in solid patches *Bromus inermis* is easily recognized at all seasons. Its early green-up makes it especially easy to detect during the spring months.

Leaves: The leaf blades are smooth, flat, 4-5 inches long and 1/4-3/8 inches wide with a conspicuous "M"- or "W"-shaped constriction in the middle.

Fruit: Lemmas are all unawned or with very short awn.

Flowers: The inflorescence is an erect, open panicle with ascending branches that are sometimes reflexed, blooming May - July.

Origin: *Bromus inermis* is a Eurasian species ranging from France to Siberia, apparently introduced in the United States by the California Experiment Station in 1884. Within the United States smooth brome has been introduced in the northeastern and northern Great Plains states as far south as Tennessee, New Mexico and California. It has become naturalized from the maritime provinces to the Pacific coast north to Alaska to California and through the plains states. Within the United States, "northern" and "southern" agricultural strains have been developed. The southern strain is more tolerant of drought and heat than the northern strain.

Mechanical Control

Both experimental studies and management experience indicate that burning or cutting smooth brome in the boot stage is perhaps the most effective means of control. Smooth brome is in boot stage between mid-

April and late May when the plant has reached a height of 18 to 24 inches and the flowering head is still enclosed within the sheath. This is somewhat later than would be recommended for other management purposes such as control of Kentucky bluegrass. Research indicates that a well-timed burn that treats *Bromus inermis* in boot or early flower may be more effective than mowing at the same susceptible period. It appears that late May burns would be optimal in the northern plains for reduction of smooth brome. One close mowing when the plants are 18-24 inches tall (followed ideally by 3 repetitions), may improve chances of selectively controlling this species. The best conditions for damage are hot, moist weather at the time of cutting, followed by a dry period.

Chemical Control

Its habit of occurring frequently in nearly pure swards renders *Bromus inermis* a good target for selective control by timed, close mowing or use of herbicides. An early study of brome control found Tordon (picloram) most effective at rates of 1.1 to 2.2 kg/ha, or treatment with Roundup (glyphosate) at 0.5 to 1.1 kg/ha before flowering. It appears that April or May applications of glyphosate at 2 kg/ha may be an effective management technique for controlling smooth brome in pure patches.

Sources:

NatureServe. 2003. NatureServe Explorer: An online encyclopedia of life [web application]. Version 1.8. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>. (Accessed: July 14, 2003).

Element Stewardship Abstract for *Bromus inermis*, The Nature Conservancy, 1987 (updated May 2000)

Minnesota invasive non-native terrestrial plants, an identification guide for resource managers, MN DNR, 2003

Appendix B: Species Lists for Proposed Restoration Target Communities

The following species lists contain the common plants of intact remnant communities in Minnesota. These species lists have been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

- Dry prairie species list (Appendix B Table 1)
- Oak savanna species list (Appendix B Table 2)
- Maple-basswood forest species list (Appendix B Table 3)
- Dry oak forest species list (Appendix B Table 4)
- Lowland hardwood forest species list (Appendix B Table 5)
- Floodplain forest species list (Appendix B Table 6)
- Mixed emergent marsh species list (Appendix B Table 7)
- Willow swamp species list (Appendix B Table 8)

Appendix B Table 1. Dry Prairie species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Shrubs	
<i>Amorpha canescens</i>	Lead-plant
<i>Prunus americana</i>	American plum
<i>Prunus virginiana</i>	Choke cherry
<i>Rhus glabra</i>	Smooth sumac
<i>Rosa arkansana</i>	Prairie rose
<i>Symphoricarpos occidentalis</i>	Wolfberry

Forbs	
<i>Anemone cylindrica</i>	Long-fruited Thimbleweed
<i>Antennaria plantaginifolia</i>	Large-leaved pussytoes
<i>Apocynum cannabinum</i>	Indian Hemp
<i>Apocynum sibiricum</i>	Clasping Indian Hemp
<i>Artemisia ludoviciana</i>	White Sage
<i>Asclepias syriaca</i>	Common Milkweed
<i>Asclepias tuberosa</i>	Butterfly Milkweed
<i>Asclepias verticillata</i>	Whorled Milkweed
<i>Asclepias viridiflora</i>	Green milkweed
<i>Aster ericoides</i>	Heath Aster
<i>Aster oolentangiensis</i>	Sky-blue aster
<i>Campanula rotundifolia</i>	Harebell
<i>Comandra umbellata</i>	Bastard toadflax
<i>Coreopsis palmata</i>	Stiff tickseed or bird foot coreopsis
<i>Dalea purpureum</i>	Purple prairie clover
<i>Delphinium virescens</i>	Prairie larkspur

<i>Euphorbia corollata</i>	Flowering spurge
<i>Gnaphalium obtusifolium</i>	Sweet everlasting
<i>Hedera hispida</i>	Mock pennyroyal
<i>Helianthemum bicknellii</i>	Hoary frostweed
<i>Helianthus pauciflorus (rigidus)</i>	Stiff sunflower
<i>Heuchera richardsonii</i>	Alum root
<i>Lespedeza capitata</i>	Round-headed bush-clover
<i>Liatris aspera</i>	Rough blazing star
<i>Liatris punctata</i>	Dotted blazing star
<i>Lithospermum canescens</i>	Hoary puccoon
<i>Lithospermum incisum</i>	Narrow-leaved puccoon
<i>Physalis heterophylla</i>	Ground-cherry
<i>Potentilla arguta</i>	Prairie cinquefoil
<i>Rosa arkansana</i>	Prairie rose
<i>Solidago nemoralis</i>	Gray goldenrod
<i>Solidago ptarmicoides</i>	White aster
<i>Solidago rigida</i>	Stiff goldenrod
<i>Tradescantia occidentalis</i>	Western spiderwort
<i>Viola palmata</i> var. <i>pedatifida</i>	Prairie Violet

Grasses and Sedges	
<i>Andropogon gerardii</i>	Big bluestem
<i>Bouteloua gracilis</i>	Blue grama grass
<i>Bouteloua hirsuta</i>	Hairy grama grass
<i>Bouteloua curtipendula</i>	Side-oats grama
<i>Carex heliophila</i>	A species of sedge
<i>Cyperus lupulinus</i>	Hop-like-cyperus
<i>Elymus canadensis</i>	Canada wild rye
<i>Koeleria macrantha</i>	Junegrass
<i>Panicum oligosanthos</i>	Scribner's panic grass
<i>Muhlenbergia cuspidata</i>	Plains muhley
<i>Schizachyrium scoparium</i>	Little Bluestem
<i>Sorghastrum nutans</i>	Indian Grass
<i>Sporobolus heterolepis</i>	Prairie dropseed
<i>Stipa spartea</i>	Porcupine grass

Appendix B Table 2. Oak Savanna species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees	
<i>Quercus macrocarpa</i>	Bur oak
<i>Quercus ellipsoidalis</i>	Northern pin oak
<i>Populus tremuloides</i>	Quaking aspen
<i>Prunus serotina</i>	Black cherry

Shrubs	
<i>Amorpha canescens</i>	Leadplant
<i>Ceanothus americanus</i>	New Jersey tea
<i>Cornus racemosa</i>	Gray dogwood
<i>Corylus americana</i>	American hazelnut
<i>Rhus glabra</i>	Smooth sumac
<i>Rosa arkansana</i>	Prairie rose
<i>Prunus virginiana</i>	Chokecherry
<i>Amelanchier sanguinea</i>	Round-leaf serviceberry
<i>Symphoricarpos occidentalis</i>	Wolfberry
<i>Salix humilis</i>	Prairie willow

Forbs:	
<i>Anemone cylindrica</i>	Thimbleweed
<i>Antennaria neglecta</i>	Pussytoes, white
<i>Antennaria plantaginifolia</i>	Plantain-leaved or large-leaved pussytoes
<i>Aristida tuberculosa</i>	Butterfly weed
<i>Artemisia ludoviciana</i>	Prairie sage
<i>Artemisia frigida</i>	Prairie sagewort
<i>Asclepias tuberosa</i>	Butterfly milkweed
<i>Asclepias verticillata</i>	Whorled milkweed
<i>Asclepias viridiflora</i>	Green milkweed
<i>Aster ericoides</i>	Heath aster
<i>Aster oolentangiensis</i>	Azure aster
<i>Aster sericeus</i>	Silky aster
<i>Astragalus crassicaarpus</i>	Buffalo-bean, ground-plum
<i>Besseyia bullii</i>	Kitten-tails
<i>Calylophus serrulata</i>	Toothed-leaved evening primrose
<i>Campanula rotundifolia</i>	Harebell
<i>Coreopsis palmata</i>	Stiff tickseed or bird-foot coreopsis
<i>Dalea candidum</i>	White prairie clover
<i>Dalea purpureum</i>	Purple prairie clover
<i>Delphinium virescens</i>	Prairie larkspur
<i>Desmodium illinoense</i>	Illinois tick-trefoil
<i>Euphorbia corollata</i>	Flowering spurge
<i>Fragaria virginiana</i>	Wild strawberry
<i>Galium boreale</i>	Northern bedstraw
<i>Geum triflorum</i>	Prairie smoke

<i>Gnaphalium obtusifolium</i>	Sweet everlasting
<i>Helianthemum bicknellii</i>	Hoary frostweed
<i>Helianthus hirsutus</i>	Woodland sunflower
<i>Helianthus occidentalis</i>	Western sunflower
<i>Helianthus rigidus</i>	Rigid sunflower
<i>Heliopsis helianthoides</i>	Early sunflower
<i>Heterotheca villosa</i>	Hairy golden aster
<i>Heuchera richardsonii</i>	Alum root
<i>Hieracium longipilum</i>	Long-bearded hawkweed
<i>Lespedeza capitata</i>	Round-headed bush-clover
<i>Liatris aspera</i>	Rough blazing star
<i>Liatris punctata</i>	Dotted blazing star
<i>Lithospermum canescens</i>	Hoary puccoon
<i>Lithospermum caroliniense croceum</i>	Hairy puccoon
<i>Lobelia spicata</i>	Rough-spiked lobelia
<i>Monarda fistulosa</i>	Wild bergamot
<i>Oenothera biennis</i>	Evening primrose
<i>Oxalis violacea</i>	Violet wood sorrel
<i>Penstemon gracilis</i>	Slender beard-tongue
<i>Penstemon grandiflorus</i>	Large-flowered beard-tongue
<i>Physalis virginiana</i>	Ground cherry
<i>Rudbeckia hirta pulcherrima</i>	Black-eyed Susan
<i>Sisyrinchium campestre</i>	Blue-eyed grass
<i>Smilacina stellata</i>	Starry false Solomon's seal
<i>Solidago nemoralis</i>	Gray goldenrod
<i>Solidago ptarmicoides</i>	White aster
<i>Solidago rigida</i>	Stiff goldenrod
<i>Teucrium canadense</i>	Germander
<i>Tradescantia occidentalis</i>	Western spiderwort
<i>Verbena stricta</i>	Hoary vervain
<i>Viola pedatifida</i>	Prairie violet

Grasses and Sedges	
<i>Andropogon gerardii</i>	Big bluestem
<i>Aristida basiramea</i>	Three-awn grass
<i>Bouteloua curtipendula</i>	Side-oats grama
<i>Bouteloua hirsuta</i>	Hairy grama
<i>Carex muhlenbergii</i>	Muhlenberg's sedge
<i>Carex pennsylvanica</i>	Pennsylvania sedge
<i>Elymus canadensis</i>	Canada wild rye, nodding wild-rye
<i>Koeleria macrantha</i>	June grass
<i>Muhlenbergia cuspidata</i>	Plains muhley
<i>Panicum oligosanthos</i>	Scribner's panic grass
<i>Panicum virgatum</i>	Switch grass
<i>Schizachyrium scoparium</i>	Little bluestem
<i>Sorghastrum nutans</i>	Indian grass
<i>Sporobolus heterolepis</i>	Prairie dropseed
<i>Stipa comata</i>	Needle grass
<i>Stipa spartea</i>	Porcupine grass

Appendix B Table 3. Maple-basswood Forest species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees - Canopy	
<i>Acer saccharum</i>	Sugar maple
<i>Celtis occidentalis</i>	Hackberry
<i>Juglans cinerea</i>	Butternut
<i>Juglans nigra</i>	Black walnut
<i>Prunus serotina</i>	Black cherry
<i>Quercus alba</i>	White oak
<i>Quercus macrocarpa</i>	Bur oak
<i>Quercus rubra</i>	Northern red oak
<i>Tilia americana</i>	Basswood
<i>Ulmus americana</i>	American elm
<i>Ulmus rubra</i>	Slippery elm

Trees – Sub-canopy	
<i>Betula papyrifera</i>	Paper-birch
<i>Carpinus caroliniana</i>	Blue beech
<i>Carya cordiformis</i>	Bitternut hickory
<i>Fraxinus nigra</i>	Black ash
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Ostrya virginiana</i>	Ironwood
<i>Pinus strobus</i>	White pine
<i>Prunus americana</i>	Wild plum
<i>Prunus virginiana</i>	Chokecherry

Shrubs	
<i>Cornus alternifolia</i>	Pagoda dogwood
<i>Cornus foemina</i>	Gray dogwood
<i>Dirca palustris</i>	Leatherwood
<i>Ribes americanum</i>	Wild black currant
<i>Ribes cynosbati</i>	Prickly gooseberry
<i>Ribes missouriense</i>	Missouri gooseberry
<i>Sambucus canadensis</i>	Common elder
<i>Sambucus pubens</i>	Red-berried elder
<i>Staphylea trifolia</i>	Bladdernut

Vines	
<i>Celastrus scandens</i>	Climbing bittersweet
<i>Parthenocissus inserta</i>	Five-leafed Virginia creeper
<i>Parthenocissus quinquefolia</i>	Virginia creeper

Forbs	
<i>Actaea rubra</i>	Red baneberry
<i>Adiantum pedatum</i>	Maidenhair fern

<i>Allium burdickii</i>	Burdick's leek
<i>Allium tricoccum</i>	Wild leek
<i>Amphicarpaea bracteata</i>	Hog-peanut
<i>Anemone quinquefolia</i>	Wood-anemone
<i>Anemone virginiana</i>	Virginia thimbleweed
<i>Anemonella thalictroides</i>	Rue-anemone
<i>Aquilegia canadensis</i>	Columbine
<i>Aralia nudicaulis</i>	Wild sarsaparilla
<i>Arisaema triphyllum</i>	Jack in the pulpit
<i>Asarum canadense</i>	Wild ginger
<i>Aster cordifolius</i>	Heart-leaved aster
<i>Athyrium angustum</i>	Lady-fern
<i>Botrychium virginianum</i>	Rattlesnake fern
<i>Campanula americana</i>	Tall bellflower
<i>Caulophyllum thalictroides</i>	Blue cohosh
<i>Cirsium discolor</i>	Field thistle
<i>Claytonia caroliniana</i>	Carolina spring-beauty
<i>Claytonia virginica</i>	Virginia spring-beauty
<i>Corallorhiza</i>	Coral-root
<i>Cypripedium calceolus</i>	Yellow lady-slipper
<i>Cystopteris bulbifera</i>	Bulblet bladder-fern
<i>Cystopteris fragilis</i>	Fragile bladder-fern
<i>Desmodium glutinosum</i>	Pointed-leaved tick-trefoil
<i>Dicentra cucullaria</i>	Dutchman's breeches
<i>Dryopteris cristata</i>	Crested fern
<i>Equisetum pratense</i>	Meadow horsetail
<i>Erythronium albidum</i>	White trout-lily
<i>Eupatorium rugosum</i>	Common snakeroot
<i>Galearis spectabilis</i>	Showy orchis
<i>Galium concinnum</i>	Elegant bedstraw
<i>Galium triflorum</i>	Three-flowered bedstraw
<i>Geranium maculatum</i>	Wild geranium
<i>Helianthus hirsutus</i>	Woodland sunflower
<i>Hepatica acutiloba</i>	Sharp-lobed hepatica
<i>Hydrophyllum virginianum</i>	Virginia waterleaf
<i>Isopyrum biternatum</i>	False rue-anemone
<i>Lilium michiganense</i>	Michigan lily
<i>Lonicera canadensis</i>	Fly honeysuckle
<i>Maianthemum canadense</i>	Canada mayflower
<i>Matteuccia struthiopteris</i>	Ostrich-fern
<i>Menispermum canadense</i>	Canada moonseed
<i>Onoclea sensibilis</i>	Sensitive fern
<i>Osmorhiza claytonii</i>	Clayton's sweet cicely
<i>Osmorhiza longistylis</i>	Anise-root
<i>Osmunda claytoniana</i>	Interrupted fern
<i>Panax quinquefolium</i>	American ginseng
<i>Phlox divaricata</i>	Blue phlox
<i>Polygonatum commutatum</i>	Giant Solomon's-seal
<i>Polygonatum pubescens</i>	Hairy Solomon's seal
<i>Prenanthes alba</i>	White rattlesnake-root
<i>Ranunculus abortivus</i>	Kidney-leaf buttercup
<i>Rubus occidentalis</i>	Black raspberry

<i>Rubus strigosus</i>	Red raspberry
<i>Rudbeckia laciniata</i>	Goldenglow
<i>Sanguinaria canadensis</i>	Bloodroot
<i>Thalictrum dioicum</i>	Early meadowrue
<i>Trillium cernuum</i>	Nodding trillium
<i>Uvularia grandiflora</i>	Yellow bellwort
<i>Uvularia sessilifolia</i>	Pale bellwort
<i>Viola pratincola</i>	Meadow violet
<i>Viola pubescens</i>	Yellow violet
<i>Viola sororia</i>	Common blue violet

Grasses and Sedges	
<i>Carex blanda</i>	Charming sedge
<i>Carex pedunculata</i>	Long-stalked sedge
<i>Carex pennsylvanica</i>	Pennsylvania sedge
<i>Carex rosea</i>	Stellate sedge
<i>Carex sprengelii</i>	Sprengel's sedge
<i>Elymus hystrix</i>	Bottlebrush grass
<i>Elymus villosus</i>	Downy wild rye
<i>Festuca obtusa</i>	Nodding fescue
<i>Milium effusum</i>	Woodland millet grass
<i>Oryzopsis asperifolia</i>	Mountain rice-grass
<i>Oryzopsis asperifolia</i>	Mountain rice-grass
<i>Oryzopsis racemosa</i>	Black-fruited rice-grass
<i>Schizachne purpurascens</i>	False medic grass

Appendix B Table 4. Dry Oak Forest species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees – Canopy	
<i>Ostrya virginiana</i>	Ironwood
<i>Populus tremuloides</i>	Quaking Aspen
<i>Quercus alba</i>	White Oak
<i>Quercus ellipsoidalis</i>	Northern Pin Oak
<i>Quercus macrocarpa</i>	Bur Oak

Trees – Sub-canopy	
<i>Prunus serotina</i>	Black Cherry
<i>Quercus macrocarpa</i>	Bur Oak
<i>Acer rubrum</i>	Red maple

Shrubs	
<i>Amelanchier laevis</i>	Smooth Juneberry
<i>Amelanchier sanguinea</i>	Round-leaf Juneberry
<i>Cornus racemosa</i>	Gray Dogwood
<i>Corylus americana</i>	American Hazelnut
<i>Diervilla lonicera</i>	Bush Honeysuckle
<i>Prunus virginiana</i>	Choke Cherry
<i>Ribes cynosbati</i>	Prickly Gooseberry
<i>Ribes missouriense</i>	Missouri Gooseberry
<i>Rosa blanda</i>	Smooth Wild Rose
<i>Sambucus canadensis</i>	Common Elder
<i>Symphoricarpos alba</i>	Snowberry
<i>Vaccinium angustifolium</i>	Lowbush Blueberry
<i>Viburnum lentago</i>	Nannyberry
<i>Viburnum rafinesquianum</i>	Downy Arrow-wood

Vines	
<i>Lonicera prolifera</i>	Yellow Vine Honeysuckle

Forbs	
<i>Agrimonia gryposepala</i>	Common Agrimony
<i>Amphicarpa bracteata</i>	Hog-peanut
<i>Anemone cylindrica</i>	Long-headed Thimbleweed
<i>Anemone quinquefolia</i>	Wood Anemone
<i>Anemone riparia (virginiana)</i>	Tall Thimbleweed
<i>Anemonella thalictroides</i>	Rue Anemone
<i>Apocynum androsaemifolium</i>	Spreading Dogbane
<i>Aquilegia canadensis</i>	Canada Columbine
<i>Aralia nudicaulis</i>	Wild Sarsaparilla
<i>Arenaria lateriflora</i>	Grove Sandwort
<i>Aster macrophyllus</i>	Large-leaved Aster
<i>Aster sagittifolius (urophyllus)</i>	Arrow-leaved Aster
<i>Athyrium filix-femina</i>	Lady Fern
<i>Botrychium dissectum</i>	Dissected Grape-fern

<i>Botrychium virginianum</i>	Rattlesnake Fern
<i>Cerastium nutans</i>	Nodding Chickweed
<i>Desmodium glutinosum</i>	Pointed-leaved Tick-trefoil
<i>Euphorbia corollata</i>	Flowering Spurge
<i>Fragaria virginiana</i>	Thick-leaved Wild Strawberry
<i>Galium aparine</i>	Cleavers
<i>Galium boreale</i>	Northern Bedstraw
<i>Galium concinnum</i>	Shining Bedstraw
<i>Galium triflorum</i>	Sweet-scented Bedstraw
<i>Geranium maculatum</i>	Wild geranium
<i>Hackelia virginiana</i>	Virginia Stickseed
<i>Helianthus hirsutus</i>	Woodland Sunflower
<i>Helianthus strumosus</i>	Rough-leaved Sunflower
<i>Heuchera richardsonii</i>	Alum-root
<i>Maianthemum canadense</i>	Canada Mayflower
<i>Monarda fistulosa</i>	Wild Bergomat
<i>Osmorhiza claytonii</i>	Clayton's Sweet-cicely
<i>Phryma leptostachya</i>	Lopseed
<i>Polygonatum biflorum (commutatum)</i>	Giant Solomon's Seal
<i>Pteridium aquilinum</i>	Bracken Fern
<i>Pyrola elliptica</i>	Common Shinleaf
<i>Sanicula gregaria</i>	Clustered Snakeroot
<i>Sanicula marilandica</i>	Black Snakeroot
<i>Smilacina racemosa</i>	False Solomon's Seal
<i>Smilacina stellata</i>	Starry false Solomon's Seal
<i>Smilax ecirrhata</i>	Cat-briar
<i>Smilax herbacea</i>	Cat-briar
<i>Solidago hispida</i>	Hairy Goldenrod
<i>Trientalis borealis</i>	Starflower
<i>Triosteum perfoliatum</i>	Perfoliate Horse-gentian
<i>Uvularia sessilifolia</i>	Sessile-leaved Bellwort
<i>Veronicastrum virginicum</i>	Culver's Root
<i>Viola pubescens</i>	Yellow Violet

Grasses and Sedges	
<i>Carex cephalophora</i>	Oval-headed Sedge
<i>Carex gracillima</i>	Graceful Sedge
<i>Carex hirtifolia</i>	Hairy-leaved Sedge
<i>Carex peckii</i>	Peck's Sedge
<i>Carex pensylvanica</i>	Pennsylvania Sedge
<i>Carex rosea</i>	Stellate Sedge
<i>Elymus hystrix (Hystrix patula)</i>	Bottlebrush Grass
<i>Oryzopsis asperifolia</i>	Rough-leaved Ricegrass
<i>Schizachne purpurascens</i>	False Melic Grass

Appendix B Table 5. Lowland Hardwood Forest species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees - Canopy	
<i>Acer rubrum</i>	Red maple
<i>Betula papyrifera</i>	Paper birch
<i>Celtis occidentalis</i>	Hackberry
<i>Fraxinus nigra</i>	Black ash
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Populus tremuloides</i>	Trembling aspen
<i>Quercus alba</i>	White oak
<i>Quercus macrocarpa</i>	Bur oak
<i>Quercus rubra</i>	Red oak
<i>Tilia americana</i>	Basswood
<i>Ulmus americana</i>	American elm
<i>Ulmus rubra</i>	Red elm

Trees – Sub-canopy	
<i>Acer saccharum</i>	Sugar maple
<i>Betula papyrifera</i>	Paper birch
<i>Carpinus caroliniana</i>	American hornbeam (blue beech)
<i>Carya cordiformis</i>	Bitternut
<i>Celtis occidentalis</i>	Hackberry
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Juglans cinerea</i>	Butternut
<i>Ostrya virginiana</i>	Ironwood
<i>Ulmus rubra</i>	Red elm

Shrubs	
<i>Alnus incana</i>	Speckled alder
<i>Cornus alternifolia</i>	Pagoda dogwood
<i>Cornus foemina</i>	Gray dogwood
<i>Corylus americana</i>	American hazelnut
<i>Euonymus atropurpureus</i>	Wahoo
<i>Ilex verticillata</i>	Winter berry
<i>Prunus virginiana</i>	Chokecherry
<i>Ribes americanum</i>	Wild black current
<i>Ribes missouriense</i>	Missouri gooseberry
<i>Sambucus canadensis</i>	Common elder
<i>Zanthoxylum americanum</i>	Prickly ash

Vines	
<i>Menispermum canadense</i>	Canada moonseed
<i>Parthenocissus spp.</i>	Virginia creeper
<i>Rhus radicans</i>	Poison ivy
<i>Vitis riparia</i>	Wild grape

Forbs	
<i>Amphicarpa bracteata</i>	Hog peanut
<i>Adiantum pedatum</i>	Maidenhair fern
<i>Aralia nudicaulis</i>	Wild sarsaparilla
<i>Arisaema triphyllum</i>	Jack-in-the-pulpit
<i>Asarum canadense</i>	Wild ginger
<i>Aster lateriflorus</i>	Side-flowering aster
<i>Athyrium filix-femina</i>	Lady fern
<i>Campanula americana</i>	Tall bellflower
<i>Circaea lutetiana</i>	Enchanter's nightshade
<i>Cryptotaenia canadensis</i>	Honewort
<i>Equisetum sylvaticum</i>	Woodland horsetail
<i>Galium aparine</i>	Cleavers
<i>Galium triflorum</i>	Three-flowered bedstraw
<i>Geum canadense</i>	White avens
<i>Hydrophyllum virginianum</i>	Virginia waterleaf
<i>Impatiens capensis</i>	Spotted touch-me-not
<i>Laportea canadensis</i>	Wood nettle
<i>Maianthemum canadense</i>	Canada mayflower
<i>Matteuccia struthiopteris</i>	Ostrich fern
<i>Osmunda claytoniana</i>	Interrupted fern
<i>Osmunda cinnamomea</i>	Cinnamon fern
<i>Panax quinquefolium</i>	Ginseng
<i>Pilea pumila</i>	Clearweed
<i>Rudbeckia laciniata</i>	Goldenglow
<i>Scutellaria lateriflora</i>	Skullcap
<i>Silphium perfoliatum</i>	Cup plant
<i>Smilacina racemosa</i>	False Solomon's seal
<i>Smilax spp.</i>	Carrion flower
<i>Stachys palustris</i>	Woundwort
<i>Thalictrum dioicum</i>	Early meadow-rue
<i>Teucrium canadense</i>	Germander
<i>Urtica dioica</i>	Stinging nettle

Grasses and Sedges	
<i>Carex gracillima</i>	A species of sedge
<i>Carex typhina</i>	A species of sedge
<i>Carex pedunculata</i>	A species of sedge
<i>Elymus virginicus</i>	Virginia wild rye
<i>Festuca subverticillata</i>	Nodding fescue
<i>Leersia virginica</i>	White grass
<i>Muhlenbergia frondosa</i>	Swamp satin grass
<i>Oryzopsis asperifolia</i>	Mountain ricegrass

Appendix B Table 6. Floodplain Forest species list

This species list has been compiled from Curtis (1959), Wovcha et al. (1994) and from plant inventory lists compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees - Canopy	
<i>Acer saccharinum</i>	Silver maple
<i>Acer negundo</i>	Boxelder
<i>Fraxinus nigra</i>	Black ash
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Populus deltoides</i>	Cottonwood
<i>Quercus macrocarpa</i>	Bur oak
<i>Ulmus americana</i>	American elm
<i>Ulmus rubra</i>	Red elm

Trees – Sub-canopy	
<i>Acer negundo</i>	Boxelder
<i>Acer saccharinum</i>	Silver maple
<i>Celtis occidentalis</i>	Hackberry
<i>Salix nigra</i>	Black willow
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Populus deltoides</i>	Cottonwood
<i>Tilia americana</i>	Basswood
<i>Ulmus americana</i>	American elm

Shrubs	
<i>Cephalanthus occidentalis</i>	Buttonbush
<i>Salix interior</i>	Sandbar willow
<i>Staphylea trifolia</i>	Bladdernut
<i>Zanthoxylum americanum</i>	Prickly ash

Vines	
<i>Parthenocissus spp.</i>	Virginia creeper
<i>Rhus radicans</i>	Poison ivy
<i>Vitis riparia</i>	Wild grape

Forbs	
<i>Amphicarpa bracteata</i>	Hog-peanut
<i>Apios americana</i>	Groundnut
<i>Aster ontarionis</i>	Ontario aster
<i>Bidens spp.</i>	Beggar-ticks
<i>Boehmeria cylindrica</i>	False nettle
<i>Cryptotaenia canadensis</i>	Honewort
<i>Echinocystis lobata</i>	Wild cucumber
<i>Eupatorium rugosum</i>	White snakeroot
<i>Galium aparine</i>	Cleavers
<i>Hydrophyllum virginianum</i>	Virginia waterleaf
<i>Impatiens capensis</i>	Spotted touch-me-not
<i>Laportea canadensis</i>	Wood nettle

<i>Lobelia cardinalis</i>	Cardinal flower
<i>Lycopus virginicus</i>	Virginia water horehound
<i>Physostegia virginiana</i>	False dragonhead
<i>Pilea pumila</i>	Clearweed
<i>Rudbeckia laciniata</i>	Goldenglow
<i>Scutellaria lateriflora</i>	Skullcap
<i>Sicyos angulatus</i>	Bur cucumber
<i>Stachys hispida</i>	Smooth hedge nettle
<i>Stachys tenuifolia</i>	Narrow-leaved hedge nettle
<i>Urtica dioica</i>	Stinging nettle

Grasses and Sedges	
<i>Carex crinita</i>	A species of sedge
<i>Carex tribuloides</i>	A species of sedge
<i>Carex typhina</i>	Cattail sedge
<i>Cares lupulina</i>	A species of sedge
<i>Echinochloa walteri</i>	Walter's barnyard grass
<i>Elymus virginicus</i>	Virginia wild rye
<i>Leersia oryzoides</i>	Rice cut-grass
<i>Leersia virginica</i>	White grass

Appendix B Table 7. Mixed Emergent Marsh species list

This species list has been compiled by Cynthia Lane, Ph.D., former staff ecologist with Great River Greening.

Latin Name	Common Name
Trees	
<i>Acer saccharinum</i>	Silver maple
<i>Celtis occidentalis</i>	Hackberry
<i>Fraxinus pennsylvanica</i>	Green ash
<i>Salix exigua interior</i>	Sandbar willow

Shrubs	
<i>Amorpha fruticosa</i>	False indigo
<i>Betula pumila</i>	Bog-birch
<i>Cornus sericea</i>	Red-osier dogwood
<i>Salix eriocephala</i>	Heart-leaved willow
<i>Salix petiolaris</i>	Meadow willow
<i>Sambucus canadensis</i>	Common elder
<i>Spirea tomentosa rosea</i>	Steeple-bush

Vines	
<i>Cuscuta spp.</i>	Dodder
<i>Decodon verticillatus laevigatus</i>	Water willow
<i>Echinocystis lobata</i>	Wild cucumber
<i>Menispermum canadense</i>	Common moonseed
<i>Smilax hispida</i>	Green-briar

Forbs	
<i>Acorus calamus</i>	Sweet flag
<i>Alisma subcordatum</i>	Heart-leaved water-plantain
<i>Alisma triviale</i>	Ordinary water-plantain
<i>Amaranthus tuberculatus</i>	Tall water hemp
<i>Ambrosia trifida</i>	Great ragweed
<i>Apocynum androsaemifolium</i>	Spreading dogbane
<i>Artemisia serrata</i>	Leafy mugwort
<i>Asclepias incarnata</i>	Swamp milkweed
<i>Aster ontarionis</i>	Ontario aster
<i>Bidens spp.</i>	Beggar-ticks
<i>Boehmeria cylindrica</i>	False nettle
<i>Campanula aparinoides</i>	Marsh bellflower
<i>Cicuta bulbifera</i>	Bulb-bearing water-hemlock
<i>Cicuta maculata</i>	Spotted water-hemlock
<i>Epilobium spp.</i>	Willow-herb
<i>Eupatorium maculatum</i>	Spotted Joe-pye weed
<i>Eupatorium perfoliatum</i>	Common boneset
<i>Eupatorium purpureum</i>	Sweet Joe-pye weed
<i>Eupatorium rugosum</i>	Common snakeroot
<i>Galium labradoricum</i>	Marsh bedstraw

<i>Galium tinctorium</i>	Small bedstraw
<i>Galium trifidum</i>	Three-cleft bedstraw
<i>Impatiens</i> spp.	Spotted touch-me-not
<i>Iris versicolor</i>	Northern blue flag
<i>Laportea canadensis</i>	Wood-nettle
<i>Lathyrus palustris</i>	Marsh vetchling
<i>Lemna</i> spp.	Lesser duckweed
<i>Lobelia cardinalis</i>	Cardinal flower
<i>Ludwigia palustris</i>	Water purslane
<i>Lycopus americanus</i>	Cut-leaved bugleweed
<i>Lycopus asper</i>	Bugleweed
<i>Lycopus uniflorus</i>	Northern bugleweed
<i>Lycopus virginicus</i>	Virginia bugleweed
<i>Lysimachia ciliata</i>	Fringed loosestrife
<i>Lysimachia terrestris</i>	Yellow loosestrife
<i>Lysimachia thyrsiflora</i>	Tufted loosestrife
<i>Mentha arvensis glabrata</i>	Common mint
<i>Mimulus ringens</i>	Purple monkey-flower
<i>Nymphaea</i> cmx.	Water lily
<i>Oenothera biennis</i>	Common evening-primrose
<i>Oxalis</i> cmx.	Wood-sorrel
<i>Physostegia virginiana</i>	Obedient plant
<i>Pilea</i> spp.	Clearweed
<i>Polygonum amphibium stipulaceum</i>	Water smartweed
<i>Polygonum amphibium</i>	Swamp smartweed
<i>Polygonum lapathifolium</i>	Nodding smartweed
<i>Polygonum pensylvanicum</i>	Pennsylvania smartweed
<i>Polygonum punctatum</i>	Dotted smartweed
<i>Polygonum sagittatum</i>	Arrow-leaved tearthumb
<i>Polygonum virginianum</i>	Virginia knotweed
<i>Ranunculus pensylvanicus</i>	Bristly buttercup
<i>Rorippa palustris</i>	Yellow cress
<i>Rudbeckia laciniata</i>	Goldenglow
<i>Rumex maritimus fueginus</i>	Golden dock
<i>Rumex orbiculatus</i>	Great water dock
<i>Sagittaria latifolia</i>	Broad-leaved arrowhead
<i>Scutellaria galericulata</i>	Marsh skullcap
<i>Scutellaria lateriflora</i>	Mad-dog skullcap
<i>Sium suave</i>	Water-parsnip
<i>Solidago gigantea</i>	Giant goldenrod
<i>Sparganium androcladum</i>	Bur reed
<i>Sparganium emersum</i>	Bur-reed
<i>Sparganium eurycarpum</i>	Giant bur-reed
<i>Spirodela polyrhiza</i>	Greater duckweed
<i>Stellaria longifolia</i>	Long-leaved chickweed
<i>Teucrium canadense</i>	Germander
<i>Typha</i> spp.	Cattail
<i>Urtica dioica gracilis</i>	Stinging nettle
<i>Verbena hastata</i>	Blue vervain

Grasses and Sedges	
<i>Calamagrostis canadensis</i>	Bluejoint
<i>Carex aquatilis</i>	Water sedge
<i>Carex comosa</i>	Bristly sedge
<i>Carex diandra</i>	Lesser-panicled sedge
<i>Carex haydenii</i>	Hayden's sedge
<i>Carex lacustris</i>	Lake-sedge
<i>Carex pellita</i>	Woolly sedge
<i>Carex stricta</i>	Tussock-sedge
<i>Cyperus bipartitus</i>	Nut grass (a type of sedge)
<i>Cyperus diandrus</i>	Nut grass (a type of sedge)
<i>Cyperus erythrorhizos</i>	Nut grass (a type of sedge)
<i>Cyperus odoratus</i>	Nut grass (a type of sedge)
<i>Cyperus strigosus</i>	Nut grass (a type of sedge)
<i>Elymus virginicus</i>	Virginia wild rye
<i>Equisetum fluviatile</i>	Water horsetail
<i>Dulichium arundinaceum</i>	Three-way sedge
<i>Echinochloa muricata</i>	Barnyard grass
<i>Eleocharis ovata</i>	Spike rush
<i>Eleocharis pauciflora fernaldii</i>	Spike rush
<i>Glyceria canadensis</i>	Rattlesnake grass
<i>Leersia oryzoides</i>	Rice cut grass
<i>Leersia virginica</i>	White grass
<i>Phalaris arundinacea</i>	Reed canary-grass
<i>Phragmites australis</i>	Common reed
<i>Scirpus acutus</i>	Hard-stemmed bullrush
<i>Scirpus cyperinus</i>	Wool-grass
<i>Scirpus fluviatilis</i>	River bulrush
<i>Scirpus validus creber</i>	Softstem bullrush
<i>Spartina pectinata</i>	Prairie cord-grass
<i>Thelypteris palustris</i>	Northern marsh-fern
<i>Zizania palustris</i>	Wild rice

Appendix B Table 8. Willow Swamp species list

This species list has been compiled from Wovcha et al. (1994).

Latin Name	Common Name
Shrubs	
<i>Salix gracilis</i>	Slender willow
<i>Salix discolor</i>	Pussy willow
<i>Salix bebbiana</i>	Bebb's willow
<i>Alnus incana rugosa</i>	Speckled alder
<i>Cornus stolonifera</i>	Red-osier dogwood
<i>Betula glandulifera</i>	Bog birch

Forbs	
<i>Thelypteris palustris</i>	Northern marsh fern
<i>Sagittaria latifolia</i>	Broad-leaved arrowhead
<i>Campanula aparinoides</i>	Marsh bellflower
<i>Cicuta bulbifera</i>	Bulb-bearing water-hemlock
<i>Eupatorium maculatum</i>	Joe-pye weed
<i>Potentilla palustris</i>	Marsh cinquefoil
<i>Rumex orbiculatus</i>	Great water dock
<i>Impatiens capensis</i>	Spotted touch-me-not
<i>Lysimachia thyrsiflora</i>	Tufted loosestrife


Grasses and Sedges	
<i>Calamagrostis canadensis</i>	Blue-joint grass
<i>Carex lacustris</i>	Lake sedge
<i>Carex stricta</i>	Tussock sedge

Appendix C: Cherokee Park Inventory Results

Introduction

This report presents the results of an ecological inventory of Cherokee Park conducted during the growing season of 2002. Cherokee Park is a city park that is located on the south side of the Mississippi River in St. Paul. Land-cover maps developed through the Minnesota Land Cover Classification System (MLCCS) were used to develop species lists. Separate species lists were compiled for each land-cover type within the project area. Plants are listed by vegetation form and are listed alphabetically by scientific name. Written descriptions summarizing the ecological condition of each land-cover type are included in the report.

Maple-basswood forest - 12.35 acres



The maple-basswood forest at the project site generally faces northeast and is on a relatively steep portion of bluff. Maple-basswood forest species are most dominant within ravines while ridge tops are often dominated with dry to mesic oak forest species. Sugar maple trees and ironwood dominate the maple-basswood forest along with basswood, red oak and hackberry.

Generally, the shrub and ground layers are diverse. Shrub species that are common but not generally seen in other Twin Cities woodlands include bladdernut, roundleaf dogwood, and leatherwood. Also uncommon in the Twin Cities area are spring ephemerals. It is believed that combination of erosion, trampling and non-native earthworms are causing spring ephemeral populations to decline. Within the maple-basswood forest at Cherokee Park many spring ephemeral can be found. Some species include sharp-lobed hepatica, bloodroot, rue-anemone and wild sarsaparilla. Sedges are also common in the maple basswood forest and include Pennsylvania sedge, woodland sedge, Sprengell's sedge and *Carex eburina*. *Carex eburina* is a sedge with very fine leaf blades and it is found in dense groupings on some steeper portions of the bluff. These sedges play an important role in preventing erosion. Although erosion is a natural process of steep slopes and ravines, hiking/animal trails and increased water runoff from above are significantly accelerating erosion in the park.

The most common invasive species include common buckthorn and Tartarian honeysuckle. Common buckthorn comprises about 20% of the mid-story and Tartarian honeysuckle makes up about 1% of the shrub layer.

Canopy trees

<i>Latin Name</i>	<i>Common Name</i>	<i>Percent Cover</i>
<i>Acer saccharum</i>	Sugar maple	25%
<i>Carya cordiformis</i>	Bitternut hickory	4%
<i>Celtis occidentalis</i>	Hackberry	8%
<i>Juglans cinerea</i>	Butternut	7%
<i>Populus deltoides</i>	Eastern cottonwood	8%
<i>Quercus alba</i>	White oak	5%
<i>Quercus ellipsoides</i>	Pin oak	5%
<i>Quercus macrocarpa</i>	Bur oak	3%
<i>Quercus rubra</i>	Red oak	10%
<i>Tilia americana</i>	American basswood	15%
<i>Ulmus americana</i>	American elm	5%
<i>Ulmus rubra</i>	Red elm	10%

Mid-Story trees

<i>Ostrya virginiana</i>	Ironwood	25%
<i>Rhamnus cathartica</i>	Common buckthorn	20%

Shrub layer

<i>Amelanchier laevis</i>	Smooth serviceberry	1%
<i>Catalpa speciosa</i>	Catalpa	<1%
<i>Cornus alternifolia</i>	Pagoda dogwood	<1%
<i>Cornus racemosa</i>	Grey dogwood	4%
<i>Cornus rugosa</i>	Round-leaved dogwood	<1%
<i>Dirca palustris</i>	Leatherwood	<1%
<i>Juniperus virginiana</i>	Red Cedar	<1%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	1%
<i>Prunus americana</i>	Chokecherry	10%
<i>Rhus typhina</i>	Staghorn sumac	<1%
<i>Ribes cynosbati</i>	Gooseberry (thornless)	<1%
<i>Ribes missouriensis</i>	Prickly gooseberry?	<1
<i>Sambucus spp.</i>	Elderberry	<1%
<i>Staphylea trifolia</i>	Bladdernut	1%
<i>Symphoricarpos occidentalis</i>	Wolfberry	1%
<i>Viburnum lentago</i>	Nannyberry	<1%
<i>Viburnum rafenesquianum</i>	Downy arrowwood viburnum	3%
<i>Xanthoxylum americanum</i>	Prickly ash	<1%

Groundlayer vines

<i>Amphicarpa bracteata</i>	Hog peanut	1%
<i>Parthenocissus inserta</i>	Virginia creeper	1%
<i>Rhus toxicodendron</i>	Poison ivy	1%
<i>Vitis riparia</i>	Riverbank grape	<1%

Forbs or seedling trees


<i>Anemone cylindrica</i>	Thimbleweed	<1%
<i>Aquilegia canadense</i>	Columbine	<1%
<i>Aralia nudicaulis</i>	Wild sarsaparilla	<1%
<i>Arctium minus</i>	Burdock	<1%

<i>Arisaema atrorubens</i>	Jack-in-the pulpit	<1%
<i>Asarum canadense</i>	Wild ginger	<1%
<i>Aster laevis</i>	Smooth aster	<1%
<i>Desmodium glutinosum</i>	Pointed-leaved tick trefoil	<1%
<i>Equisetum pratense</i>	Horsetail	<1%
<i>Galium boreale</i>	Northern bedstraw	<1%
<i>Helianthus divaricatus</i>	Woodland sunflower	<1%
<i>Hepatica acutiloba</i>	Sharp lobed hepatica	<1%
<i>Hydrophyllum virginianum</i>	Virginia waterleaf	<1%
<i>Lactuca canadensis</i>	Wild lettuce	<1%
<i>Maianthemum stellatum</i>	False Solomon's seal	<1%
<i>Melilotus officinalis</i>	Yellow sweet clover	<1%
<i>Sanguinaria canadensis</i>	Bloodroot	<1%
<i>Smilax rotundifolia</i>	Greenbriar	<1%
<i>Solidago flexicaulis</i>	Zig-Zag goldenrod	1%

Grasses and sedges

<i>Carex blanda</i>	Woodland sedge	<1%
<i>Carex eburina</i>	A species of sedge	1%
<i>Carex pennsylvanica</i>	Pennsylvania sedge	5%
<i>Hystrix patula</i>	Bottlebrush grass	<1
<i>Oryzopsis racemosa</i>	Black-seeded rice grass	<1%

Short grasses and mixed trees with 26-50% impervious cover - 13.35 acres



This land-cover area along Cherokee Heights Boulevard and Cherokee Avenue is made up of mown lawn with scattered boulevard trees. Of the wide variety of tree species planted species planted along Cherokee Heights Boulevard and Chippewa Avenue, hackberry is the most common. Other common deciduous trees include white oak, red oak and little-leaf linden. A number of evergreens have also been planted here including Scotch pine, Norway spruce, Colorado blue spruce and red cedar.

The lawn is a typical park lawn composed of a variety of grasses and weeds. Much of the lawn extends to the edge of the bluff, allowing rainwater to flow at an accelerated rate over the edge, adding to erosion on the bluff face. In this regard, converting a band of lawn adjacent to the wooded slopes to prairie, savanna and forest edge species would help to reduce runoff and slow erosion of the bluff.

Canopy trees

<i>Latin Name</i>	<i>Common Name</i>	<i>Percent Cover</i>
<i>Acer Platanoides</i>	Norway maple	3%
<i>Celtis occidentalis</i>	Hackberry	50%
<i>Fraxinus americana</i>	White ash	5%
<i>Fraxinus pennsylvanica</i>	Green ash	4%
<i>Juniperus virginiana</i>	Red cedar	1%
<i>Picea abies</i>	Norway spruce	2%
<i>Picea glauca</i>	White spruce	3%
<i>Pinus strobus</i>	White pine	5%
<i>Pinus sylvestris</i>	Scotch pine	1%
<i>Quercus alba</i>	White oak	25%
<i>Quercus macrocarpa</i>	Bur oak	1%
<i>Quercus rubra</i>	Red oak	2%
<i>Tilia cordata</i>	Little-leaf linden	8%
<i>Ulmus americana</i>	American elm	2%

Mid-Story trees

<i>Acer ginnala</i>	Amur maple	1%
<i>Malus sp.</i>	Crab apple	3%
<i>Rhamnus cathartica</i>	Common buckthorn	<1%

Shrub layer

<i>Syringa sp.</i>	Lilac	<1%
--------------------	-------	-----

Groundlayer vines

<i>None</i>		
-------------	--	--


Forbs or seedling trees

<i>Alisma plantago</i>	Common Plantain	2%
<i>Arctium minus</i>	Burdock	1%
<i>Glechoma sanguinalis</i>	Creeping Charlie	2%
<i>Melilotus alba</i>	Yellow sweet clover	<1%
<i>Melilotus officinale</i>	White sweet clover	<1%
<i>Taraxacum officinale</i>	Dandelion	2%
<i>Trifolium repens</i>	White clover	2%

Grasses and sedges

<i>Agropyron repens</i>	Quack grass	3%
<i>Digitaria sanguinalis</i>	Crabgrass	3%
<i>Poa pratensis</i>	Kentucky bluegrass	90%

Short grasses and sparse tree cover on upland soils - 9.06 acres



This land-cover area makes up the recreational portion of Cherokee Park with picnic tables, a playground and restrooms. This land-cover has many species in common with the land-cover type along Cherokee Heights Blvd. and Chippewa Ave. (short grasses and mixed trees with 26-50% impervious cover) but contains a large stand of red and bur oak trees. Other common tree species include hackberry and two non-native species, little leaf linden and

Norway maple. Generally the lawn areas have enough variation in topography to retain stormwater. Since most of this land-cover unit is heavily used by park visitors it would be difficult to incorporate many native plant reconstructions. The large grove of mature oak trees in the south part of this land cover area is a potential site for savanna restoration.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Acer platanoides</i>	Norway maple	8%
<i>Aesculus glabra</i>	Horse chestnut	1%
<i>Celtis occidentalis</i>	Hackberry	5%
<i>Fraxinus americana</i>	White ash	5%
<i>Fraxinus pennsylvanica</i>	Green ash	5%
<i>Picea abies</i>	Norway spruce	2%
<i>Pinus resinosa</i>	Red Pine	2%
<i>Pinus sylvestris</i>	Scotch pine	2%
<i>Quercus alba</i>	White oak	3%
<i>Quercus macrocarpa</i>	Bur oak	10%
<i>Quercus rubra</i>	Red oak	5%
<i>Tilia cordata</i>	Little-leaf linden	3%
<i>Ulmus rubra</i>	Red elm	3%

Mid-Story trees

<i>Acer ginnala</i>	Amur maple	1%
<i>Acer rubrum</i>	Red maple	1%
<i>Malus sp.</i>	Crabapple	1%
<i>Syringa reticulata</i>	Japanese tree lilac	1%

Shrub layer

<i>Spiraea sp.</i>	Spiraea	1%
--------------------	---------	----

Groundlayer vines

None		
------	--	--

Forbs or seedling trees

<i>Alisma plantago</i>	Common Plantain	2%
<i>Arctium minus</i>	Burdock	1%
<i>Glechoma hederacea</i>	Creeping Charlie	2%
<i>Melilotus alba</i>	Yellow sweet clover	<1%
<i>Melilotus officinale</i>	White sweet clover	<1%
<i>Taraxacum officinale</i>	Dandelion	2%
<i>Trifolium repens</i>	White clover	2%

Grasses and sedges

<i>Agropyron repens</i>	Quack grass	3%
<i>Digitaria sanguinalis</i>	Crabgrass	3%
<i>Poa pratensis</i>	Kentucky bluegrass	90%

Boxelder - green ash disturbed native forest (Northwest portion) – 10.56 acres



Between 1883 and 1973 a brick yard operated adjacent to the southwest edge of this land cover area. Extensive disturbance as a result of this operation included two railroad tracks that have since been abandoned, ditches and brick/soil disposal piles. Due to the resulting irregular soil surface, there are a variety of moisture conditions within the forest and most trees are relatively young.

Ditches are dominated by species such as reed canary grass, red-osier dogwood and American elm. Disturbed upland soils such as those along the railroad berms are dominated by species such as quaking aspen, cottonwood, Siberian elm, smooth brome and Canada goldenrod. The disturbed nature of the forest has made it ideal habitat for invasive species. Garlic mustard, Siberian elm, common buckthorn, Tartarian honeysuckle, smooth brome, black locust, creeping Charlie, reed canary grass and Kentucky bluegrass are all found at the site. Since this area is generally flat, erosion is not a significant problem. Deposition of eroded materials from the slopes above is extensive at some points along the base of the bluff.

Canopy trees

<i>Latin Name</i>	Common Name	Percent Cover
<i>Acer nigra</i>	Boxelder	5%
<i>Acer saccharum</i>	Sugar maple	2%
<i>Catalpa speciosa</i>	Catalpa	<1%
<i>Celtis occidentalis</i>	Hackberry	1%
<i>Fraxinus pennsylvanica</i>	Green ash	60%

<i>Populus deltoides</i>	Cottonwood	5%
<i>Populus tremuloides</i>	Quaking aspen	10%
<i>Quercus alba</i>	White oak	<1%
<i>Robinia pseudoacacia</i>	Black locust	<1%
<i>Salix nigra</i>	Black willow	<1%
<i>Ulmus americana</i>	American elm	15%

Mid-Story trees

<i>Rhamnus cathartica</i>	Common buckthorn	10%
<i>Ulmus pumila</i>	Siberian elm	1%

Shrub layer

<i>Amorpha fruticosa</i>		<1%
<i>Cornus sericea</i>	Red-osier dogwood	10%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	3%
<i>Prunus virginiana</i>	Chokecherry	1%
<i>Ribes</i>	Gooseberry	1%
<i>Rubus sp.</i>	Raspberry	<1%
<i>Salix exigua</i>	Sandbar willow	1%
<i>Viburnum lentago</i>	Nannyberry	<1%
<i>Viburnum rafinesquianum</i>	Arrowwood viburnum	<1%

Groundlayer vines

<i>Amphicarpa bracteata</i>	Hog peanut	<1%
<i>Parthenocissus inserta</i>	Virginia creeper	<1%
<i>Rhus toxicodendron</i>	Poison ivy	<1%

Forbs or seedling trees


<i>Alliaria petiolata</i>	Garlic mustard	<1%
<i>Arctium minus</i>	Burdock	<1%
<i>Aster puniceus</i>	Red -stem aster	<1%
<i>Aster sp.</i>	Aster	4%
<i>Aster-novae angliae</i>	New England aster	<1%
<i>Cornus serotina</i>	Black cherry	<1%
<i>Equisetum sp.</i>	Equisetum	1%
<i>Eupatorium rugosum</i>	White snakeroot	1%
<i>Glechoma hederacea</i>	Creeping Charlie	3%
<i>Helianthus divaricatus</i>	Woodland sunflower	<1%
<i>Nepeta cataria</i>	Catmint	<1%
<i>Pilea pumila</i>	Clearweed	<1%
<i>Rhus glabra</i>	Smooth sumac	<1%
<i>Rudbeckia laciniata</i>	Giant coneflower	<1%
<i>Solidago canadensis</i>	Canada goldenrod	3%
<i>Solidago canadensis</i>	Canada goldenrod	<1%
<i>Sonchus sp.</i>	Sow thistle	<1%
<i>Ulmus americana</i>	American elm	<1%

Grasses and sedges

<i>Bromus inermis</i>	Smooth brome	5%
<i>Carex blanda</i>	Woodland sedge	<1%
<i>Elymus canadensis</i>	Canada wild rye	<1%

<i>Leersia oryzoides</i>	Rice-cut grass	<1%
<i>Phalaris arundinacea</i>	Reed canary grass	1%
<i>Poa pratensis</i>	Kentucky bluegrass	10%
<i>Setaria glauca</i>	Yellow foxtail	1%

51% to 75% Impervious cover with deciduous trees (Northern portion) - .45 acres



This is a small land-cover unit located along West Water Street at the base of the bluff. It is an area of open grasses surrounded by relatively high quality maple-basswood forest. It appears that this area experienced soil slumping in the past or may have been an area where soils were mined. Due to past disturbance, the non-native grass species smooth brome and Kentucky bluegrass dominate the land-cover unit. The invasive legume species, crown vetch and

alfalfa are also abundant, indicating that the site may have been planted with a slope stabilization mix in the past. Early successional tree species are starting to colonize the site. Eastern cottonwood, boxelder, black locust, common buckthorn and green ash are all present. It is likely that the trees will eventually create too much shade for the smooth brome and Kentucky bluegrass to persist. Weedy understory species will most likely replace the grasses.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Acer negundo</i>	Boxelder	30%
<i>Fraxinus pennsylvanica</i>	Green ash	3%
<i>Populus deltoides</i>	Cottonwood	20%
<i>Robinia pseudoacacia</i>	Black locust	3%

Mid-Story trees

<i>Rhamnus cathartica</i>	Common buckthorn	5%
---------------------------	------------------	----

Shrub layer

<i>Cornus sericea</i>	Red-osier dogwood	3%
<i>Rhus typhina</i>	Staghorn sumac	3%

Groundlayer vines

<i>Vitis riparia</i>	River-bank grape	<1%
----------------------	------------------	-----

Forbs or seedling

<i>Arctium minus</i>	Burdock	20%
----------------------	---------	-----

<i>Coronilla varia</i>	Crown vetch	40%
<i>Medicago sativa</i>	Alfalfa	20%
<i>Solidago canadensis</i>	Canada goldenrod	3%

Grasses and sedges

<i>Bromus inermis</i>	Smooth brome	100%
<i>Poa pratensis</i>	Kentucky bluegrass	20%
<i>Typha latifolia</i>	Broad-leaved cattail	2%

51% to 75% Impervious cover with deciduous trees (Southern portion) - .22 acres



This small land-cover unit is West Water Street southwest of the bluff where the land flattens out in the floodplain. The site has signs of significant soil disturbance that may have occurred when an adjacent parking area was constructed. The site is dominated by boxelder trees that occupy nearly 100% of the canopy. Other canopy species include eastern cottonwood and green ash. None of the tree species appear over 30 years old. Little understory is present probably as a result of the dense shade produced by the boxelder. Riverbank grape and Canada goldenrod were the only two ground layer species present during the inventory.

Canopy trees

<i>Latin Name</i>	<i>Common Name</i>	<i>Percent Cover</i>
<i>Acer negundo</i>	Boxelder	100%
<i>Celtis occidentalis</i>	Hackberry	<1%
<i>Fraxinus pennsylvanica</i>	Green ash	5%
<i>Juglans nigra</i>	Black walnut	<1%
<i>Populus deltoides</i>	Cottonwood	5%

Mid-Story trees

<i>Rhamnus cathartica</i>	Common buckthorn	<1%
---------------------------	------------------	-----

Shrub layer

None		
------	--	--

Groundlayer vines

<i>Vitis riparia</i>	Riverbank grape	<1%
----------------------	-----------------	-----

Forbs or seedling trees

<i>Solidago gigantea</i>	Giant goldenrod	<1%
--------------------------	-----------------	-----

Grasses and sedges

None		
------	--	--

Disturbed deciduous woodland (Northern portion) – 4.43 acres



This area of disturbed deciduous woodland is found in the northernmost portion of the project area. The woodland is located just south of the Smith Avenue bridge and is on a very steep slope. It is likely that a combination of factors including bridge construction, erosion, the presence of invasive species and tree cutting have all contributed to the disturbed nature of the woodland. Both common buckthorn and Tartarian honeysuckle

are common on the slope. Other invasive species present during the inventory include black locust, Siberian elm, smooth brome, creeping Charlie and reed canary grass. Young sugar maple and ironwood trees are common within the woodland indicating that it may be developing into maple-basswood forest. There is little understory vegetation common to maple-basswood forests.

Canopy trees

<i>Latin Name</i>	Common Name	Percent Cover
<i>Acer negundo</i>	Boxelder	7%
<i>Acer platanoides</i>	Norway maple	4%
<i>Acer saccharum</i>	Sugar maple	30%
<i>Carya cordiformis</i>	Bitternut hickory	<1%
<i>Catalpa speciosa</i>	Catalpa	<1%
<i>Fraxinus pennsylvanica</i>	Green ash	7%
<i>Pinus nigra</i>	Austrian pine	4%
<i>Populus deltoides</i>	Cottonwood	3%
<i>Quercus ellipsoidalis</i>	Northern pin oak	2%
<i>Robinia pseudoacacia</i>	Black locust	<1%
<i>Tilia americana</i>	American basswood	12%
<i>Ulmus americana</i>	American elm	3%
<i>Ulmus pumila</i>	Siberian elm	<1%
<i>Ulmus rubra</i>	Red elm	3%

Mid-Story trees

<i>Crataegus sp.</i>	Hawthorn	<1%
<i>Ostrya virginiana</i>	Ironwood	30%
<i>Prunus americana</i>	Choke cherry	4%
<i>Rhamnus cathartica</i>	Common buckthorn	10%

Shrub layer

<i>Lonicera tartarica</i>	Tartarian honeysuckle	3%
<i>Ribes cynosbati</i>	Gooseberry (no thorns)	<1%
<i>Ribes missouriense</i>	Black currant	<1%
<i>Salix exigua</i>	Sandbar willow	1%
<i>Sambucus pubens</i>	Elderberry	<1%
<i>Xanthoxylum americanum</i>	Prickly ash	<1%

Groundlayer vines

<i>Parthenocissus inserta</i>	Virginia creeper	<1%
<i>Vitis riparia</i>	Riverbank grape	4%

Forbs or seedling trees


<i>Arctium minus</i>	Burdock	<1%
<i>Asarum canadense</i>	Wild ginger	<1%
<i>Aster cordifolius</i>	Heart leaved aster	<1%
<i>Glechoma hederacea</i>	Creeping Charlie	<1%
<i>Glechoma hederacea</i>	Creeping Charlie	<1%
<i>Hydrophyllum virginiana</i>	Virginia waterleaf	<1%
<i>Impatiens capensis</i>	Jewelweed	4%
<i>Leonurus cardiaca</i>	Motherwort	<1%
<i>Smilax rotundifolia</i>	Greenbriar	<1%
<i>Solidago canadensis</i>	Canada goldenrod	<1%
<i>Solidago flexicaulis</i>	Zig-zag goldenrod	1%
<i>Solidago gigantea</i>	Giant goldenrod	<1%
<i>Violet sp.</i>	Violet	<1%

Grasses and sedges

<i>Bromus inermis</i>	Smooth brome	4%
<i>Carex blanda</i>	Woodland sedge	1%
<i>Carex pennsylvanica</i>	Pennsylvania sedge	1%
<i>Phalaris arundinacea</i>	Reed canary grass	<1%

Disturbed deciduous woodland (Southern portion) – 10.64 acres

This area of disturbed deciduous woodland is located at the south end of the site. The woodland extends uphill (east) from a road that parallels the base of the bluff. The road was part of the brick yard operation located southwest of the project area. Overall, this area has had a significant amount of human disturbance for a long time. A number of sandstone caves found at the base of the bluff were primarily used for storage in the past.



The woodland extends approximately 1/3 of the way up the bluff where it meets a plateau supporting oak forest and boxelder-green ash disturbed native forest. None of the canopy trees in the unit are very old, showing that this is a relatively young forest. Common buckthorn dominates the shrub layer of this woodland. Other invasive species such as garlic mustard and Tartarian honeysuckle are also common.

American elm and cottonwood are common

canopy trees. There is little ground layer vegetation most likely due to trampling and the presence of invasive species. Garlic mustard is a common invasive species in the ground layer. The lack of ground layer vegetation on the steep slope contributes to erosion, particularly in ravines and where trails lead down the slope.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Acer saccharum</i>	Sugar maple	20%
<i>Betula papyrifera</i>	Paper birch	1%
<i>Carya cordiformis</i>	Bitternut hickory	1%
<i>Catalpa speciosa</i>	Catalpa	<1%
<i>Celtis occidentalis</i>	Hackberry	8%
<i>Fraxinus pennsylvanica</i>	Green ash	20%
<i>Populus deltoides</i>	Cottonwood	20%
<i>Quercus rubra</i>	Red oak	7%
<i>Tilia americana</i>	American basswood	1%
<i>Ulmus americana</i>	American elm	10%
<i>Ulmus rubra</i>	Red elm	10%

Mid-Story trees

<i>Ostrya virginiana</i>	Ironwood	5%
<i>Prunus americana</i>	Choke cherry	<1%
<i>Rhamnus cathartica</i>	Common buckthorn	50%

Shrub layer

<i>Amelanchier laevis</i>	Smooth serviceberry	1%
<i>Cornus alternifolia</i>	Pagoda dogwood	<1%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	3%
<i>Staphylea trifolia</i>	Bladdernut	<1%
<i>Viburnum lentago</i>	Nannyberry	<1%
<i>Viburnum rafinesquianum</i>	Arrowwood viburnum	1%
<i>Xanthoxylum americanum</i>	Prickly ash	<1%

Groundlayer vines

<i>Amphicarpa bracteata</i>	Hog peanut	1%
<i>Echinocystis lobata</i>	Wild cucumber	<1%
<i>Parthenocissus inserta</i>	Virginia creeper	<1%

<i>Rhus toxicodendron</i>	Poison ivy	4%
---------------------------	------------	----


Forbs or seedling trees

<i>Alliaria petiolata</i>	Garlic mustard	10%
<i>Aquilegia canadensis</i>	Columbine	<1%
<i>Asarum canadense</i>	Wild ginger	<1%
<i>Desmodium glutinosum</i>	Pointed leaved tick trefoil	<1%
<i>Hepatica acutiloba</i>	Sharp lobed hepatica	<1%
<i>Hydrophyllum virginiana</i>	Virginia waterleaf	1%
<i>Solidago flexicaulis</i>	Zig-zag goldenrod	1%

Grasses and sedges

<i>Carex blanda</i>	Woodland sedge	<1%
<i>Carex eburina</i>	A species of sedge	1%
<i>Carex pennsylvanica</i>	Pennsylvania sedge	1%
<i>Oryzopsis racemosa</i>	Black-seeded rice grass	<1%

Boxelder - green ash disturbed native forest (Southeast portion) – 6.82 acres



This area of forest follows a ravine that leads from the picnic area of Cherokee Park, under Cherokee Heights Boulevard and down about 2/3 of the bluff. From the ravine, the boxelder-green ash disturbed native forest continues northeast along a moist plateau.

Sugar maple, ironwood, green ash, boxelder, and basswood dominate the forest canopy of the ravine. About ten percent of the shrub layer of the forest is occupied by common buckthorn. Other invasive species present during the inventory include Siberian elm, Tartarian honeysuckle and reed canary grass. In areas where buckthorn is not dominant and where trampling is less severe, ground layer sedges and forbs occur. The most common vine and ground layer species include greenbriar, Virginia creeper, Virginia waterleaf, and zigzag goldenrod. Wild ginger is also common in a portion of the ravine just north of Cherokee Heights Boulevard.

The head of the ravine closest to the picnic area is heavily used and contains few ground layer species due to trampling. The ravine also has a high degree of disturbance north of Cherokee Heights Boulevard near the location of the brick yard operation. Some adjacent slopes are still experiencing erosion due to the operation while others are re-vegetating.

Canopy trees

<i>Latin Name</i>	<i>Common Name</i>	<i>Percent Cover</i>
<i>Acer negundo</i>	Boxelder	25%
<i>Carya cordiformis</i>	Bitternut hickory	<1%
<i>Celtis occidentalis</i>	Hackberry	6%
<i>Fraxinus pennsylvanica</i>	Green ash	8%
<i>Juglans nigra</i>	Black walnut	<1%
<i>Pinus strobus</i>	Eastern white pine	<1%
<i>Populus deltoides</i>	Cottonwood	10%
<i>Quercus alba</i>	White oak	18%
<i>Quercus bicolor</i>	Swamp white oak	<1%
<i>Quercus rubra</i>	Red oak	3%
<i>Salix nigra</i>	Black willow	<1%
<i>Tilia americana</i>	American basswood	25%
<i>Ulmus americana</i>	American elm	5%

Mid-Story trees

<i>Ostrya virginiana</i>	Ironwood	2%
<i>Prunus americana</i>	Choke cherry	2%
<i>Prunus americana</i>	Choke cherry (purple cultivar)	<15%
<i>Rhamnus cathartica</i>	Common buckthorn	3%
<i>Ulmus pumila</i>	Siberian elm	1%

Shrub layer

<i>Cornus foemina</i>	Grey dogwood	<1%
<i>Cornus sericea</i>	Red-osier dogwood	15%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	4%
<i>Rhus typhina</i>	Staghorn sumac	<1%
<i>Ribes cynosbati</i>	Gooseberry (no thorns)	<1%
<i>Rosa sp.</i>	rose	<1%
<i>Sambucus sp.</i>	Elderberry	<1%
<i>Viburnum lantana</i>	Wayfaring bush	<1%
<i>Viburnum lentago</i>	Nannyberry	<1%
<i>Viburnum rafinesquianum</i>	Arrowwood viburnum	<1%

Groundlayer vines

<i>Rhus toxicodendron</i>	Poison ivy	<1%
<i>Parthenocissus inserta</i>	Virginia creeper	<1%

Forbs or seedling trees

<i>Anemone virginiana</i>	Cylindrical thimbleweed	<1%
<i>Apocynum androsaemifolium</i>	Dogbane	<15
<i>Aquilegia canadensis</i>	Columbine	<1%
<i>Arctium minus</i>	Burdock	<1%
<i>Arisaema atrorubens</i>	Jack in the pulpit	<1%
<i>Asarum canadense</i>	Wild ginger	2%
<i>Asparagus officinalis</i>	Asparagras	<1%
<i>Aster cordifolius</i>	Heart leaved aster	<1%
<i>Desmodium glutinosum</i>	Pointed leaved tick trefoil	<1%
<i>Equisetum sp.</i>	Equisetum	<1%

<i>Eupatorium rugosum</i>	White snakeroot	<1%
<i>Helianthus divaricata</i>	Woodland sunflower	<1%
<i>Hydrophyllum virginianum</i>	Virginia waterleaf	<1%
<i>Laportea canadensis</i>	Wood nettle	<1%
<i>Monarda fistulosa</i>	Wild bergamot	<1%
<i>Rubus spp.</i>	Raspberry	<1%
<i>Scirpus atrovirens</i>	Green bulrush	<1%
<i>Solanum nigrum</i>	Black nightshade	<15
<i>Solidago canadensis</i>	Giant goldenrod	<1%
<i>Solidago flexicaulis</i>	Zig-zag goldenrod	<1%
<i>Taraxacum officinale</i>	Dandelion	<1%
<i>Violet sp.</i>	Violet	<1%

Grasses and sedges

<i>Carex blanda</i>	Woodland sedge	<1%
<i>Carex eburina</i>	A species of sedge	<1%
<i>Carex pennsylvanica</i>	Pennsylvania sedge	<1%
<i>Leersia oryzoides</i>	Rice-cut grass	<1%
<i>Phalaris arundinacea</i>	Reed canary grass	<1%
<i>Typha latifolia</i>	Cattail	<1%

Lowland hardwood forest - .48 acres



This small area of lowland hardwood forest is located in an area of disturbed soil. The soil is generally mounded and contains a large amount of rock and wood, indicating that it was material dumped at the site. Green ash and American elm dominate the canopy. Common buckthorn makes up about 10% of the shrub layer. Red-osier dogwood, gooseberry and chokecherry are other common shrubs in the relatively dense shrub layer. The generally sparse ground layer is dominated by

Aster spp. and riverbank grape.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Fraxinus pennsylvanica</i>	Green ash	75%
<i>Ulmus americana</i>	American elm	15%

Mid-Story trees

<i>Prunus americana</i>	Choke cherry	1%
<i>Rhamnus cathartica</i>	Common buckthorn	10%

Shrub layer

<i>Cornus sericea</i>	Red-osier dogwood	40%
-----------------------	-------------------	-----

<i>Ribes Missouriense</i>	Missouri gooseberry	2%
<i>Ribes sp.</i>	Gooseberry	1%

Groundlayer vines

<i>Vitis riparia</i>	Riverbank grape	5%
----------------------	-----------------	----


Forbs or seedling trees

<i>Arctium minus</i>	Burdock	<1%
<i>Aster sp.</i>	Aster	<1%
<i>Rubus spp.</i>	Raspberry	<1%

Grasses and sedges

<i>None</i>		
-------------	--	--

Mixed emergent marsh (Northern portion) - .81 acres



This mixed emergent marsh is quickly becoming a willow swamp. The marsh is dominated by the invasive species, reed canary grass. Reed canary grass forms dense stands that spread readily by rhizomes and seed. More desirable species such as wild iris, lake sedge, smartweed, river bulrush, and Joe-pye weed are present. Sandbar willow is the dominant shrub in the marsh and it is spreading quickly, currently covering about

20% of the marsh. Red-osier dogwood is another shrub species that is present in the marsh. The shrubs may be spreading due to a change in hydrology within the marsh. In areas where the willows have become thick, the reed canary grass is less robust. A powerline runs along the northern edge of the marsh. Trimming of trees and shrubs likely occurs in the power line right-of-way.

Canopy trees

<i>Latin Name</i>	<i>Common Name</i>	<i>Percent Cover</i>
<i>None</i>		

Mid-Story trees

<i>None</i>		
-------------	--	--

Shrub layer

<i>Cornus serotina</i>	Red-osier dogwood	2%
<i>Salix exigua</i>	Sandbar willow	15%

Groundlayer vines

None		
------	--	--

Forbs or seedling trees

<i>Arctium minus</i>	Burdock	<1%
<i>Asclepias incarnata</i>	Swamp milkweed	<1%
<i>Eupatorium maculatum</i>	Joe-pye weed	4%
<i>Hibiscus palustris</i>	Hibiscus	5%
<i>Iris versicolor</i>	Wild iris	<1%
<i>Polygonatum sp.</i>	Smartweed	1%
<i>Urtica dioica</i>	Stinging nettle	<1%

Grasses and sedges

<i>Carex lacustris</i>	Lake sedge	<1%
<i>Elymus canadensis</i>	Canada wild rye	<1%
<i>Phalaris arundinacea</i>	Reed canary grass	90%
<i>Scirpus fluviatilis</i>	River bulrush	1%
<i>Typha angustifolia</i>	Narrow leaved cattail	3%

Mixed emergent marsh (Southern portion) - .70 acres



This area of mixed emergent marsh is at the southern end of the site and is at the northern end of Pickerel Lake. The wetland is dominated by narrow-leaved cattail but also contains other native species typical of mixed emergent marsh including giant bur-reed, river bulrush, lake sedge, sandbar willow, smartweed and iris. Invasive species include reed canary grass and purple loosestrife. A hibiscus species not native to the area is very common in the wetland with a

cover of about 20%. The wetland seems to receive a significant amount of nutrients and fluctuating water levels, which have led to low diversity, and species that can handle these conditions. About 20% of the entire northeast portion of the wetland is dominated by sandbar willow.

Canopy trees

Latin Name	Common Name	Percent Cover
None		

Mid-Story trees

<i>Ulmus pumila</i>	Siberian elm	<1%
---------------------	--------------	-----

Shrub layer

<i>Cornus serotina</i>	Red-osier dogwood	2%
------------------------	-------------------	----

<i>Salix exigua</i>	Sandbar willow	20%
---------------------	----------------	-----

Groundlayer vines

<i>Parthenocissus inserta</i>	Virginia creeper	<1%
<i>Vitis riparia</i>	Riverbank grape	<1%

Forbs or seedling trees

<i>Anemone canadensis</i>	Canada anemone	<1%
<i>Aster sp.</i>	Aster	<1%
<i>Equisetum sp.</i>	Equisetum	2%
<i>Hibiscus palustris</i>	Hibiscus	20%
<i>Impatiens capensis</i>	Jewelweed	<1%
<i>Iris versicolor</i>	Wild Iris	<1%
<i>Lycopus sp.</i>	Bugleweed	<1%
<i>Lythrum salicaria</i>	Purple loosestrife	<1%
<i>Pilea pumila</i>	Clearweed	<1%
<i>Polygonatum sp.</i>	Smartweed	3%

Grasses and sedges

<i>Carex lacustris</i>	Lake sedge	1%
<i>Phalaris arundinacea</i>	Reed canary grass	15%
<i>Scirpus atrovirens</i>	Green bulrush	1%
<i>Sparganium eurycarpum</i>	Giant burreed	1%
<i>Spartina pectinata</i>	Prairie cord grass	<1%
<i>Typha angustifolia</i>	Narrow leaved cattail	30%

Willow swamp - .34 acres



This area of willow swamp is connected to the two areas of mixed emergent marsh at the project site (one to the north and one to the south). The willow swamp has many species in common with the emergent marshes but has more area of open water. Sandbar willow is the dominant shrub species, covering about 15% of the swamp. Reed canary grass dominates the ground layer with a cover of about 30% but clearweed, wild iris, river bulrush and

germander are also present. The exotic hibiscus found in the emergent marsh (southern portion) is also found in the willow swamp.

Canopy trees

Latin Name	Common Name	Percent Cover
None		

Mid-Story trees

None		
------	--	--

Shrub layer

<i>Cornus sericea</i>	Red-osier dogwood	5%
<i>Salix exigua</i>	Sandbar willow	15%

Groundlayer vines

None		
------	--	--


Forbs or seedling trees

<i>Cirsium canadensis</i>	Canada thistle	<1%
<i>Hibiscus palustris</i>	Hibiscus	20%
<i>Impatiens capensis</i>	Jewelweed	1%
<i>Iris versicolor</i>	Wild iris	<1%
<i>Lycopus sp.</i>	Bugleweed	<1%
<i>Lythrum salicaria</i>	Purple loosestrife	<1%
<i>Pilea pumila</i>	Clearweed	<1%
<i>Teucrium canadense</i>	Germander	<1%
<i>Urtica dioica</i>	Stinging nettle	<1%

Grasses and sedges

<i>Phalaris arundinacea</i>	Reed canary grass	30%
<i>Scirpus fluviatilis</i>	River bulrush	10%

Floodplain forest - 2.68 acres



The floodplain forest within the project site is dominated by silver maple. The silver maple covers about 90% of the canopy. Other canopy species include green ash, black willow, hackberry, cottonwood and red oak. The invasives, common buckthorn and Tartarian honeysuckle, are the dominant shrubs. The ground layer is rather sparse but desirable native species that are present include, clearweed, Virginia wild rye, riverbank grape,

Equisetum spp., sedges, rice-cut grass and *Aster spp.* The floodplain forest is adjacent to an area of boxelder-green ash disturbed forest where a significant amount of regrading occurred. However, the floodplain forest appears largely intact with some mature tree species.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Acer saccharinum</i>	Silver maple	90%
<i>Carya cordiformis</i>	Bitternut hickory	<1%
<i>Celtis occidentalis</i>	Hackberry	1%
<i>Fraxinus pennsylvanica</i>	Green ash	5%
<i>Populus deltoides</i>	Cottonwood	10%
<i>Quercus rubra</i>	Red oak	<1%
<i>Salix nigra</i>	Black willow	10%

Mid-Story trees

<i>Rhamnus cathartica</i>	Common buckthorn	3%
---------------------------	------------------	----

Shrub layer

<i>Amorpha fruticosa</i>	Indigobush	<1%
<i>Cornus sericea</i>	Red-osier dogwood	10%
<i>Xanthoxylum americanum</i>	Prickly ash	<1%

Groundlayer vines

<i>Vitis riparia</i>	Riverbank grape	<1%
----------------------	-----------------	-----

Forbs or seedling trees

<i>Arctium minus</i>	Burdock	<1%
<i>Aster ontarionis</i>	Ontario aster	1%
<i>Catalpa speciosa</i>	Catalpa	<1%
<i>Equisetum sp.</i>	Horsetail	1%
<i>Pilea pumila</i>	Clearweed	<1%

Grasses and sedges

<i>Carex tribuloides</i>	A species of sedge	<1%
<i>Elymus virginica</i>	Virginia wild rye	1%

Dry prairie/savanna - sand gravel subtype (including prairie edges) - .62 acres

The prairie at the project site is currently a little less than an acre in size yet exhibits good diversity with around 50 native prairie species. It is quickly being invaded by trees and shrubs and potentially will turn into forest within twenty years without management. Both native and non-native species are invading the prairie with prickly ash, green ash, common buckthorn and staghorn sumac being common. There are signs that the area of prairie experienced sluffing in the past, which may have kept it open. In addition there are reports of the prairie burning periodically over the past century. Since it appears that the prairie was adjacent to savanna, there was probably a sufficient seed source for its re-establishment after sluffing occurred. A trail currently traversing the prairie is a significant threat to the long-term existence of this rare native plant community. The trail has caused considerable erosion and appears to be widening and becoming deeper. It is a



popular trail for the neighborhood residents so an alternate route will be necessary if the trail is closed.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Quercus macrocarpa</i>	Bur oak	10%
<i>Quercus rubra</i>	Red oak	5%

Mid-Story trees

<i>Betula papyrifera</i>	Paper birch	2%
<i>Crataegus spp.</i>	Hawthorn	1%
<i>Fraxinus pennsylvanica</i>	Green ash	3%
<i>Ulmus americana</i>	American elm	2%

Shrub layer

<i>Cornus foenea</i>	Gray dogwood	1%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	3%
<i>Rhamnus cathartica</i>	Common buckthorn	2%
<i>Rhus glabra</i>	Smooth sumac	3%
<i>Symphoricarpos occidentalis</i>	Western Snowberry (Buck brush)	1%
<i>Viburnum rafinesquianum</i>	Arrowwood viburnum	1%

Groundlayer vines

<i>Celastrus scandens</i>	Bittersweet	1%
<i>Vitis riparia</i>	River bank grape	1%

Forbs or seedling trees


<i>Aquilegia canadensis</i>	Wild columbine	<1%
<i>Amorpha canescens</i>	Leadplant	1%
<i>Anemone cylindrica</i>	Cylindrical thimbleweed	1%
<i>Apocynum cannabinum</i>	Dogbane	1%
<i>Aster ericoides</i>	Heath aster	1%
<i>Aster oblongifolius</i>	Aromatic aster	1%

<i>Campanula rotundifolia</i>	Harebell	1%
<i>Comandra umbellata</i>	Bastard (star) toadflax	1%
<i>Coreopsis palmata</i>	Prairie coreopsis	1%
<i>Dalea candida</i>	White prairie clover	1%
<i>Dalea purpurea</i>	Purple prairie clover	1%
<i>Desmodium canadense</i>	Showy tick trefoil	1%
<i>Eupatorium rugosum</i>	Black snakeroot	1%
<i>Euphorbia esula</i>	Flowering spurge	1%
<i>Fragaria virginiana</i>	Common strawberry	1%
<i>Galium boreale</i>	Northern bedstraw	1%
<i>Helianthus divaricatus</i>	Woodland sunflower	2%
<i>Heliopsis helianthoides</i>	Common oxeye	1%
<i>Lactuca canadensis</i>	White lettuce	1%
<i>Maianthemum stellatum</i>	False solomon's seal	1%
<i>Melilotus alba</i>	White sweet clover	1%
<i>Melilotus officinalis</i>	Yellow sweet clover	1%
<i>Mirabilis nyctaginea</i>	Wild four-o'clock	1%
<i>Monarda fistulosa</i>	Wild bergamot	1%
<i>Ostrya virginiana</i>	Ironwood	1%
<i>Physalis virginiana</i>	Ground cherry	1%
<i>Potentilla arguta</i>	Tall cinquefoil	1%
<i>Pycnanthemum tenuifolium</i>	Narrow-leaved mountain mint	1%
<i>Ratibida pinnata</i>	Grey-headed coneflower	1%
<i>Sanguinaria canadensis</i>	Bloodroot	<1%
<i>Solidago canadensis</i>	Canada goldenrod	1%
<i>Solidago hispida</i>	Hairy goldenrod	1%
<i>Solidago nemoralis</i>	Gray goldenrod	1%
<i>Solidago rigida</i>	Stiff goldenrod	1%
<i>Taraxacum officinale</i>	Dandelion	1%
<i>Tilia americana</i>	Basswood	<1%
<i>Ulmus americana</i>	American elm	1%
<i>Uvularia grandiflora</i>	Large flowered bellwort	<1%
<i>Veronicastrum virginicum</i>	Culver's root	1%

Grasses and sedges

<i>Andropogon gerardii</i>	Big bluestem	2%
<i>Bouteloua curtipendula</i>	Sideoats grama	3%
<i>Carex blanda</i>	Woodland sedge	1%
<i>Carex eburina</i>	A species of sedge	1%
<i>Elymus canadensis</i>	Canada wild rye	3%
<i>Panicum oligosanthos var. scriberianum</i>	Scribner's panic grass	1%
<i>Panicum sp.</i>	Panic grass	1%
<i>Poa pratensis</i>	Kentucky bluegrass	3%
<i>Schizachyrium scoparium</i>	Little bluestem	60%
<i>Sorghastrum nutans</i>	Indiangrass	40%

Oak forest - 17.70 acres



Oak forest makes up a large portion of the project area. Species composition within the oak forest seems to vary considerably between ravines and ridges. Ravines contain many sugar maples and have many species in common with maple-basswood forests while ridges often have relatively widely spaced oak trees and have some characteristics of oak savanna. Overall, red oaks and sugar maple are the dominant canopy trees with 30% cover for each. American basswood, hackberry, white oak, green ash and bur oak are also common. Due to the large number of trees common to maple-basswood forests it appears that the oak forest may be making a successional change to maple basswood-forest. The most prevalent invasive species in the forest are common buckthorn (20% cover) and Tartarian honeysuckle (2% cover).



Erosion is a serious problem in several portions of the oak forest. One large ravine is experiencing serious erosion and other small ravines are eroding to a lesser degree. The Ramsey Soil and Water Conservation District is aware of the ravine with severe erosion and is planning future stabilization efforts. There are many trails within the oak forest and some are causing significant erosion. The trails are also causing soil compaction and habitat fragmentation. A trail plan should be encouraged for the park to determine which trails can be closed, relocated or stabilized.

Canopy trees

Latin Name	Common Name	Percent Cover
<i>Acer platanoides</i>	Norway maple	<1%
<i>Acer saccharum</i>	Sugar maple	30%
<i>Betula papyrifera</i>	Paper birch	<1%
<i>Carya cordiformis</i>	Bitternut hickory	1%
<i>Celtis occidentalis</i>	Hackberry	7%
<i>Fraxinus pennsylvanica</i>	Green ash	15%
<i>Juglans cinerea</i>	Butternut	3%
<i>Juglans nigra</i>	Black walnut	<1%
<i>Populus deltoides</i>	Cottonwood	3%
<i>Prunus serotina</i>	Black cherry	1%

<i>Quercus alba</i>	White oak	8%
<i>Quercus ellipsoidalis</i>	Northern pin oak	4%
<i>Quercus macrocarpa</i>	Bur oak	5%
<i>Quercus rubra</i>	Red oak	25%
<i>Tilia americana</i>	American basswood	20%
<i>Ulmus americana</i>	American elm	2%

Mid-Story trees

<i>Amelanchier laevis</i>	Smooth serviceberry	<1%
<i>Ostrya virginiana</i>	Ironwood	15%
<i>Prunus americana</i>	Choke cherry	8%
<i>Rhamnus cathartica</i>	Common buckthorn	20%
<i>Sorbus sp.</i>	Mountain ash	<1%

Shrub layer

<i>Cornus alternifolia</i>	Pagoda dogwood	<1%
<i>Cornus foemina</i>	Grey dogwood	2%
<i>Cornus rugosa</i>	Round leaved serviceberry	1%
<i>Lonicera tartarica</i>	Tartarian honeysuckle	2%
<i>Rhus glabra</i>	Smooth sumac	>1%
<i>Rhus typhina</i>	Staghorn sumac	<1%
<i>Ribes cynosbati</i>	Prickly gooseberry	<1%
<i>Ribes missouriensis</i>	Gooseberry	<1%
<i>Rosa blanda</i>	Wild rose	<1%
<i>Sambucus pubens</i>	Red berried elder	<1%
<i>Staphylea trifolia</i>	Bladdernut	<1%
<i>Symphoricarpos occidentalis</i>	Wolfberry	<1%
<i>Symphoricarpos occidentalis</i>	Wolfberry	<1%
<i>Viburnum lentago</i>	Nannyberry	<1%
<i>Viburnum rafinesquianum</i>	Arrowwood viburnum	1%
<i>Xanthoxylum americanum</i>	Prickly ash	2%

Groundlayer vines

<i>Amphicarpa bracteata</i>	Hog peanut	2%
<i>Celastrus scandens</i>	Bittersweet	<1%
<i>Parthenocissus inserta</i>	Virginia creeper	<1%
<i>Rhus toxicodendron</i>	Poison ivy	<1%
<i>Vitis riparia</i>	Riverbank grape	1%

Forbs or seedling trees

<i>Actaea rubra</i>	Baneberry	<1%
<i>Anemone virginiana</i>	Cylindrical thimbleweed	<1%
<i>Anamonella thalictroides</i>	Rue anemone	<1%
<i>Aquilegia canadensis</i>	Columbine	<1%
<i>Aralia nudicaulis</i>	Wild sarsaparilla	<1%
<i>Arctium minus</i>	Burdock	<1%
<i>Arisaema atrorubens</i>	Jack in the pulpit	<1%
<i>Asarum canadense</i>	Wild ginger	<1%
<i>Aster prenanthoides</i>	Crooked stem aster	<1%
<i>Campanula rotundifolia</i>	Harebell	<1%
<i>Desmodium glutinosum</i>	Pointed leaved tick trefoil	<1%

<i>Echinocystis lobata</i>	Wild cucumber	<1%
<i>Eupatorium rugosum</i>	White snakeroot	<1%
<i>Fragaria virginiana</i>	Wild strawberry	<1%
<i>Galium boreale</i>	Northern bedstraw	<1%
<i>Geranium maculatum</i>	Wild geranium	<1%
<i>Helianthus divaricatus</i>	Woodland sunflower	<1%
<i>Hepatica acutiloba</i>	Sharp lobed hepatica	<1%
<i>Impatiens spp</i>	Jewelweed	<1%
<i>Lactuca canadensis</i>	White lettuce	<1%
<i>Maianthemum canadense</i>	Wild lily-of-the-valley	<1%
<i>Maianthemum stellatum</i>	False Solomon's seal	<1%
<i>Melilotus alba</i>	Yellow sweet clover	<1%
<i>Melilotus officinale</i>	White sweet clover	<1%
<i>Oxalis spp.</i>	Wood sorrel	<1%
<i>Rubus spp.</i>	Raspberry	<1%
<i>Sanguinaria canadense</i>	Bloodroot	<1%
<i>Smilax rotundifolia</i>	Greenbriar	<1%
<i>Solidago flexicaulis</i>	Zig-zag goldenrod	<1%
<i>Solidago gigantea</i>	Giant goldenrod	<1%
<i>Thalictrum dioicum</i>	Woodland meadowrue	<1%
<i>Uvularia grandiflora</i>	Large-flowered bellwort	<1%


Grasses and sedges

<i>Carex blanda</i>	Woodland sedge	<1%
<i>Carex eburina</i>	A species of sedge	5%
<i>Carex pennsylvanica</i>	Pennsylvania sedge	3%
<i>Carex sprengelii</i>	Sprengel's sedge	<1%
<i>Hystrix patula</i>	Bottlebrush grass	<1%
<i>Juncus tenuis</i>	Path rush	<1%
<i>Oryzopsis racemosa</i>	Black-seeded rice grass	<1%
<i>Poa pratensis</i>	Kentucky bluegrass	<1%


Appendix D: Earthworms

There are many species of earthworms found in North America, both native and exotic. Severe infestations of exotic earthworms damage woodland and forest ecosystems by consuming the humus layer of the forest floor changing its structure, composition and function. Below is an excerpt from the web site for the Minnesota Worm Watch at the University of Minnesota in Duluth.

(From Minnesota Worm Watch, 2002 - 2003, University of Minnesota Duluth, www.nrri.umn.edu/worms/Default.htm) Photo credits University of Minnesota Agricultural Experiment Station



Without earth worms:



Forest structure without worms

The structure of a woodland or forest is determined by several layers of plants: the **canopy** layer is made up of the tallest trees, the **subcanopy** is composed of shorter tree species and tree saplings, the **understory** contains most of the visible plant life found between the sapling layer and forest floor. The **forest floor** is where one would find the roots, bulbs, fungi, seeds, years of accumulated leaves and twigs. Hardwood trees produce tons of **leaf litter** each year, which is high in nutrients. This litter is **decomposed** by bacteria and fungi in the forest floor. The combination of high **productivity** and slow **decomposition** results in the development of a thick forest floor with a unique set of **soil layers** beneath.

The understory is sometimes understood as anything below the canopy. However, the definition used here includes all plants other than tree species and they usually occupy the area from the forest floor to about 6 feet up. We find the following at this level:

- **Shrubs** are woody species of plants that do not grow into trees. They tend to grow as small to medium sized bushes. There are many shrubs that grow in hardwood forests with some of the more familiar being Raspberry, Gooseberry and Hazelnut.



Understory without worms

- **Herbaceous plants** include grasses and grass-like plants, ferns, flowers and all other non-woody plant species that grow in the forest. Among the herbaceous plants, there are several categories as follows:
 - **Spring ephemerals** begin growing very early in the spring to take advantage of the sun breaking through the leafless trees. Spring

ephemerals will grow, flower, produce seeds, and die back by the time the trees start budding and the summer plants start coming up.

- **Annuals** are plants that grow, produce seeds and die in the same year.
- **Biennials** take two years to grow to maturity, produce seeds and die.
- **Perennials** may take two or more years to grow to maturity, produce seeds every year or only occasionally, and continue to grow year after year.
- **Mosses** are common in hardwood forests and are different than herbaceous plants because they have no vascular tissue. **Vascular tissue** inside the stems of plants pumps nutrients and water up from the roots to leaves. Mosses transfer nutrients and water from one cell to another. However, this process is limited by gravity, which explains why moss is found growing low to the ground in moist places.

When looking at how an ecosystem functions, one component that sometimes gets overlooked is what happens IN the ground. The tendency is to take notice of the plants and animals that are above ground as defining the system. However, the soil and forest floor are two of the most important aspects of a hardwood forest ecosystem because they are the foundation on which all life above ground depends. For example, root systems are the foundation of most species of plants. Plants get their nutrients and water from the soil and forest floor through



Soil without worms

their root systems. The roots also anchor the plants in the forest floor or soil. A given plant community depends upon a specific soil. A change in the soil can dramatically change the plants that make up that community.

While we may not have paid much attention to the soil, there have been soil scientists studying different soils all over the world for hundreds of years. One of the interesting things they discovered is that as time passes, soils form layers and that each layer has different characteristics and functions in the ecosystem. The layers in a hardwood forest ecosystem can be broken down into the following:

- The **O horizon** is the layer that makes up the forest floor. This layer is composed of fresh and partially decomposed litter that has accumulated over many years. The litter contains twigs, leaves, seeds, bark, and wood from small fragments to large logs. In the hardwood forests of the Great Lakes region, the O horizon can be up to 10 cm (4 inches) thick. This layer is full of organisms and is very important to the overall functioning of the ecosystem.
- The **A horizon** is a thin layer just below the O horizon in hardwood forests and is considered the top layer of soil. This layer is usually 1 centimeter or less in thickness and a very dark brown or black in color. The color comes from the decomposed litter that is no longer distinguishable, much like the soil that comes out of a compost pile. This is what gardeners and farmers might call "good, black dirt."
- The **E horizon** develops beneath the A horizon. The total thickness may be 10 to 20 centimeters. It is composed of soil deposited both during the retreat of the glaciers and before it was covered by forest. This soil may contain various amounts of clay, sand, silt and rocks. The top of the E horizon is dark black or gray in color and gets lighter in color as soil depth increases. The dark color comes from organic molecules carried down from the A horizon. This process is called leaching.
- The **B horizon** is below the E horizon and is composed of the same material. This layer can be very thick or thin depending on the site and is usually some shade of yellow, brown or red coloring. The coloring comes from the natural color of the soil as it was deposited but also can be affected by dissolved molecules of iron or salts that leach down with rainwater.
- The **C horizon** is below the B horizon and is made up of the same material as E and B horizons but has not been changed by leaching and is virtually identical to what would have been seen after glacial retreat. Because of this, it is often referred to as "parent material."
- **Mineral soil** is a general term that often refers to the E, B & C horizons collectively. These lower layers of soil that have been little changed from the nature of soil that was deposited by glaciers or by rivers and lakes since glacial retreat. The most important distinction is that mineral soil doesn't contain much decomposed litter. So, the color is usually much lighter than the black color of the A horizon.

The forest floor and top layer of soil (the O and A horizons, respectively) are found between the vegetation and the mineral soil. It is the centerpiece of the hardwood forest ecosystem. These two horizons are where most of the **nutrient cycling** takes place and where all the plants **germinate** and grow. One important characteristic of soils is their bulk density. In hardwood forests, the forest floor and upper soil have low **bulk density**, meaning they are very loose and spongy so roots can grow easily through them. The forest floor and upper soil also hold a lot of moisture. The combination of moisture and shade from the canopy create a generally cool **microclimate**, which is an important factor in a hardwood forest ecosystem. A microclimate refers to the unique temperature and moisture conditions created in a small space due to influences of plants, which can be very different than temperature and moisture conditions in open spaces nearby. Many plants and animals rely on this microclimate for their survival.

In winter, the forest floor acts as a blanket that helps protect organisms from freezing conditions. Most of the plants and animals living in this layer have adapted to survive and grow in the particular conditions of the forest floor. Big changes in this layer could mean big changes for all the organisms that depend on a stable hardwood forest ecosystem. Let's take a look at some of the components of the forest floor:

- **Logs** fall to the forest floor and decompose very slowly. Dead fall, or logs, contain a great deal of nutrients and are home to a number of insects, fungi, and bacteria. The older and more rotted logs often have a layer of moss and other plants growing on them. As a log ages and decomposes, it sinks deeper and deeper into the forest floor, providing habitat for amphibians such as salamanders and small mammals like red-backed voles. The log not only provides protection and moisture for these animals but also food in the form of insects and fungi. A log can be a habitat onto itself for some creatures.
- **Plant roots** grow densely in the forest floor because of the high concentration of nutrients and its loose spongy texture. Very few roots extend below the forest floor. The ones that do are usually for anchoring large plants as opposed to taking up nutrients and water. Besides the fine roots used to take up nutrients, many forest plants also use their roots to store food and reproduce. Perennial plants, for example, store food in different kinds of fleshy roots called bulbs, rhizomes, or corms. As they grow the bulbs or rhizomes will spread and divide, growing new plants. This process is called **vegetative reproduction**.
- **Fungi** grow densely in the forest floor and there are more species than have been identified. Fungi are not green because they don't have chlorophyll and therefore do not produce their own food through **photosynthesis**. Instead, fungi eat dead plant material. Mushrooms growing on a dead log are an example of this.

There are some fungi that don't get enough food through this process so they work with green plants by attaching themselves to the roots. The fungi form an extensive network of root-like strings, called hyphae, spreading out from the plant roots. This relationship is mutually beneficial because the fungi provide more nutrients and water to the plant and the plant, in exchange, provides extra carbohydrates (made through photosynthesis) to the fungi. Fungi that work in this **symbiotic relationship** are called **mycorrhizae**. There are some hardwood forest

plants that have a hard time absorbing enough food from the forest floor because their roots are very thick, not dense and hairy like grass roots. Plants such as these depend upon mycorrhizae fungi for survival. An example of this would be many ORCHID species. However, most hardwood forest plants, including the trees, have mycorrhizae associated with their roots. Without the presence of mycorrhizae, the diversity of plants that make up the understory would be dramatically reduced.

- **Seeds** are deposited by plants into the forest floor. Because the forest floor is made up of loose, organic material such as leaf litter, many of the seeds are protected from **predation** by small mammals and birds. Also, because the forest floor is moist and full of nutrients, the seeds have a perfect place to germinate. The forest floor and the protection it provides is especially important for some herbaceous plants because their seeds germinate slowly, taking two or more years to develop into a small plant. If not protected from predators or from drying out over a long period of time, the seeds would have no chance of growing into a plant.
- **Leaves and twigs** fall to the forest floor creating a thick layer on top of the soil. The youngest leaves on top are typically brown and easy to identify. However, as one goes deeper, the leaves turn black and are broken apart making identification difficult or impossible. This is due to the work of **bacteria** and fungi, critical partners in the nutrient cycling process.

If one has a compost pile for yard leaves, grass, and vegetable kitchen scraps, then decomposition is a familiar occurrence. If not, most people have picked up or kicked a pile of leaves. The top leaves are dry and easily identifiable. However, the bottom may be moist and black in color. There are probably hundreds of different kinds of bacteria and perhaps millions living in a single handful of leaf litter.

Everything in the forest is a source of nutrients. However, only the nutrients in the forest floor and upper soil are being broken down so they are FREE to be taken up by plants. Going back to our compost example, if vegetable kitchen scraps are thrown into a compost pile and allowed to be broken down by fungi and bacteria, the black, organic mixture that results can be applied to the garden and the plants will respond by taking up the available nutrients. However, if the scraps were thrown directly into the garden, the garden plants would not be able to take up the nutrients. In fact, the scraps may sit on the soil for some time before they break down. The nutrients exist within the scraps but are not available or free to be taken up by the garden plants. In other words, if it weren't for bacteria and fungi, the nutrients in the forest would not get broken down and eventually the forest would run out of nutrients. If that were to happen, then plants could no longer grow and survive!

In a hardwood forest floor, the composting process is controlled by the fungi and bacteria. The nutrients are slowly released over time and taken up by living plants

as fast as the nutrients are produced. Because of this balance between nutrient release and plant absorption, there is little to no loss of nutrients from the system.

Critters are diverse and numerous in hardwood forests. From Moose to spiders, many creatures use the understory for **habitat**.

- **Animals** living in the forest floor and upper soil are numerous. Hundreds of microscopic animals like protozoa, nematodes, flatworms, and water bears (a tiny animal living in the water film found on the surface of leaves, mosses, and leaf litter - phylum tardigrada) live in this narrow portion of ground. There are also dozens of land snails and spiders such as Orb Weaver spiders that spin large "orb" webs between trees and branches, "Jumping" spiders that pounce on prey, and "Forest Wolf" spiders which do not spin webs but rather burrow in the forest floor and upper soil and hunt at night.
- **Insects** crawl and fly through the understory eating plants and each other. Insects play an important role by pollinating plant life in the forest and surrounding areas. They are also a food source for many birds and some mammals. Ants, beetles, butterflies, flies, bees, and wasps are all important pollinators for hardwood forest plant life. For example, "Ichneumonid" wasps rely on rotting logs for food and shelter. These wasps lay their eggs on the larvae of other insects living in logs. Without the rotting logs, it would be difficult for these wasps to survive. In fact, bees, ants, beetles, and wasps all rely on rotting logs for survival. A rotting log is a crowded place!
- **Birds** that winter in South and Central America fly thousands of miles to nest and raise young in the hardwood forests of Minnesota and the Great Lakes region. For example, the ovenbird makes its home in the understory of hardwood forests. Ovenbirds build their nest in the thick forest floor. Their nest of leaves, moss and twigs always has a roof so the entrance to the nest is just a tiny slit. Ovenbirds eat a diverse diet of insects, spiders, snails and seeds that they find in the forest floor.
- **Mammals** of all sizes inhabit and make use of hardwood forests. Raccoons, white-tailed deer and bear are just some of the mammals that use the FOREST understory for both cover and a source of food. Bear for example, find hazelnuts and raspberries a great source of food, as they put fat on for winter. Shrews and moles eat insects and tubers and nest under forest debris. The Eastern Chipmunk eats bulbs, fruit, seeds and insects and burrows underground. Deer Mice and Red-backed Voles eat insects, seeds, fruit and fungi, preferring damp conditions and nesting under forest litter, logs, and roots. White-tailed deer browse various plants including tree buds and leaves and many of the herbaceous plants. Eating plants, whether it is from an insect or mammal is called **herbivory**. Herbivory in a hardwood forest ecosystem usually has little impact on plant species in the understory. The total number of plant species is high as are the number of plants within each species. Because of this, the percent of total plants grazed is low, as is the impact due to grazing.
- **Amphibians and reptiles** live in the hardwood forest but are usually hard to see since they tend to make their homes inside or under old, rotting logs or in piles of rotting plant material. Salamanders, like the blue spotted salamander, are

especially adapted to the moist cool conditions of the forest floor. Salamanders do not have lungs and therefore must breathe through their skin, which must be wet for this to happen. They feed on insects and other small organisms that live IN(on) the forest floor. There are also several snake species living in the moist, cool and well-protected forest floor of Minnesota's hardwood forests. One example is the beautifully colored milk snake.

With earth worms:

The canopy and sub-canopy do not change much immediately after the worms invade. However, regeneration is very low after the worms invade. So as the canopy and sub-canopy trees age and begin to die, it is possible there will be few, if any, younger trees to replace them. However, because we haven't been able to study the long term effects of these worms on hardwood forest regeneration, it is unclear at this time what the exact effect will be.



Forest structure with worms

Tree Roots extend through all soil horizons, depending on the tree species. The large roots are primarily for anchoring the trees so they don't fall over and can extend a long way from the tree, deep into the soil horizons. The roots that take up water and nutrients are tiny, whitish roots the size of human hair or string, growing from little branches off the large roots. They tend to grow in the upper soil horizons where most of the water and nutrients are.



Understory with worms

Saplings have most of their roots in the forest floor. When earthworms invade, the first thing they do is eat all of the litter in the forest floor and mix it into the deeper soil layers. This activity both disturbs and exposes the sapling's roots. As a result, many of them die and fall over.

Seedlings also root in the forest floor and most die when the earthworms invade because the earthworms eat the forest floor right out from under their tiny roots. Where

previously there were 100 or more tree seedlings in a square meter, now there may only be 1 or 2 and in many areas none are left growing.

Most **native species** (species which are indigenous to a given ecosystem) that make up the understory do not survive after the invasion of earthworms. In a forest that previously had 20 to 40 native species, there may now only be one or just a few remaining. In addition, there is now very little plant cover, as little as 0-20% where there had been 100%. Most of the understory is now bare soil rather than a lush carpet of green plants.

Most **exotic species** (species not indigenous to a given ecosystem) we have in Minnesota came from Europe, including earthworms. This means that European plants have co-evolved with earthworms and are better adapted to living with them than our native species that evolved with no worms. In some forests, after the earthworms invade and the native species die back, some exotic species begin to invade and can start the process of taking over the understory.

Shrubs, Herbaceous plants and mosses all decrease after the earthworms invade. Like the tree saplings and seedlings, these plants had been rooting almost exclusively in the forest floor. When the earthworms eat the forest floor, the plant roots are left exposed. The microclimate (cool and moist) protected these root systems from warm and dry environment is not gone.



Soil with worms

after the worms invade.

We know that some of these native species can grow in soil containing worms because a lot of us grow them in our gardens and most gardens have earthworms. The difference is that in a garden, plant roots are put directly into the soil. In the forest, worms eat the forest floor so fast that most of the plants don't have a opportunity to get their roots into soil and thus die. It could be the case that if these native plants could establish themselves in the soil, they could recover

The first thing that Earthworms do when they invade a forest is to eat the O horizon. Within a matter of a few years (3-5), they can consume the whole layer of litter and all the organisms that live in it. In many forests, this layer is completely eliminated so that all that is left is bare soil with small piles of cast material by the entrance to the earthworm burrows. Each fall the trees deposit a new supply of leaves to the forest floor. The earthworms will eat some of these leaves in the fall before winter arrives and they become dormant. During spring and early summer, the worms can usually eat the rest of the litter so by late summer, only bare soil remains.

The *A horizon* was very thin before the earthworms arrived (1 cm), but now it gets very thick, between 10 and 15 centimeters. The soil that makes up this new horizon is composed of the earthworm casts produced after eating the litter. It is a dark black layer with earthworm burrows throughout. The original A horizon was kept loose and moist because of the amount of organic material present. The new A horizon lacks this organic material and therefore is compacted in comparison.

A new *E horizon* develops beneath the new A horizon. It looks pretty much the same as it did before, but now it is lower in the soil.

Earthworm burrows can be seen on the top of the soil and, if you were to dig a hole, all through the A horizon. Each kind of earthworm has its own type of burrow system. The small, reddish worms living in the litter and at the surface usually don't burrow down very far. However, they will create burrows along the surface underneath the litter or logs. If you dig under these a log for example, you will see their "tracks" that are usually 1-2 millimeters in diameter.

The large red worms (night crawlers) create large burrows that go almost straight down into the soil. You can see the holes at the surface of the soil, usually 3-4 millimeters in diameter, surrounded by a small pile of cast material called a **midden**. Night crawlers also line their burrow with cast material. To see this, cut a cross section of a burrow with a hand shovel. Each burrow is home to one night crawler so estimating the population can be done by counting the number of holes and middens in an area.

The whitish gray worms create branching burrows that wind through the A horizon. They are smaller than night crawler burrows, usually 1-2 millimeters in diameter. The burrow will come to the surface occasionally, typically under a log and may connect to night crawler burrows.

The *forest floor*, centerpiece of the hardwood forest ecosystem, has been radically changed and for all practical purposes is gone due to earthworms eating the O horizon. All of the processes that used to occur in the forest floor have been moved into the deeper soil layers. Many of the organisms that used to live in the forest floor have lost their habitat, including food sources. They will either leave or die trying to find another habitat they can live in. The loose, spongy layer of litter is now gone. Plant roots have a harder time growing in the new A horizon than they did in the O horizon. Without the forest floor to insulate the soil, it will get warmer and drier in the summer and colder in the winter. These conditions may make it difficult to survive for organisms that had adapted to the particular conditions of what was the forest floor.

Earthworms do not eat *logs* directly, but once the forest floor is gone, they can begin to dry out and get hard. The hard wood makes it difficult for insects to burrow into them and the log no longer provides the moist, protected habitat and food sources some animals need. The mosses and other plants that require moisture to survive may also die back if the log has dried out. If the log has not dried out, mosses will still grow in under the log for as long as moisture is present. Another exception is a **tip-up mound**, defined as the soil still clinging to the root system of downed tree. The effect earthworms have on this soil is limited, allowing for some plants and mosses to survive.

Plant roots do not grow as densely in the new A horizon as they did in the forest floor. As a result, the remaining plants may become stressed more easily when the weather turns warm and dry. There continues to be plenty of nutrients in the soil because of the nutrient rich casts left behind by the earthworms. However, some plants with poor root systems may not be able to get to the nutrients with the same efficiency. In addition, the tiny roots that plants use to absorb nutrients and water can easily get damaged by earthworms grazing around or on them. Earthworms many not want to eat the root itself,

but they like to eat the bacteria and fungi close to the roots. Earthworms can also cause damage to the bulbs, rhizomes or corms that native perennial plants use to store food. When these fleshy roots are damaged and the stored food is lost or used up, the plant can no longer divide and grow new plants through vegetative reproduction.

Fungi are a preferred food of earthworms and they graze it heavily, which could dramatically impact their abundance in the soil. By grazing fungi on or near plant roots, the earthworms not only can damage the roots, but they prevent the plant and fungi from forming the symbiotic relationship where mycorrhizal fungi exchange nutrients and water for carbohydrates with green plants. If the fungi can't get enough food, they will die back even further. For some of the native plants that need mycorrhizal fungi, especially when the plant is young and small, survival will be difficult if earthworms prevent this relationship from being formed.

Seeds produced by the few surviving plants are no longer protected by the forest floor, allowing animals, including worms, to find and eat the seeds. If the seeds survive to germinate, they are no longer protected from temperature extremes. The seeds will be more vulnerable to death, especially native herbaceous plants that germinate very slowly, taking two or more years to grow into a small plant. Seed that one buys in a garden store will germinate upon putting them in the soil. In a hardwood forest ecosystem, most native plant seeds don't germinate fully in one season. Most need to go through a freezing and warming cycle (winter and summer) at least once and sometimes twice before growing into a small plant. The forest floor protects these seeds and tiny plants from predators and extremes in temperature and moisture, making the loss of the forest floor devastating to native plant production.

Leaves and twigs continue to fall to the forest floor each year but are rapidly eaten by the earthworms. Thus the forest floor never redevelops.

Earthworms change the *nutrient cycling* in the forest by increasing the rate at which litter disappears. They do this in two ways. First, earthworms break up the litter into tiny pieces and second, those tiny pieces get broken down by bacteria. Imagine eating a tootsie roll sucker and your favorite part is the chewy center. To get to the chewy center, you would have to eat through the hard candy exterior. Bacteria prefer the "center" of the litter (sugars and carbohydrates) found on the forest floor but first have to get through the fiber (lignin and cellulose). If you imagine licking your way to the center of a tootsie roll, it would be much slower than biting through the hard candy exterior. Likewise, it takes time for bacteria to get through the fiber except when earthworms are part of the system. Earthworms act as the teeth and expose the sugars and carbohydrates to the bacteria, allowing for a relatively quick breakdown of the litter compared to bacteria consuming the litter alone. The end result is leaf litter will be consumed at a rate faster than it is produced.

Nutrients needed for plants to grow are now found in the new A horizon composed of earthworm casts. Earthworm casts don't have more nutrients than the forest floor (since it came from the forest floor, it couldn't) but as the litter passes through the earthworm gut,

a lot of it is converted to forms of nutrients that plants can easily absorb. Although the total amount of nutrients does not exceed the forest floor, earthworms cause more nutrients to be available to plants at any given time. However, if the plants don't absorb these nutrients quickly, they can be washed away or leached when it rains. Two things increase the likelihood of nutrient leaching. First, there are not enough plants or root systems to absorb the amount of nutrients available. Second, with all of those earthworm burrows, water can wash the dissolved nutrients down through the soil, below the plant roots or out into rivers and streams. Nutrients that would have been cycled within the hardwood forest ecosystem can either be lost underground or transferred out to another system.

Bacteria still primarily breakdown the litter into nutrients that plants can use. However, now most of that activity takes place in the earthworm gut and not in the litter, much like a compost pile.

Most *insects* living in hardwood forests rely upon the forest floor for food and protection. Since earthworms eliminate the forest floor, we would expect that not only the numbers of insects will decrease, but the variety of insects will also decrease.

Amphibians and Reptiles that live in hardwood forests are especially adapted to the moist and cool conditions of the forest floor. When the forest floor is removed, they no longer have this protection from predators and from drying out. With a decrease in the number of insects, a critical food source is diminished which can lead to additional stresses. Taking into account the loss of the forest floor, which impacts plant reproduction, insect, amphibian, and reptile habitat, we can conclude the addition of earthworms into a hardwood forest ecosystem severely impacts the diversity of that system.

Different *mammals* will each be affected according to their dependence on various aspects of the ecosystem. Like amphibians and reptiles, small mammals like voles rely on the forest floor for protection and food. With fewer insects and fungi, their preferred foods, these small mammals will probably die back after earthworms remove the forest floor. If the voles disappear, weasels are threatened because voles are a food source for them. Medium sized mammals like raccoons, hare, and porcupine will also find less food after earthworms invade and will probably look for other habitats to supply their needs. Large mammals like white-tailed deer and bear are only occasional visitors to hardwood forests. If they find their food source has disappeared, they will simply move on and look elsewhere and spend less time in the forest.

Birds that breed and nest in the hardwood forests rely on the forest floor for both food (mostly insects and seeds) and nest sites. With fewer plants, there will be fewer seeds. Other birds use the layers of vegetation that seedling, sapling and shrubs provide for nesting and food. When these layers die back, important nest sites disappear along with important sources of berries and seeds for food. As a result, we would expect to see and hear fewer birds in our forests after earthworms invade.

Herbivory in a hardwood forest ecosystem occurs all the time but doesn't control what plants are or are not present. However, herbivory can have a severe impact when earthworms reduce the diversity and quantity of the remaining plant species. White-tailed deer are an important mammalian herbivore in hardwood forests in Minnesota. They particularly like to eat many of the native herbaceous plants that grow in these forests because they do not have bitter or toxic substances in them. When plants are numerous, deer can eat many of them and still not negatively impact the population over time. We can illustrate how this might happen in this hypothetical scenario:

- Let us suppose there are 10,000 plants in a given area and those 10,000 produce 1,000 new plants every year. If the deer in the area eat 1000 plants in a year, the plant population remains constant.
- When earthworms are added to the equation, their impact on plants is significant because they decrease the number of mature plants and the number of new plants produced each year by eating the seeds and damaging the bulbs, rhizomes and corms.
- After earthworms invade our hypothetical ecosystem, 1000 plants remain with those 1000 plants only producing 50 new plants a year. The deer population stays the same and thus will continue to consume 1000 plants a year as long as they are available. In two years, these plants would be eliminated from this ecosystem.

Jack-in-the-pulpit is one native plant species that has bitter tasting or toxic substances in their leaves, as do many exotic plant species. Deer and other herbivores will avoid eating these plants. After some time, they may take over the forest understory where earthworms are present

Sampling methods for earthworms

If site conditions suggest a worm infestation, the following methods can be used to estimate densities and identify the species of earthworms present.

To collect worm specimens for identification and sample the worm densities, mix 1/3 C. dry yellow mustard with 1 gal water. Using a 1 – 2 ft² or 1/3 m² frame with 2inch high sides pressed into the ground a ways, pour the solution slowly and evenly over the area within the frame. Collect the worms as they “fly” out and identify.

Lumbricus terrestris (the big earthworms) are the only burrowing worms that leave castings. Counting the castings/m² can give an estimate of worm densities.

Planting well-rooted shrubs and forbs into a woodland or forest damaged by worms or after buckthorn removal is a potential way to restore understory diversity and cover. Seed germination is difficult in worm infested areas because they can knock them over and also may consume the young seedlings.

If planting seedlings, place an acidic mulch such as wood chips around the base of each plant. This may minimize worm activity at least for the first growing season.

Appendix E: Resources

Contacts:

National Park Service

Mississippi National River and Recreation Area

Susan Overson

111 Kellogg Blvd.

St. Paul, MN 55101-1256

651-290-3030 ext. 225

susan_overson@nps.gov

Minnesota Department of Natural Resources (DNR)

Division of Wildlife:

Brian Lueth, Area Wildlife Manager

5463-C W. Broadway

Forrest Lake, MN 55025

brian.lueth@dnr.state.mn.us

Hannah Dunevitz, Regional Plant Ecologist

1200 Warner Rd.

St. Paul, MN 55106

651-772-7570

hannah.dunevitz@dnr.state.mn.us

Division of Forestry:

Art Widerstrom, Area Forester

(651) 982-9820 X224

art.widerstrom@dnr.state.mn.us

Ramsey County

Ramsey Soil and Water Conservation District

2015 Rice St. Roseville, MN

651-488-1476

West Side Citizens Organization

Bluff Task Force

Equipment:

Tree planting, seeding and fire fighting tools and equipment:

Forestry Suppliers, Inc.

205 West Rankin St.

Jackson, MS 39201

(800) 647-5368

www.forestry-suppliers.com

Princeton, MN 55371
(763) 633-4342
www.prairieresto.com

Minnesota Native Landscapes
14088 Hwy. 95 NE
Foley, MN 56329
(320) 968-4222
www.mnNativeLandscapes.com

Smith Overlook, 2002



Barge Terminal No. 1, 2002




Flint Hills Resources: Before Brush Removal, Summer 2003



Flint Hills Resources: After Brush Removal, Fall 2003



Cherokee Prairie Implementation 2003



Eagle Creek, 2003



South St. Paul Levee 2004





News - August 2002

Greening Helps Shape Regional Vision for Protecting Natural Resources

Great River Greening has identified the most significant natural resources along the Mississippi River corridor from the Ford Dam in Saint Paul to Hastings. Greening's computer-based ecological analysis will inform the Metropolitan Council's work with the communities in the corridor on river-related projects.

Greening found that areas of high ecological significance are concentrated in four general areas: Vermillion River Bottoms, Pine Bend, the Pig's Eye area and lands around the confluence of the Minnesota and Mississippi rivers. These areas still support remnant natural vegetative communities and provide crucial habitat for wildlife; many of the plants and animals found in these areas are rare.

Great River Greening is participating in a similar ecological study that will help target natural areas for protection along the entire Mississippi National River and Recreation Area, 54,000 acres along the river corridor from Anoka County to the Vermillion River Bottoms in Hastings.

Welcome to Great River Greening's new e-mail postcard, which will offer periodic updates on Greening news and events. For more on this story and other Great River Greening news, click on [River Valley Reader](#) for a PDF file of our Fall 2002 newsletter. Or click on the menu above to visit another section of our website.

To remove your name from this e-mail list, please reply to sender with the word "remove" in the subject line.



News - January 2004

GREAT RIVER GREENING—RESTORING THE LAND, RENEWING COMMUNITIES

January 2, 2004

To Partners and Friends of Great River Greening:

Great River Greening is excited to start the New Year thanking our volunteers and partners who make our work possible. At Greening, we are grateful to be able to live out our beliefs and values by focusing on what is important—the restoration of our land and the renewal of our communities.

With 14,000 volunteers and growing stronger, we are convinced that individuals and communities are eager to get involved in well-organized and truly significant on-the-ground environmental work. In the most direct way possible, our efforts:

- contribute to the health and bio-diversity of ecosystems
- beautify our surroundings
- enhance our community's natural heritage and sense of place

This year's line-up of volunteer events provides just that—a variety of opportunities across the metro area to practice land stewardship. On February 21, Greening and its partners will host the first of many events to restore Crosby Regional Park, an often-overlooked backwater sanctuary along the Mississippi River in Saint Paul. On May 1, volunteers will plant a section of River Park on the Mississippi in Brooklyn Park, helping to create 12.5 acres of prairie, oak forest, an interpretive garden, and a stormwater system. Designed by Greening landscape ecologists, River Park is one of the few professionally designed natural parks in the metro area.

The 2004 restoration event schedule is below. The detailed event schedule will be up soon on the Great River Greening, www.greatrivergreening.org. To sign up for an event contact Jane Stubblefield at jstubblefield@greatrivergreening.org.


2004 GREAT RIVER GREENING RESTORATION EVENTS

January 31	Bucks and Buckthorn	Warner Nature Center, St. Croix Greenway
February 21	Crosby Regional Park	Saint Paul
8/6/2004		

March 13	Cherokee Bluffs	Ohio St. bluff, Saint Paul
March 20	Hastings River Flats	Mississippi River bottom, Hastings
April 3	Supervisors Adv. Training	Raptor Center, Saint Paul U of M campus
April 17	Bucks and Buckthorn	St. Croix Watershed Research Station, St. Croix Greenway
May 1	River Park	Brooklyn Park
May 6/7 or Wk of May 10	Heritage Park	West of downtown Minneapolis
May 22	Prospect Crest	Above Wabasha St. Caves, Saint Paul
June 5	Mississippi River Gorge	West River Parkway, Minneapolis
June 12	South St. Paul Leyee	North of 494, South St. Paul
August 21	Bucks and Buckthorn	Site TBD
September 7	Crosby Regional Park or Mounds Park	Site TBD
September 18	MN Valley National Wildlife Refuge	Louisville Swamp Unit, Shakopee
October 16	Rosemount, Inc.	Eden Prairie
October 23	Pine Bend Bluffs	Flint Hills Resources, Rosemount
Date TBD	Arcola Mills Historic and Natural Site	St. Croix River, north of Stillwater

Great River Greening's e-mail postcard offers periodic updates on Greening news and events. For more information about Great River Greening, click on the menu above to visit our website.

To remove your name from this e-mail list, please reply to sender with the word "remove" in the subject line. If you have trouble reading this html document, please notify sender, and you will receive a text-only version.



News - February 2004

February 2, 2004

To Partners and Friends of Great River Greening:

Great River Greening had a very successful 2003 as a direct result of the participation and support of its partners and friends! It is truly amazing to see what individuals can do when they pull together as a community with a common goal.

Greening's list of accomplishments showcases the great interest and commitment of individuals to restore valuable and endangered land within their communities. These accomplishments also provide a springboard for Greening to continue to achieve its long term goals in 2004. Thanks again to our volunteers, partners and supporters!

Accomplishment Highlights - 2003

Volunteer Restoration Events:

- Engaged **1,160 volunteers** in **11 events** for **3,820 hours** of restoration work.
- **Removed 18 acres** of invasive species.
- **Planted a total of 1,332 trees and shrubs** and installed **8,891 prairie plants**.
- Have engaged a running total of 13,000 volunteers in restoration activities since 1995.

Volunteer Training:

- Successfully created a partnership with Saint Paul Parks and Recreation to test a pilot of Greening's Stewardship Program. The pilot used resident stewards to monitor the health of and educate communities about designated Saint Paul natural areas.
- Conducted 812 hours of advanced training for 100 supervisors.

Awards:

- Significant Contribution to Conservation: from the Garden Club of America
- The Marion Thompson Fuller Brown Conservation Award: for an outstanding Conservation Exhibit at the Garden Club of America Zone XI annual meeting.

Ecological/Technical Work:

- Completed 4 major management plans for high quality, natural area remnants, including a native prairie, which contains one of the largest Minnesota populations of the rare *Kitten Tails* plant.
- Completed 8 landscape designs achieving ecological goals such as biodiversity, stormwater infiltration, and sustainability.
- Conducted a vegetation and hydrology analysis of stormwater systems in the Twin Cities areas for the book *Plant for Stormwater Design* by Dan Shaw, Greening Landscape Ecologist and Rusty Schmidt,

8/6/2004

Independent Designer.

Fundraising:

- Exceeded overall fundraising goal by 20%.
- Received several new grants including Greening's first national grant from the Carolyn Foundation.

For a summary of 2003's accomplishments, as well as a history of Greening's accomplishments every year since its inception, visit www.greatrivergreening.org.

Great River Greening's e-mail postcard offers periodic updates on Greening news and events. For more information about Great River Greening, click on the menu above to visit our website.

To remove your name from this e-mail list, please reply to sender with the word "remove" in the subject line. If you have trouble reading this html document, please notify sender, and you will receive a text-only version.



News - June 2004

GREENING'S RESTORATION EXPERTISE AT WORK

June 10, 2004

To Partners and Friends of Great River Greening:

Crosby Regional Park is one of many restoration projects in which Great River Greening is currently involved. In addition to providing design guidance for Crosby Park, Greening will be conducting major habitat restoration work, a natural resources inventory, authoring a restoration management plan and developing a plan for the bluff trails. "This project is an innovative method to address trail issues from both a cultural and ecological perspective," said Greening Landscape Ecologist Dan Shaw. "We're hoping it will be a model for natural area trail planning in the region."

Crosby Park is a sanctuary of wetlands and floodplain forest on the Mississippi River and is Saint Paul's largest nature preserve. It provides valuable habitat to a diverse community of plant and animal life including rare species such as the Prothonotory Warbler and Northern Shrike. At this point in the project Greening has removed a substantial amount of buckthorn through crew work and a volunteer event held this past winter. Greening is well into the trail plan which will be completed by the end of June and the management plan work has been started.

The project is based on a \$50,000 National Park Service (NPS) grant secured by the Mississippi National River and Recreation Area (MNRRA) office of NPS and Great River Greening. Other funding comes from Capitol Region Watershed District, City of Saint Paul Division of Parks and Recreation, and Great River Greening through a Carolyn Foundation grant and through funds from the Environmental and Natural Resources Trust Fund as recommended by the Legislative Commission on Minnesota Resources. Ramsey Soil and Water Conservation District is also a project partner.

The innovative component of the trail study will be its sustainable perspective. It will offer design guidance on materials, trail layout, methods to control erosion, and methods to revegetate portions of the bluff and control the flow of water going over the bluff. The focus is on the bluff and the trail as a force within the park. Greening ecologists are also putting together a management plan for the whole park including a complete plant inventory and management recommendations for ecosystem restoration. As one of the largest floodplain forests and river backwater remnants in the metro area, Crosby's 500 acres represent an ecosystem that is vital to the life of the Mississippi River.

For more information about Greening's ecological work please visit our website: www.greatrivergreening.org

8/6/2004

Remember to reserve your seat for the Great River Greening Gala July 3


You won't want to miss the party of the season during the Grand Excursion. The evening will begin with a dinner cruise, music and silent auction and will conclude with a fireworks display in celebration of the arrival of the Grand Excursion Flotilla in Saint Paul. The Gala will raise money for restoring Twin Cities rivers and watersheds through the work of Greening. Visit the Greening website now to reserve your place.

Volunteers needed for Greening's June 12th South St. Paul Planting Event

Volunteers are still needed to help create a park, designed by Greening, in South St. Paul by planting trees, shrubs and prairie plants. The park will add wildlife habitat, stormwater filtering and erosion control. To register email or call Jane at jstubblefield@greatrivergreening.org or 651-665-9500 ext. 11.

Great River Greening recently installed a new database, which brought together information from several places. Please let us know if you would like to be removed from our E-Postcard list. Just reply to the sender with the word "remove" in the subject line.

Great River Greening's e-mail postcard offers periodic updates on Greening news and events. For more information about Great River Greening, visit our website www.greatrivergreening.org.




The

River Valley Reader

It's habitat forming

Greening ecologists design unique River Park addition

THANKS TO NATIONAL PARK SERVICE (NPS) FUNDING AND GREAT RIVER GREENING ECOLOGISTS, the City of Brooklyn Park will soon be home to one of the few professionally designed natural parks in the metro area. A 13-acre northern addition to River Park, on the banks of the Mississippi, is being restored to create prairie, oak forest, an interpretive garden and a natural stormwater system for runoff from the neighborhood.



"The land was once a farm field and then a grassy front yard," said Brooklyn Park Recreation & Parks Manager Jay Lotthammer. "We took the opportunity to jump to the next level of progressive park development—we're restoring urban land, promoting natural park design and supporting the health of the Mississippi River."

Project partners NPS, Greening and Brooklyn Park had several goals for the addition to River Park. The Greening design creates a passive recreation area next to an active area, inviting children and adults to enjoy and learn about native plant communities along the Mississippi River. The Park recreates the native vegetation corridor along this stretch of river, improving wildlife habitat, reducing erosion and lessening stormwater impact.

A series of rain garden-like areas creates a natural stormwater system receiving runoff from the nearby residential neighborhood, filtering it and releasing the cleaned water into the river. Previously, stormwater traveled the traditional route from the streets, through a pipe, to the river. In addition to improving the quality of

water entering the Mississippi from the neighborhood, the infiltration plants will add color and attract birds, butterflies and other wildlife to the area. Directing

stormwater to parkland for natural treatment is a unique aspect of Greening's River Park design and one of the larger demonstrations of the system.

Great River Greening staff is thrilled to be working in Brooklyn Park for the first time and to be partnering again with NPS. To launch

the creation of River Park, 250 volunteers will help with planting on Saturday, May 1. See page 5.

GRAND EXCURSION 2004™

Great River Greening Gala

GET A FRONT ROW SEAT for the Grand Excursion fireworks finale on the Mississippi in Saint Paul and support Great River Greening. Purchase tickets now for the July 3 *Great River Greening Gala* fundraising event. This evening will be the culmination of years of planning for the arrival of the Grand Excursion Flotilla in Saint Paul.

- Saturday evening, July 3, 2004
- Grand Excursion and Taste of Minnesota fireworks
- Musical entertainment and more

Visit www.greatrivergreening.org for details or call Katherine Nielsen at 651-665-9500, x28.

Message from the Executive Director

"EXOTIC PESTS RUN WILD IN FLORIDA" WAS A HEADLINE RECENTLY IN THE NEW YORK TIMES.

It seems southern Florida is teeming with Burmese pythons, African monitor lizards and South American Monk parakeets—just some of the more than 200 nonnative species at large in Florida because of careless or unknowledgeable breeders, dealers or owners. Monitors eat the eggs of burrowing owls which are an endangered species, pythons displace native snakes and parrots compete with native owls and woodpeckers for nesting sites.

All across the country, every state has problems with exotic animals and plants which affect the ecosystem. Minnesota is no different. Asian carp have been in our headlines recently as they steadily move up the Mississippi and its tributaries, advancing 35 to 50 miles a year and wreaking havoc on the aquatic ecosystems they colonize.

Minnesota's nonnative plants such as Eurasian buckthorn, tartarian honeysuckle and garlic mustard are no less destructive. Of the more than 2000 species in Minnesota, about 20 percent are nonnative. Jay Rendall, Minnesota Department of Natural Resources (DNR)



Greening's Executive Director, Deborah Karasov, leads the battle against buckthorn.

exotic species coordinator, says that nonnative invasive species are one, if not the top, threat to native habitats in the state. One reason is that invasive species harm aquatic habitats and protected habitats in parks and preserves that are not subject to destruction.

Greening fights invasive species

Great River Greening volunteers and staff have been fighting

invasive species and promoting native species since our beginning. We have six buckthorn removal events on our 2004 winter/spring calendar. For two years, we have been one of a select group nationally to receive a grant from the National Fish and Wildlife Foundation to manage invasive plants through a coordinated program of public/private partnership. The West Side Bluffs in Saint Paul will benefit.

Greening helps people experience the huge scope of the problem through invasive species removal events. Once volunteers spend hours hacking away at thickets of buckthorn, we believe they will be ever more sensitive as voting citizens to policy, budget and resource questions related to invasives, including the need for ongoing monitoring and management.

—Deborah Karasov

Big Rivers Partnership meets goals

AS JUNE 2004 APPROACHES, GREAT RIVER GREENING IS WRAPPING UP WORK ON THE BIG RIVERS PARTNERSHIP (BRP), one of the first important restoration collaborative organizations in the state. It was launched in 1999 and continued in 2002 with funds from the Minnesota Environmental Trust Fund as recommended by the Legislative Commission on Minnesota Resources (LCMR). The Big Rivers Partnership brought together nonprofit organizations, government agencies and private landowners to restore critical river valley habitat as well as to engage community volunteers in the metro

area's most important natural resource feature—its three major rivers, the Mississippi, Minnesota and St. Croix.

What has BRP accomplished?

Great River Greening has coordinated all of the BRP initiatives and is proud to announce that all of the goals set by the LCMR have been met or exceeded. For 2002-2004, the LCMR appropriated \$910,000 to BRP. A previous \$800,000 grant has already been completed for 1999-2001.

continued on next page

As of December 2003, BRP implemented projects on 1,488 acres, triple the goal. More than 3000 volunteers participated in BRP habitat restoration projects and more than \$1.3 million in cost-share funds has been raised.


Partnership members, including the Minnesota Department of Natural Resources (DNR), National Park Service and Great River Greening have been completing ecological rankings of the metropolitan stretches of the Mississippi and Minnesota river valleys to complement the regionally significant areas identified by the Minnesota County Biological Survey (MCBS). The Big Rivers Partnership has focused on areas with potential for restoration, not just the highest quality natural areas identified by MCBS.

How is the BRP model important?

The Big Rivers Partnership is significant because it provides a model for organizational collaboration on

restoration implementation, site prioritization and leveraging of funds. The success of BRP proves that multiple organizations, both private and governmental, can work together to achieve conservation goals.

Big Rivers Partnership is already a model for another conservation and restoration collaborative, Metro Wildlife Corridors (MWC). The goal of MWC is to bring together nonprofit organizations and government agencies to protect and restore natural areas, with priority for regionally important wildlife habitat. Metro Wildlife Corridors also encompasses land acquisition which was not a part of BRP's mission. Greening is an active participant in MWC.



Greening to restore Crosby Park

ON FEBRUARY 21, 50 LOCAL RESIDENTS ENTHUSIASTICALLY KICKED OFF A MAJOR habitat restoration project at Crosby Regional Park with a buckthorn removal event. This regionally significant sanctuary of wetlands and floodplain forest along the Mississippi River is part of the City of Saint Paul Division of Parks and Recreation.

Great River Greening will conduct major habitat restoration work and a natural resources inventory, and write a restoration management plan for Crosby Park in 2004. The project is based on a \$50,000 National Park Service (NPS) grant secured by Saint Paul Division of Parks and Recreation, and Great River Greening. Other partners in the project are Capitol Region Watershed District; Mississippi National River and Recreation Area office of NPS; and Ramsey Soil and Water Conservation District.

Located near the confluence of the Mississippi and Minnesota Rivers, Crosby Park is Saint Paul's largest



nature preserve. It provides valuable habitat to a diverse community of plant and animal life including rare species such as the Prothonotary Warbler and Northern Shrike. As one of the largest floodplain forests and river backwater remnants in the metro area, Crosby's 500 acres represent an ecosystem that is vital to the life of the Mississippi River.

continued on next page

continued from previous page

As is the case with most urban natural areas, portions of Crosby Park are degraded by recreational use, erosion and the invasion of exotic species. It features extensive hiking and biking trails, marsh ponds, wildflowers and outstanding bird watching. Past farming activities also contributed to degradation. Crosby Park was originally staked out as farmland in 1858 by Thomas Crosby and was farmed by a succession of families until the 1960s when it was purchased for the Saint Paul Division of Parks and Recreation.

Funding for the project comes from the Capitol Region Watershed District; City of Saint Paul Division of Parks and Recreation; Great River Greening, through a Carolyn Foundation grant and through funds from the Environmental and Natural Resources Trust Fund as recommended by the Legislative Commission on Minnesota Resources; as well as the NPS grant.

Project funding also provides for Great River Greening to conduct a focused analysis of bluff erosion and nature trail deterioration.



Greening announces four new board members

FOUR NEW BOARD MEMBERS ADD GREATLY TO GREENING'S reputation, and business and technical expertise. We are pleased to welcome Susan Galatowitsch, Ann McMillan, Kristine Sundberg and Alvin-o Williams. Each brings a unique set of skills and background, and all are incredible assets to the organization.

Susan Galatowitsch is an Associate Professor of Horticultural Science and holds a joint appointment in the Department of Landscape Architecture at the University of Minnesota. She has a Ph.D. in Ecology and Evolutionary Biology from Iowa State University as well as an M.S. in Botany. Sue brings to Great River Greening a nationally-recognized expertise in restoration and ecology with a particular emphasis in wetland restoration.

One of the many assets Ann McMillan brings to Greening is her Landscape Architecture background. Ann has an M.S. in Landscape Architecture from the University of Minnesota. She has served on the boards of the Minnesota Landscape Arboretum and the Saint Paul Garden Club, and was an early supporter of Greening. She currently serves on the University of Minnesota

College of Architecture and Landscape Architecture national board.

Kristine Sundberg joins Greening with more than 20 years of communications, investor relations and public affairs achievement. Kristine has an M.S. in Environmental Studies from the University of Illinois. She is currently Executive Counsel for the Sage Group, a full-service public relations, marketing communications, investor relations and communications firm in Minneapolis. Among her clients are AT&T; Global Bridge, Inc.; and Lee Financial.

In Alvin-o Williams, Great River Greening gains considerable experience in both law and business management expertise. He has a B.S. from the University of Minnesota and a J.D. from Hamline University. Alvin-o is co-owner of Castaneda Williams, Ltd., a successful Edina advertising agency that provides total business solutions to clients. Among his clients are Boston Scientific, Target Corporation and Minnesota Minority Supplier Development Council.



Save

Volunteers Field Guide

Opportunities to help green the Twin Cities

Spring Restoration Events

Tools, equipment and refreshments provided. Pre-registration required. Contact Jane at 651-665-9500 x11 or register online at www.greatrivergreening.org under "Volunteers."

Bucks and Buckthorn, St. Croix Watershed Research Station, Marine on St. Croix

SATURDAY, APRIL 17 • 8:30 A.M. – NOON

Volunteers needed: 150 Supervisors needed: 15

SPEND AN INVIGORATING SPRING MORNING in the St. Croix Valley for the largest Bucks and Buckthorn restoration event to date and the first public event of the program. Volunteers will remove Buckthorn from land at the St. Croix Watershed Research Station (SCWRS) to help restore the "St. Croix Greenway," an important continuous corridor of 2,300 acres of undeveloped land extending from the St. Croix River toward Big Marine Lake. This marks Great River Greening's first large event on the St. Croix River so we're asking Greening veterans to come out and help us introduce Greening to Valley volunteers. This is a great family event and a wonderful opportunity to discover the SCWRS, a field research station of the Science Museum of Minnesota.

River Park, 83rd Ave. & West River Road, Brooklyn Park

SATURDAY, MAY 1 • 8:30 A.M. – NOON

Supervisors needed: 22

THIS PLANTING EVENT HAS FILLED WITH 250 ENTHUSIASTIC VOLUNTEERS. WE STILL NEED SUPERVISORS.

Prospect Crest, Above Wabasha Caves off Prospect Blvd. in Saint Paul

SATURDAY, MAY 22 • 8:30 A.M. – NOON

Volunteers needed: 60 Supervisors needed: 6

PART OF THE GRAND EXCURSION CELEBRATION in Saint Paul, volunteers will plant prairie grasses on the top of the bluff to help reduce erosion and restore a viewing area historically important to the neighborhood. The steep bluff face will be planted by members of the Minnesota Climbers Association. This event is not suitable for children under 14.



Mississippi River Gorge, 36th Street and West River Parkway, Minneapolis

SATURDAY, JUNE 5 • 8:30 A.M. – NOON

Volunteers needed: 80 Supervisors needed: 8

RESTORATION WORK CONTINUES ON NATURAL PARKLAND in the Longfellow community along the only true gorge on the Mississippi River. This project is a partnership with the Minneapolis Park Board, Friends of the Mississippi River, the Longfellow Community Council and Greening. Volunteers will plant 4000 prairie plants and may help remove some small brush. Suitable for families with school-age children and groups.

South St. Paul Levee, Just north of Highway 494 in South St. Paul 12

SATURDAY, JUNE 7 • 8:30 A.M. – NOON

Volunteers needed: 225 Supervisors needed: 22

VOLUNTEERS WILL PLANT TREES AND PRAIRIE PLANTS in a beautiful park along the Mississippi River and along a major bicycle path that follows the levee. The park, designed by Greening, will add wildlife habitat, storm water filtering and erosion control along the river. Suitable for families with school-age children and groups.



Supervisors Field Guide

Opportunities for Greening's volunteer supervisors

2004 Supervisor Recruitment

Are you ready to "branch out?"

EVER WONDERED HOW YOU CAN BECOME more involved in the work of Great River Greening? Are you looking for a place to share your knowledge of the environment with both adults and children? Are you eager to meet new people in communities across the Twin Cities who share your interests and values? If your answer is "yes," we invite you to become one of our Great River Greening Supervisors.

Greening Supervisors receive three hours of Basic Training from Greening ecologists on species identification, restoration techniques, management of volunteers and the Great River Greening organization. To help new Supervisors become comfortable with their role, they are paired with an experienced Supervisor for their first event.

We ask all Supervisors to commit to working at least two restoration events per year. No previous experience or environmental knowledge is required. We just ask for a genuine interest in restoring the environment and a willingness to work with diverse volunteers of all ages. Many of our Supervisors are also Master Gardeners or Tree Care Advisors and are willing to share their expertise – a great benefit for those with less experience.

Supervisor Role

Supervisors are trained to lead a group of 10 to 15 volunteers at our Saturday morning planting and removal events.

They are responsible for:

- Introducing volunteers to the work of Greening
- Sharing information about the site and the significance of the work
- Providing instructions for proper planting and removal
- Instructing volunteers in the safe and proper use of tools
- Motivating volunteers to complete the assigned activities
- Sharing their own knowledge about the environment
- Ensuring that everyone has a good time!

An updated Basic Training course will be held on Saturday, July 24 for new Greening Supervisors and for those who may not have supervised for a few seasons and would like a "refresher."

Additional Advanced Training sessions are offered several times a year. Recent courses include Native Landscape Design for Homeowners, Shrub Identification and Prairie Restoration Beginning with the July training, there will be a small fee for each course: \$15 for Basic Training and \$10 for all Advanced Training.

New for 2004!

In response to feedback from our Supervisor Survey last November, we have some new activities planned this year. (Thanks to all of you who sent in your survey!) There will be opportunities to help educate youth and help promote Greening at community events. We're planning

an informal e-newsletter to communicate information, tips and news. Advanced training courses will cover more in-depth plant identification and habitat restoration, and at year's end we will celebrate the accomplishments of all Supervisors with an appreciation event.

Thank you for your interest in Greening. We look forward to adding you to our valued corps of Supervisors!

Supervisor Training

Saturday, July 24

Updated Basic Training for Supervisors
Time and location TBD

Fee: \$15

For more information on becoming a Supervisor or to register for the training course, contact Jane at 651-665-9500 x11 or jstubblefield@greatrivergreening.org

A special thank you to our 2003 event Supervisors:

Dianne Ballentine	Eli Lewis
Julia Bohnen	Cindy Matiski
Gina Bonsignore	Lisa McDonald
Darlene Charboneau	Doug Mensing
Nancy Hansen	Lynn Meyer
Gordon Hanson	Bev Roberts
Mimi Hottinger	Nell Ruemmele
Bret Hubert	Karen Schik
Patsy Huberty	Mary Stade
Linda Hustad	Lorrie Stromme
Maxine Hughes	Fred Tio
Gloria Jackson	Roger Tix
David Kelley	John Vickery
Bark Kirkpatrick	Sarah White
Ray Kivlahan	June Young
Jane Klein	

Thank you to Greening's 2003 Corporate Sponsors!

Great River Greening continues to build strong partnerships with area businesses, just as we did with our very first project. Together we are—restoring the land, renewing communities—through our mutual goal of maintaining a healthy and vibrant river valley and watershed.

In 2003, the generous support of businesses like Bailey Nurseries, Ford Motor Company, Wells Fargo Bank and 3M Foundation, helped Greening raise nearly \$50,000 in corporate contributions. Great River Greening's first fundraising event, the Great Rivers Marathon, was also a big success and attracted additional corporate sponsors, including Wellington Management and Superior Minerals Company.

Thank you to our generous 2003 Corporate Sponsors and Patrons who help make our work possible.

\$5,000 or more

3M Foundation, Inc.
Bailey Nurseries
Ford Motor Company
Wells Fargo Bank Minnesota

\$1,000 – 4,999

Alter River Terminal*
Briggs & Morgan, P.A.*
Cemstone Products Company*
CenterPoint Energy Minnegasco*
Custom Drywall, Inc.
Hawkins, Inc.
Leonard, Street and Deinard
Lowry Redevelopment Partners, LP
Loucks Associates*
Saint Paul Port Authority*
Saint Paul Riverfront Corporation*
Superior Minerals Company*
Wellington Management*

\$500-999

Aggregate Industries*
Asset Recovery Corporation*
Bidwell Maintenance, Inc.*
CHS Cooperatives*
Ceridian Corporation*
Land O'Lakes, Inc.
Lindquist & Vennum P.L.L.P.*
LHB Engineers & Architects*
Rexam Beverage Can Company*
SKB Environmental*
Wabasha Business Center*

\$100-499

Barton Sand & Gravel
Bjorklund Trucking
Capitol City Beverage Sales

Cermak Rhoades Architects
Close Landscape Architecture
District Energy St. Paul, Inc.
Michels Corporation
Natural Resources Restoration
The Awes Agency, Inc.
The St. Paul Companies, Inc.
Thor Construction
U.S. Bancorp Foundation

\$1-99

American Express Foundation
Cub Foods
Stefan Helgeson Associates, Inc.
Nalco Printing Company
Sauer-Danfoss Company
UPS Foundation
Viking Office Supplies
*Great Rivers Marathon Sponsor

In-Kind Contributors

All-Wood Products
Another Sun Nursery
Bahn's Landscape & Design
Bailey Nurseries
Beisswenger's How-to-Hardware
Divine Swine
Houston's Neighborhood Supermarket
Ideal Printers, Inc.
Minnesota Nursery & Landscape Association
Minnesota Timberwolves/Lynx
Outback Nursery
REI
Unique Balloons and Catering
Valley Creek Express Blower Service
Wabasha Deli & Cafe
Youth Frontiers

2004 Corporate Sponsorship program supports restoration events season

IN THE THIRD YEAR OF OUR SUCCESSFUL Corporate Sponsorship program, we are focusing directly on the heart of Great River Greening's work—our volunteer events. This program provides an opportunity for businesses to contribute funds to directly support Greening's 2004 restoration event season. The dollars go directly to the actual work of planting and restoring a healthy river corridor and watershed. By sponsoring Greening's restoration events, companies are helping to create a more attractive and inviting urban environment.

Thank you to our 2004 Sponsors who have already signed on: Ceridian Corporation, CHS Cooperatives, Custom Drywall, Inc., Faegre & Benson, Saint Paul Port Authority, The Toro Foundation, Upper River Service and the Wabasha Business Center.

To learn more about Corporate Sponsorship, contact Pam McLellan, at 651-665-9500, x18.

Thank you to 100 new donors!

Great River Greening thanks 100 new donors and funders in 2003; and a heartfelt thanks to those who continue to support Greening every year. Your generosity is helping to restore the prairies, savannas and forests that once decorated the blufflands and natural areas of the Twin Cities. For this and all you do to preserve Minnesota's natural heritage, we are grateful.

Bluffland Benefactors

\$1,000 or more

Elmer L. & Eleanor J. Andersen Foundation
Katherine B. Andersen Fund of The Saint Paul Foundation
Gordon and Margaret Bailey Foundation
F.R. Bigelow Foundation
Bush Foundation
Carolyn Foundation
David D. & Vanessa D. Dayton
Julia W. Dayton
Steven J. Holmstoen
Marbrook Foundation
The McKnight Foundation
The Nash Foundation
Northern Environmental Support Trust
Ford J. and Catherine T. Nicholson
Margaret Ordway
Irwin Andrew Porter Foundation
Terhuly Foundation
The Saint Paul Foundation
Trust of Frederick T. Weyerhaeuser
Jonathan and Sarah Wilmschurst
Unity Avenue Foundation

Savanna Sponsors

\$500-999

Art & Martha Kaemmer
Joseph & Kathy Kingman
Mielke Family Foundation, Inc.
James and Susan Vest

Prairie Patrons

\$100-499

Mary Anderson
Monica Angle & Sam Magavern
Mark Bailey
Gina Bonsignore & Richard Garber
David Boyce
Susan Brewster & Edwin McCarthy
Ellen & Peter Brown
Bob & Gerry Bullard
Kathleen Cleary
Jay & Page Cowles
Lydia Crawford & Phil Davies
Tom & Patty Dear
Olivia Dodge
Beth and Kevin Dooley

Ruth Erkkila
Dian Eversole
Gary Gardner
Jim and Joan Gardner
Cathy Geist
James and Teddy Gesell
Terry and David Gilbertstadt
Kevin and Julie Hartwell
Greg and Julie Honsa
Libby Johnston
Deborah Karasov & David Olson
Phyllis Karasov and Alan Olstein
Larry and Peggy King
Theodora Lang
Margy & John Ligon
Deborah Loon Osgood
Rolland & Ella Loon
Robert & Helen Mairs
Bonnie Marron
Charlene McEvoy and Doug Olson
Ann & Doug McMillan
Mary McMillan
Jenny Mockenhaupt
Brad Moore
Eric and Linda Morrison
William and Allison Moyers
Betty Myers
Constance Otis
James Pagliarini & Elizabeth Raymond
David Pasiuk
Gary Petersen
Mary & Richard Stanley
Hollis & Karl Stauber
Edward & Virginia Stringer
Mike Teirney & Christine Ummoefer
Betty & Frank Tiffany
Greg & Erin Anderson Wenz

Forest Friends

\$1-99

Kathi Anderson
Renee Andres
Julie Andrus
David Bachman
Paul Bard
Vernon Berglund
Lori Biederman & Dave Peterson
Charles Billington

Ann Bitter & Lyn Parker
Jeff & Emily Blodgett
Glen & Margaret Booth
Bruce Broquist
Kirk & Ione Brown
Mary Brown
Rob Buffer & Sally Caudill
Cliff Carey
Carolyn Carr & Jonathan Sellman
Vera & Earl Christy
Stewart Corn
Ronald & Bobbe Daggett
Mollie Dean
Dorothy DeJong
Barbara & Robert De La Vega
Joan DeMarce
Kathy & Steve Dougherty
Laurie Drolson
Susan Dulek
Lilli Ann Eginton
Jennifer Engstrom

Special thanks to 2003 Partners

We also thank our partners who contributed significant resources to specific 2003 Greening projects, making more land restoration possible.

Bucks and Buckthorn Partnership
City of Minneapolis
City of Saint Paul
Environment and Natural Resources Trust
Fund as recommended by the Legislative
Commission on Minnesota Resources
Flint Hills Resources
Ford Motor Company
Friends of the Mississippi River
National Fish and Wildlife Foundation
Metro Wildlife Corridor Partnership
Minnesota Department of Natural Resources
Saint Paul Port Authority

John Errigo
 Kathy Farnell
 Paul Fate
 Charles & Mary Field
 Brian & Lisa Finander
 Dawn Freeman
 Friends of the Minnesota Valley
 Debra Gagner
 Paul Glander
 Norm Good
 Teresa Grant
 Audrey Green
 Rita Gundacker
 Kevin & Jane Gutknecht
 Larry Hampel & Mary Kopet
 Gregory & Kathryn Hart
 Jean Hart
 Betsy Hearn & Drew Danielson
 Greg Heberlein
 Harold Hebl
 Nicole Henry & Erik Rogers
 Patsy Huberty
 Maxine Hughes
 Ruby Hunt
 Kjeld & Karen Husebye
 Carl & Crystal Ireland
 Dana Jensen
 Tom & Colette Kelly
 Sarah Kirkwood
 Allan Klein & Harriet Lansing
 Jeff & Diane Kooistra
 Mike & Beth Kovacs
 Jim & Mary Kubiak
 Craig Larson & Beverly Gerdes
 Barbara & John Ledo
 Annette & Michael LeDuc
 Gayle Lens
 Philip & Amanda Little
 Paul & Cindy Lorah
 Lorrie Louder
 Patrick Loyas
 Maureen Lundgren
 Peggy Lynch
 Teresa Lynch
 Kenneth & Linda Malz
 Donald & Abby Marier
 Nancy Martin
 Lisa McDonald
 Malcolm & Wendy McLean
 Pamela McLellan
 Meredith McNab
 Laurie McRostie
 Edie Meissner
 John Mercer
 Peter Meyer
 Harold & Luann Muller

Marjorie Neihart
 Gayle Ober
 Dolores O'Brien
 Linda Odegård & Harlan Cavert
 Jesse Okie
 Mort Olson
 Kristen Olsrud
 Jeanne O'Neill Sprafka
 Marla Ordway
 Aurea Osgood
 Dick Osgood
 Diane Pearson
 David & Gillian Pogany
 Richard Power
 Susan Pukall
 Jeff Ramsey
 Lynn & Jim Ree
 Walter & Jodell Rockenstein
 Nell Ruemmele
 J. Anders & Ewa Rydaker
 Elizabeth Savage
 Charles Schultz
 Julian & Barbara Sellers
 Jeanne & Gordon Shepard
 Mary & Roger Shepard
 Linda Skallman
 Lisa Smith
 Jackie Sticha
 Jim Storm
 Jim Strupp
 Judy Stubbs
 Barbara Stuhler
 Bob Teetshorn & Paige Winebarger
 Jon & Lea Theobald
 Gil Thoele
 Lucy Thompson
 Tom Trow
 Jerry Voight
 Doris Wagar
 Donald Weden
 Buzz Wilson
 Mary Wilson
 Jim & Kathy Wolf
 Stan Zobel

In Memory

Ronald D. Clark
 Herbert Gerdes
 Frances Smith Williams

In-Kind Contributors

Laura Bates
 Rob Buffler & Sally Caudill
 David Cathcart
 Ellen Fuge
 Debra Gagner

Fred Harris
 Deborah Karasov & David Olson
 Pamela McLellan
 Meredith McNab
 Alicia Murphy
 Katherine Nielsen
 Veronika Phillips
 Dan Shaw & Mary Sobota
 Greg & Erin Anderson Wenz

Donor Profile: David Dayton

"I am absolutely optimistic about the future of open space and natural areas in the Twin Cities," says Greening donor David Dayton, "because of the greater public awareness and the funding partnerships that have been developed through the years." As an organization dedicated to the restoration of valuable and endangered natural areas and open spaces, we couldn't agree more.

As a committed Great River Greening donor and environmentalist, David explains why he chooses to support Great River Greening. He was first introduced to Greening by Board Chair Jonathan Wilmschurst. He then attended a presentation and was impressed. He has since become a valued Greening supporter.

David, an engineer in the metal finishing industry, came upon his environmental interest naturally. He spent his youth on a hobby farm in the western suburbs of Minneapolis, which allowed him to spend many hours out of doors. Today, he continues to cultivate his relationship with the environment. He supports several local and national organizations financially, as well as through volunteering.

As a husband and father of two teenage boys, David says he also pays attention to his own environmental impact and encourages others to do the same. "As an environmentalist, I tell people to think about what they're doing," he says. David believes that everyone can contribute to protecting and restoring the environment in many ways. We at Greening are certainly grateful for his optimism and contributions to support our environmental work.

The River Steward Review

Tips for managing your investment in native plantings

Restoration really pays off

GREAT RIVER GREENING HAS RECENTLY COMPILED information that reveals the cost effectiveness of native—sometimes referred to as sustainable—landscaping. We created a list of landscaping scenarios including traditional irrigated turf and native alternatives, and then calculated their respective installation and maintenance costs over five- and ten-year periods.

The results summarized in the chart were dramatic. Over a five-year period, a prairie costs 78% less than sodded irrigated turf. Over a five-year period, a native woodland costs 25% less than sod.

Regardless of cost, we know that ecological restoration of land is a good idea. Restoring native plant communities in the urban river watersheds improves water quality, wildlife habitat and ecological balance. It also improves air quality through the reduction of the need for mowing.

As the summary table shows, it literally pays off to landscape using sustainable design rather than turf grass with irrigation. Sodded irrigated turf costs six times more to install than a prairie. Maintaining turf is three-and-a-half times more expensive than maintaining a prairie. Woodland costs almost seven times less than sodded

COST ESTIMATE SUMMARY				
Landscaping Scenario	Installation per Year	Maintenance	Total Project Cost over 5 yrs.	Total Project Cost over 10 yrs.
Seeded Turf	\$7,250	\$5,380	\$34,150	\$61,050
Sodded Turf	\$22,050	\$5,380	\$48,950	\$75,850
Seeded Prairie	\$3,600	\$1,530	\$11,250	\$18,900
Seeded Prairie w/ Plugs	\$10,800	\$1,530	\$18,450	\$26,100
Woodland: Trees & Shrubs	\$33,450	\$800	\$37,450	\$41,450

Notes:

1. All estimates are per acre.
 2. Turf costs include installation and maintenance of an irrigation system.
- Details available. Contact Dan Shaw at 651-665-9500, x12.

turf to maintain, but one-and-a-half times more than sodded turf to install. Payoff is long-term for a woodland planting.

In addition to the cost and environmental benefits of restoration, working with Great River Greening has its own benefits. When you use Greening, the fees you pay go back into the community through our efforts organizing volunteer events to improve urban natural areas and open space. By involving volunteers, Greening is building a community of land stewards for the future. So working with Greening on native landscaping is good for the environment, good for the community and good for the bottom line.

Add native plants to container gardens, too!

THE HEART OF THE CITY CAN BE A GREAT LOCATION TO conduct business, but even in the summer it can be a pretty dull, drab place for employees and customers, not to mention the birds! Since there's generally limited green space available, many business owners add plants in containers to brighten up an entrance or outdoor patio. However, seeing the same geraniums and petunias on balconies and in front of every business on the block, gets dull, too.

How can you infuse some interest and creativity into window boxes and terra cotta planters at your business? Native plants can be incorporated into any landscape design, including container gardens, with several added benefits. Native plants require less care than most non-native plants and they are the natural food source for the butterflies and birds along our local rivers. Many native plants will thrive in containers.


continued on next page

SCIENTIFIC NAME**COMMON NAME****HEIGHT**



Aster spp.	Aster	24" (varies by spp.)
Fragaria virginiana	Wild Strawberry	6"
Geum triflorum	Prairie Smoke	6"
Liatris aspera	Rough Blazingstar	18"
Lupinus perennis	Wild Lupine	18 – 24"
Monarda fistulosa	Wild Bergamot	36"
Parthenocissus spp.	Virginia Creeper	Vine
Penstemon gracilis	Slender Beard-Tongue	18"
Polygonatum racemosa	False Solomon's Seal	18"
Rudbeckia hirta	Black-Eyed Susan	12 – 18"
Schizachyrium scoparium	Little Bluestem	18"
Solidago speciosa	Showy Goldenrod	12 – 18"
Sporobolus heterolepis	Prairie Dropseed	12 – 18"
Tradescantia spp.	Spiderwort	12 – 18"
Verbena stricta	Hoary Vervain	12 – 18"
Viola palmata var. pedatifida	Prairie Violet	6"

continued from previous page

Mixing the species in each container will also add visual interest to your plantings. Depending on container size, select and plant three to five species of differing heights but with similar water and light requirements. Include a small sign to identify each plant by its scientific and common names. To further educate your patrons and employees, consider placing a sign that describes the benefits of native plants in the midst of your new garden.



Prairie smoke



These River Steward Tips are made possible through the generosity of the Saint Paul Foundation, Katherine B. Andersen Foundation, F.R. Bigelow Foundation and the City of Saint Paul.



Support Greening!

Your tax-deductible gift will help Great River Greening restore valuable and endangered natural areas along the Mississippi, Minnesota and St. Croix river valleys and their watersheds. Donate online at www.greatrivergreening.org or send this form and check payable to Great River Greening, 35 West Water Street, Saint Paul, Minnesota 55107

Enclosed is my tax-deductible gift of: ☐ \$25 ☐ \$50 ☐ \$100 ☐ \$150 \$_____

Name _____

Organization _____

Phone (h) _____ Phone (w) _____

Address _____

City _____ State _____ Zip _____

E-mail address _____

☐ Please do not include my name in the annual list of donors.

THANK YOU FOR YOUR SUPPORT!

3/04

Greening Wish List

Doing some spring cleaning? If you run across one of these items, donate it to Great River Greening and benefit from a tax deduction for charitable giving! If you can't donate, but have information on where we could obtain a significant discount, please let us know, too. Contact Deb Gagner at 651-665-9500 x10.

- Computers: Pentium III or newer processor, 4GB or greater hard drive
- Monitors: 17" or larger, in good condition
- Color laser printer: Good quality, reliable
- File cabinets: in good condition
- Pickup truck: good condition, four-wheel drive

STAFF DIRECTORY

651-665-9500 • fax 651-665-9409
www.greatrivergreening.org

Wiley Buck, Conservation Program Director, x16

Deb Gagner, Accounting and Office Assistant, x10

Cade Hammerschmidt, Field Coordinator, x20

Fred Harris, Lead Ecologist, x19

Mindy Holahan, Marketing Intern, x24

Dana Jensen, Project Administrator, x15

Libby Johnston, Marketing & Communications Director, x23

Deborah Karasov, Executive Director, x14

Eli Lewis, Ecological Design Assistant

Michael Lind, Restoration Technician

Pam McLellan, Development Associate, x18

Katherine Nielsen, Executive Assistant, x28

Melissa Peterson, Restoration Technician

Daniel Shaw, Landscape Ecologist/Designer, x12

Jane Stubblefield, Events and Volunteer Manager, x11

Dave Viale, Restoration Technician

Gregory Wenz, Business Director, x17

All e-mail addresses for staff are first initial last name
@greatrivergreening.org

BOARD OF DIRECTORS

Mark Bailey, Bailey Nurseries

Susan Galatowitsch, University of Minnesota

Steven J. Holmstoen, CenterPoint Energy Minnegasco

Lorrie Louder, Saint Paul Port Authority

Ann McMillan

Brad Moore, Department of Natural Resources

Dick Nowlin, Lindquist & Vennum

Deborah L. Osgood, The Osgood Group

Kristine Sundberg, The Sage Group

Alvin-o Williams, Castaneda Williams, Ltd.

Jonathan Wilmshurst

Superior Minerals Company

Printed with soy ink on 100% post-consumer content recycled paper



Sign up
for Spring events!


Great River Greening restores valuable and endangered natural areas and open spaces in the greater Twin Cities by engaging individuals and communities in stewardship of the Mississippi, Minnesota and St. Croix river valleys and their watersheds.

OUR MISSION

Nonprofit Org.
U.S. Postage Paid
Saint Paul, MN
Permit #6864

GREAT RIVER GREENING
35 West Water Street, #201
Saint Paul, MN 55107





The River Valley Reader

It's habitat forming

Volunteers still the "stars" of Great River Greening

"GREENING THE GREAT MISSISSIPPI---A TINY NONPROFIT CONVINCED 7,000 VOLUNTEERS AND 450 ORGANIZATIONS TO DIG IN AND HELP."* That was the vision of Greening back in 1995 and—with 37,000 native trees and shrubs planted, 50 acres of prairie grasses and wildflowers restored and 100 acres of buckthorn cut—it remains our vision today.

Rob Buffler, who recently left to become Executive Director of the Yellowstone to Yukon Conservation Initiative, was the founding director of Greening. He took that vision and turned it into the efficient, professional, stable organization that Greening is today. While it is hard not to miss Rob's charisma, we know that his greatest contribution was his insistence on the volunteers as the "stars" of Greening. They are the future and the constant of the Greening organization while our board of directors works to select the new executive director.

Enthusiastic new volunteers joined our ranks as we expanded our reach to work in the glorious forests of the Minnesota Valley National Wildlife Refuge and

lower Mississippi. Even more volunteers will be needed next year when we start restoration work "Greening style" in the scenic St. Croix River valley and upper reaches of the Mississippi.

Our corps of trained supervisors overseeing volunteers is now at 300 and some of them have worked with us to pilot a park stewards program in Saint Paul. When fully developed, this stewards program could build capacity throughout the Twin Cities for community-based restoration work. And now more than ever, Greening's resources are being leveraged ten-fold as we join our regional partners to restore the last remaining natural areas in the Twin Cities in a legislatively-funded effort called Metro Wildlife Corridors.

This fall, we will have new kinds of volunteers for our first fundraising event, the Great Rivers Marathon (www.GreatRiversMarathon.org), organized with the Minnesota Boat Club and benefiting Great River Greening. On September 6, you can row, canoe or kayak with hundreds of others on the Minnesota and Mississippi Rivers while they are closed to commercial traffic and help support the health of our incredible river resources by raising funds for the restoration work of Great River Greening.

To get involved in this event and all the others listed in this newsletter, please visit www.greatrivergreening.org or call our Volunteer Hotline at 651-665-9500, x2.

Please remember that none of this work is possible without generous individuals like you who have a stake in our community's natural resources. McKnight Foundation recently awarded Greening a major two-year operating grant based in part on this shared stake. We hope you will continue to support us with your time, talent and donations. Greening's community vision remains as real today as the 30-foot green canopy that grows over what was a barren spot of land on the Mississippi.

—Deborah Karasov, Interim Executive Director

* Department of Natural Resources' *Minnesota Conservation Volunteer* magazine, May June 2000.



First Heritage Park event is big success

GREAT RIVER GREENING'S MOST CHALLENGING spring planting event turned out also to be our most successful.

More than 100 volunteers of all ages and cultural backgrounds from the Heritage Park neighborhood in North Minneapolis joined 10 newly trained supervisors and their veteran mentors to plant 1,700 native seedlings, trees and shrubs. After the work was done, we celebrated everyone's contribution with a barbecue lunch complete with ice cream cones, free plants to take home, prize drawings and a mass plant watering provided by the Minneapolis Fire Department (See photo).

Heritage Park is Minneapolis' landmark housing redevelopment designed for all income levels on the former sites of public housing. Community green space is a central design feature including ponds, meadows and lawns. Greening is playing several roles in the development. We bring native plants to Heritage Park to contribute to ecological balance and beauty. Through volunteer events, we help connect Heritage residents and neighbors to their



physical environment while encouraging stewardship. For the long-term, we have trained a small group of supervisors in volunteer stewardship and greening techniques so they may lead the effort throughout the greater community.

Heritage Park does not have a history of environmental stewardship or volunteerism. Greening made a commitment in its five-year strategic plan to increase our volunteer reach in under-served communities. Our work with the City of Minneapolis and Heritage Park has been both demanding and rewarding as we reach new communities with our native restoration message.

Our experience tells us that hands-on volunteer

involvement in restoration and stewardship is one of the most effective tools for environmental education—the first step in securing a healthy future for any historically environmentally battered areas. Greening will be back at Heritage Park next spring!

Greening partners with hunters for novel restoration program

THIS FALL, GREAT RIVER GREENING will partner with the St. Croix Watershed Research Station, Kiwanis Club and the Minnesota Deer Hunters Association to teach hands-on habitat restoration to young hunters on the lands where they hunt. *Bucks and Buckthorn: Engaging Young Hunters in Restoration* will affect an important continuous corridor of undeveloped land—2,300 acres—extending from the St. Croix River toward Big Marine Lake. This corridor has been preserved through the project partners and the Department of Natural Resources Metro Greenways program.

The program is funded through a \$255,000 state grant recommended by the Legislative Commission on Minnesota Resources and a \$50,000 matching grant

from the National Fish & Wildlife Foundation. In announcing the matching grant, Congresswoman Betty McCollum (MN-04) said, "I commend Great River Greening for their dedication to stewardship of our natural resources for our families and for the wildlife which inhabit them."

Governor Tim Pawlenty lent his support by attending the first day of the youth training camp in August.

Bucks and Buckthorn provides hunters and conservation volunteers the opportunity to work together for a common goal—critical habitat restoration. *Bucks and Buckthorn* partners hope to pilot the program throughout the state as a novel way to introduce a conservation ethic to youth.



Volunteers Field Guide

Opportunities to help green the Twin Cities

Save

Get out and enjoy the colors—Volunteer for Greening's fall events!

Battle Creek Regional Park, Saint Paul

TUESDAY, SEPTEMBER 2 • 8:15 A.M. – 11:45 A.M.
AND 12:15 P.M. – 3:30 P.M.

Only Supervisors needed: 4

150 MACALESTER COLLEGE STUDENTS will spend their first day of school volunteering for Greening as part of their "Into the Streets" community service program. Greening crew and Supervisors will lead these students in hauling and stacking already-cut invasive trees and shrubs. The restoration plan will reconnect high-quality natural communities and viable wildlife corridors, as well as add to the heritage of the site and surrounding landscape. *We need only supervisors for this event, but note that it is on a weekday.*

Great Northern Business Center, Saint Paul

SATURDAY, SEPTEMBER 20 – 9:00 A.M. – NOON

Volunteers needed: 60

*Field – 60 Supervisors – 6 Parking – 2 Registration – 4
(Please indicate your preference when registering)*

CO-SPONSORED BY THE SAINT PAUL PORT AUTHORITY, the project is located adjacent to a light industrial development that is incorporating ecological design principles and native plant species. Volunteers will plant shrubs, grasses and flowers on property between the development and Como Ave. Since work will be next to Como Ave., this event is not appropriate for small children.

Land 'O Lakes, Inver Grove Heights

FRIDAY, SEPTEMBER 26 • 9:00 A.M. – NOON

Only Supervisors needed: 3

UNIVERSITY OF MN STUDENTS AND LAND 'O LAKES employees will plant trees, shrubs and plugs in a beautiful area with a river view just off of Highway 52. *We need only supervisors for this event, but note that it is on a weekday.*

Minnesota Valley National Wildlife Refuge

SATURDAY, OCTOBER 11 • 9:00 A.M. – NOON

Volunteers needed: 50

Field – 50 Supervisors – 5

SPEND A BEAUTIFUL FALL AFTERNOON HELPING RESTORE flooded farmland to bottomland hardwood forest in the floodplain between Chaska and Carver. This 600-acre unit of the Refuge consists of a marsh-edged lake surrounded by farmland and floodplain forest. The intent of this restoration is to encourage many different bird species to use the area as a nesting spot during spring migration. Help plant 200-300 oaks and perhaps remove some buckthorn – then enjoy a picnic lunch on site. Be prepared for a three-fourths-mile walk to the worksite.

Flint Hills Resources, Rosemount

SATURDAY, OCTOBER 25 • 9:00 A.M. – NOON

Volunteers needed: 70

*Field – 70 Supervisors – 10 Parking – 4 Registration – 4
(Please indicate your preference when registering)*

RESTORATION OF THE PINE BEND BLUFFS NATURAL AREA along Highway 52 near 117th Street continues as Great River Greening, Friends of the Mississippi River (FMR) and Flint Hills Resources (FHR) host another event to remove exotic species from the site's oak savanna. Our goal for the site is to maximize native plant diversity by controlling invasive species and replanting native species. Volunteers will work alongside FHR employees and FMR volunteers to remove buckthorn and haul and stack already-cut brush and trees on a seven-acre parcel on top of the bluff. This is a family-friendly event and lunch will be provided. After lunch, and weather permitting, there will be a guided tour to see the sand/ gravel prairie restoration.

You must pre-register for this event by October 15 so your name will be on a check-in list for security. No last minute additions or substitutions — no one will be allowed on the property without pre-registering.

Supervisors Field Guide

Opportunities for Greening's volunteer supervisors


Supervisors are the inspiration behind volunteers

IF VOLUNTEERS ARE THE "STARS" OF GREAT RIVER GREENING, then Supervisors are the inspiration that keeps those stars shining brightly! On most Saturday mornings in spring and fall, you can find a dozen dedicated Greening Supervisors guiding and teaching volunteers in the fine arts of buckthorn removal and native species planting.

Our Supervisors are environmental enthusiasts who have completed a Basic Training course and then supervise groups of volunteers at Greening events. We are forever indebted to our Supervisor corps who makes Greening's broad community outreach possible.

Many of our Supervisors are Master Gardeners while others have received additional training directly through Greening. We have offered Advanced Training sessions on topics such as "Native Landscape Design for Homeowners" and "Prairie Restoration." These valuable training opportunities have been held at the Minnesota Landscape Arboretum, the Minnesota Valley National Wildlife Refuge and at a restored prairie. This fall/winter, our Advanced Training focus is on the "feathered benefactors" of our restoration work (See following article). Join us for two entertaining and informative sessions!

In June, Greening Supervisors played a special role in our work in the new North Minneapolis Heritage Park community. Seasoned Supervisors were paired with newly trained recruits from Heritage Park. New friendships were formed as teams of Supervisors guided the planting



"When I come back in a couple of years from now the plants are going to be all grown up and I can say that I planted those."

"Thank you for trusting us to use your tools, and have a blessed day!"

"I want to thank you for having those special helpers (Supervisors) because they really taught us a lot."

"I'm glad that you had some really nice people to help us."

"You showed me how to plant and helped me by teaching me how to help save the environment."

"I'm glad I got to plant with people who know a lot about plants."

process around a stormwater pond in the landmark redevelopment. Greening Supervisors were on the job, training new community leaders in environmental stewardship and greening techniques.

To those of you who are already Supervisors – we know you have fun at our events. But do you ever wonder if your generous donation of time and talent really makes a difference? After a planting event on Harriet Island last fall, sixth grade students from Saturn Academy in Saint Paul sent us thank-you notes. Their letters speak volumes about the value Greening Supervisors bring to the community and the environment.

We salute all our Supervisors and want them to know how much we appreciate their contribution to our mission!

If you have taken Basic Training for Supervisors but have not yet participated in an event, please join us this fall. And of course, we welcome back our veteran Supervisors. See the event schedule on page 3.



Advanced training for Supervisors focuses on feathered friends

Bird Migration and Native Habitat

SATURDAY NOVEMBER 1 • 9:00 A.M. – NOON,
MINNESOTA VALLEY NATIONAL WILDLIFE REFUGE,
BLOOMINGTON

WHO BENEFITS FROM OUR RESTORATION WORK? Find out when Mark Martell, Director of Bird Conservation at the Minnesota Chapter of the Audubon Society, talks about the birds who use the Mississippi flyway as their migration route. Learn how our restoration work contributes to a high-quality habitat and how you can get involved in the annual worldwide Christmas bird count. Weather (and birds) permitting, we will join Mr. Martell for a birding hike at the Refuge during the last half of his presentation.



Raptor Rehabilitation and Native Habitat

SATURDAY APRIL 3 • 10:00 AM – NOON,
RAPTOR CENTER, SAINT PAUL CAMPUS OF THE
UNIVERSITY OF MINNESOTA

OUR "FEATHERED FRIENDS" SERIES CONTINUES with an educational tour of the Raptor Center. Learn how veterinarians rehabilitate majestic birds of prey that are found along the Mississippi River flyway. This is a rare opportunity to get a close-up look at live eagles, hawks, owls and falcons. You'll also learn more about the role of raptors in the environment and the habitats that raptors call home – habitats you've helped create as a Greening Supervisor!

These classes are open to Greening Supervisors and employees of Greening partners. Advance registration is required and space is limited. To register, please call our Volunteer Hotline at 651-665-9500 x2 or register online at www.greatrivergreening.org.

The River Steward Review

Tips for managing your investment in native plantings

New Book: *Plants for Stormwater Design*

TREATING STORMWATER WITH NATIVE PLANTS is a relatively new technique to improve the water quality of our region's rivers and lakes. The first book to discuss thoroughly methods to achieve beauty, functionality and clean water through landscape design is now available.

Plants for Stormwater Design, Species Selection for the Upper Midwest, written by Dan Shaw, Great River Greening Ecologist and Designer, and Rusfy Schmidt, Landscape Ecologist at URS, guides designers through the selection of plants for a variety of stormwater systems ranging from rainwater gardens to wetlands and detention basins.

Plants for Stormwater Design will be available after September 15. Contact the Minnesota Pollution Control Agency at (651) 297-8679 or write to: Operations and Environmental Review Section, Regional Environmental Management Division, MPCA, 520 Lafayette Rd. N., Saint Paul, MN 55155-4194.

To our River Steward friends:

THE RAPIDLY EXPANDING DEVELOPMENT of Saint Paul's West Side makes it critically important to refurbish your plantings and ensure their long-term sustainability. At Great River Greening, we encourage landowners not only to maintain their original plantings, but also to expand them whenever possible. Please contact us to learn more about protecting your investment in native plantings. We offer technical assistance to landowners in the original project area.

Post an interpretive sign to spread the word to your employees and visitors that native landscaping benefits the entire river valley. Businesses on the West Side Flats of Saint Paul that have native plantings installed by Great River Greening can obtain a 12" x 18" sign from Greening at no cost. Contact Laura Bates (See staff directory on page 8).

The River Steward Review

Tips for managing your investment in native plantings

Prairie and woodland ecosystems offer winter “room and board”

WINTER IN THE UPPER MIDWEST IS A QUIET SEASON when plant life enters dormancy and many animals go into various levels of hibernation. While things may seem desolate, the natural world is actually alive and well, enduring the winter and preparing for spring. Our local prairie and woodland ecosystems house numerous species of mammals, birds and insects throughout the long, cold months of winter.

Many animals find protection and food in the prairie. The prairie is home to mammals such as minks, weasels and ground squirrels. Great horned owls hunt field mice burrowing in the snow and rabbits that dash across the grassland.

Birds count on seeds available above the snow on grasses like big bluestem (*Andropogon gerardii*) and indian grass (*Sorghastrum nutans*) and on forbs such as goldenrod (genus *Solidago*) to sustain them through the season. The rose hips of wild roses (*Rosa blanda*) are also a good food source for many animals. The tall grasses of the prairie provide shelter and protection for birds roosting for the winter.



Urban woodland plantings offer habitat for birds that stay in Minnesota for the winter. Dark-eyed juncos, cardinals and gold finches use shrubs and trees for shelter and food. Plant species such as mountain ash (*Sorbus decora*), highbush cranberry (*Viburnum edule*), juniper (*Juniperus communis*) and staghorn sumac (*Rhus typhina*) provide berries long into the winter for the birds. These plants depend on the birds as well to help spread their seeds and sustain their populations.

Insects rely on the protection of the litter layer made up of grasses and leaves during their different stages of development. The morning cloak butterfly hibernates as an adult under tree bark during the winter and the Prometheus moth cocoons on the branches of bushes such as chokecherry.

To see an amazing array of wildlife that depends on the ecosystem to sustain them through the winter, take every opportunity to look out your window and into the snow. You never know what interesting creature may scurry or fly by!

Plant a kaleidoscope of fall and winter color

PRAIRIE GRASSES

Little bluestem (*Schizachyrium scoparium*) and **big bluestem** (*Andropogon gerardii*) – stems turn an attractive russet color in the fall and winter

Indian grass (*Sorghastrum nutans*) – seed heads have a soft, golden-yellow appearance

Switchgrass (*Panicum virgatum*) – small pink seeds and yellow foliage

PRAIRIE FORBS

Aster (genus *Aster*) – blooms late summer/early fall in a variety of colors--purple, pink, white

Stiff goldenrod (*Solidago rigida*) – yellow flowers bloom into late fall

Butterfly milkweed (*Asclepias tuberosa*) – unique seed pods with light, silky seeds

Blazing star (genus *Liatris*) – late summer bloomer produces purple flowers that attract butterflies

SHRUBS

Highbush cranberry (*Viburnum edule*) – orange-red berries persist through the winter

Nannyberry viburnum (*Viburnum lentago*) – dark purple berries in the fall

Common juniper (*Juniperus communis*) – blue-green berries endure the winter

Wild rose (*Rosa blanda*) – orange-red rose hips; leaves stay green through late October

TREES

Downy hawthorn (*Crataegus mollis*) – red berries produced through late September

Red cedar (*Juniperus virginiana*) – blue-grey berries and rusty green foliage in the winter

Sugar maple (*Acer saccharum*) – yellow, orange and red fall foliage

White pine (*Pinus strobus*) – evergreen with dark green needles

These River Steward Tips are made possible through the generosity of the Saint Paul Foundation, Katherine B. Andersen Foundation, F.R. Bigelow Foundation and the City of Saint Paul.

Thank you Greening Sponsors!

WITH A GENEROUS GIFT OF \$5,000, Wells Fargo recently became a Great River Greening sponsor. "It's important to the employees of Wells Fargo and the Corporation to give back to the communities we're a part of—it's part of our culture," said Paul Seiferth, Wells Fargo Premier Banking Manager. Thank you to Wells Fargo for helping Greening.

Ford Motor Company's Twin Cities Assembly Plant also became a Great River Greening sponsor this year with a \$5,000 contribution. Ford's efforts to restore and preserve the Mississippi bluffslands on its Saint Paul property make it an important steward of the Mississippi River valley.

In April, more than 60 Ford employees helped remove invasive species and plant over 450 trees and shrubs. We thank Ford and Local 879 for their support at this event.

The Saint Paul Port Authority, long a sponsor and partner of Great River Greening, became one of Greening's first corporate sponsors with a gift of \$6,500. Since it was organized in 1932, the Port has redeveloped polluted and underutilized land into homes for growing businesses. Greening thanks the Port Authority for these "green gifts" to Saint Paul.

Thank you to Greening's other corporate sponsors:

Bailey Nurseries	MSP Development
Leonard, Street and Deinard	Hawkins Chemical
Custom Dry Wall	

And, a special thanks to those who have come on board to date as Great Rivers Marathon sponsors:

Supporting Sponsors:

Alter River Terminal	Briggs and Morgan, P.A.
Centerpoint Energy Minnegasco	Loucks Associates
Saint Paul Riverfront Corporation	

Associate Sponsors:

Asset Recovery Corporation	Aggregate Industries
Ceridian Corporation	Rexam Beverage Can Co.
Saint Paul Port Authority	SKB Environmental
Wabasha Business Center	

To learn more about Great River Greening sponsorship, contact Pam McLellan, Development Associate, 651-665-9500 x18, pmclellan@greatrivergreening.org.

For the Birds!

Great River Greening thanks Youth Frontiers, Inc. for their generous donation of seven wren nesting boxes and ten bird feeders, constructed out of recycled wood by youth in the *Make a Stand* program. They are being awarded to volunteers at restoration events.



Juanita the Warbler

EVERY FALL, I FLY ALONG the Mississippi River valley from my summer home in Minnesota to my winter home in Central America. When I look down at the water, I can see big fish. Really big fish. It turns out that the five biggest fish species in Minnesota live in the Mississippi River.

Lake sturgeon: These prehistoric monsters of the deep can weigh over 100 pounds. Twenty years ago, they had practically disappeared from the Mississippi River because of pollution. Now that the water is cleaner, they are back. Sturgeon look like sharks but they have no teeth. They eat tiny insects.

Flathead catfish: These fish have huge heads and can weigh over 70 pounds. They are called catfish because it looks like they have whiskers. Actually, these are organs they use to smell food.

Muskellunge: These ferocious predators eat other fish. The largest caught in Minnesota weighed over 54 pounds. Anglers who fish for muskies use giant lures as big as your foot.

Carp: These fish were imported to Minnesota from Germany in 1881. They grow to over 50 pounds. They eat mainly insects but will eat corn, bread and other tasty baits.

Northern pike: Like muskies, northern pike are toothy meat eaters that hunt other fish for food. Their long, streamlined bodies make them fast swimmers. Minnesota's largest northern pike was about 45 pounds.

Not all Mississippi River fish are big. Tiny minnows swim down there. And so do little fish called darters.

The next time you look down at the Mississippi River, imagine the 33 different kinds of fish that live there. And, look out for that big catfish!

—As told to Thomas Ibsen, Mississippi National River and Recreation Area, National Park Service.
Illustration by Nora Wildgen.



Support Greening!

Your tax-deductible gift will help Great River Greening restore valuable and endangered natural areas along the Mississippi, Minnesota and St. Croix river valleys and their watersheds. Donate online at www.greatrivergreening.org or send this form and check payable to Great River Greening, 35 West Water Street, Saint Paul, Minnesota 55107

Enclosed is my tax-deductible gift of: ☐\$25 ☐\$50 ☐\$100 ☐\$150 \$_____

Name _____

Organization _____

Phone (h) _____ Phone (w) _____

Address _____

City _____ State _____ Zip _____

E-mail address _____

☐ Please do not include my name in the annual list of donors.

THANK YOU FOR YOUR SUPPORT!

9/03

Greening receives Garden Club of America award

IN JUNE, GREAT RIVER GREENING WAS AWARDED a special non-member award for conservation work, by the Saint Paul Garden Club at the annual Midwestern zone meeting and flower show of the Garden Club of America. The award recognizes "the significant contribution to conservation" by the original Greening the Great River project as well as Greening's continued accomplishments in conservation and restoration.

Greening also won the Marion Thompson Fuller Brown Conservation Award for an outstanding conservation exhibit at the show. The exhibit presents the Greening story and demonstrates the ecological value and beauty of native restoration and plantings. Thank you to one of our very talented interns, Courtney Skybak, for designing the exhibit.

We also want to thank Saint Paul Garden Club members who were early and active supporters of Greening. We continue to be grateful for your interest and support!

STAFF DIRECTORY

651-665-9500 • fax 651-665-9409
www.greatrivergreening.org

Laura Bates, *Lead Event Technician*, x13

David L. Cathcart, *Assistant Director*, x16

Jennifer Erickson, *Restoration Intern*

Cade Hammerschmidt, *Field Manager*, x13

Fred Harris, *Lead Ecologist*, x19

Libby Johnston, *Marketing & Communications Director*, x23

Andy Jones, *Restoration Technician*

Deborah Karasov, *Interim Executive Director*, x14

Elijah Lewis, *Ecological Design Assistant*

Michael Lind, *Restoration Technician*

Pam McClellan, *Development Associate*, x18

Alicia Murphy, *Restoration Technician*

Katherine Nielsen, *Project Assistant*, x28

Kara Przybilla, *Restoration Technician*

Dan Shaw, *Restoration Ecologist*, x12

Tracy Rae Sokolski, *Ecological Design Intern*

Jane Stubblefield, *Events and Volunteer Manager*, x11

Gregory Wenz, *Business Director*, x17

Dan Shaw, *Restoration Ecologist*, x12

Jeff Zeitler, *Ecological Design Intern*

All e-mail addresses for staff are first initial last name
@greatrivergreening.org

BOARD OF DIRECTORS

Mark Bailey

David Boyce

Gary Gardner

Steven J. Holmstoen

Lorrie Louder

Brad Moore


Deborah L. Osgood

Jonathan Wilmschurst, *Chair*

Great River Greening restores
valuable and endangered
natural areas and open
spaces in the greater Twin
Cities by engaging individuals
and communities in steward-
ship of the Mississippi,
Minnesota and St. Croix river
valleys and their watersheds.

OUR MISSION

**G R E A T
R I V E R
G R E E N I N G**
35 West Water Street, #201
Saint Paul, MN 55107



Nonprofit Org.
U.S. Postage Paid
Saint Paul, MN
Permit #6864

Sign up
for Fall events, page 3!

Printed with soy ink on 100% post-consumer content recycled paper





The River Valley Reader

It's habitat forming

Greening Helps Shape Regional Vision for Protecting Natural Resources

THE MISSISSIPPI RIVER VALLEY from Saint Paul's gorge south to Hastings is one of the most important ecological corridors in Minnesota, one whose beauty astonished Easterners when they first saw it in the 1850s. Some 150 years later, this stretch of the river is also one of the state's most complex in terms of multiple, and sometimes conflicting, land uses.

Now the river's stakeholders—government agencies, nonprofit groups, and community and business leaders—are coming together to set priorities for making land-use decisions about this important stretch of the river. As environmental consultant to this project, known as the Mississippi River Initiative, Great River Greening completed a computer-based ecological analysis of the entire corridor, identifying the most significant natural resources and suggesting how they might be affected by various land uses.

Greening's analysis showed that land of high ecological significance is concentrated in four general areas: Vermillion River Bottoms, Pine Bend, Pig's Eye and around the confluence of the Minnesota and Mississippi rivers. These areas still support remnant natural vegetative communities, provide crucial habitat for wildlife and serve as homes to many rare plants and animals.

The Mississippi River Initiative is funded by the McKnight Foundation and coordinated by the Metropolitan Council with help from a team of consultants in urban design, landscape architecture and real-estate planning. Greening's analysis will be part of a framework the Met Council is developing with community input to help guide land-use decisions that balance growth with conservation.



Volunteers helped Great River Greening reconstruct a native oak savanna on the steep slopes of the West Side Bluff, with downtown Saint Paul as the backdrop. Nearly 300 volunteers made the most of the spring planting season by helping Greening plant native trees, shrubs, prairie grasses and wildflowers at three Mississippi River sites—the West Side Bluff, the Desnoyer Park neighborhood and the Saint Paul Port Authority's Barge Terminal 1.

See page 2 for our fall event schedule.

Preliminary information suggests that half of the corridor's high-quality natural area remnants and two-thirds of its rare plants and wildlife are located on privately owned land, with no assurance that these important assets will be preserved over the long term.

Great River Greening is participating in a similar ecological study that will help target natural areas for protection along the entire Mississippi National River and Recreation Area, 54,000 acres along the river corridor from Anoka County to the Vermillion River Bottoms in Hastings.

Halt the Invasion—Join Our Fall 2002 Events

Volunteers and Supervisors Needed!

River Gorge South Park

SATURDAY, SEPTEMBER 21 • 9 A.M.—12 NOON

Volunteers Needed: 200

Field - 200 Registration - 10 Parking - 6

Supervisors - 15–20

AT THIS EVENT, VOLUNTEERS will begin implementing a Great River Greening plan for restoring ecological balance to the Mississippi River Gorge near Minneapolis's Longfellow neighborhood. The Mississippi River Ecological Inventory and Restoration Management Plan, a collaboration of Great River Greening, the Longfellow Community Council and the Minneapolis Park and Recreation Board, evaluates the site's land cover and plant species, soil types, erosion-prone areas, trail networks and location of overlooks. It also describes short- and long-term restoration goals for key areas of the site, including planting native species, reducing erosion and removing invasive plants that are threatening the natural landscape.

Volunteers will remove buckthorn and tartarian honeysuckle, non-native species that have invaded an area along West River Road from the Ford Dam at the north end of Minnehaha Park to the railroad bridge at

27th Street. Volunteers will also help haul the brush to a site where it will be chipped then hauled away and burned. Lunch, entertainment and a prize drawing will conclude this event along the Mississippi's only true gorge.

Minnesota Valley National Wildlife Refuge

SATURDAY, OCTOBER 19 • 9 A.M.—12 NOON

Volunteers Needed: 200

Field - 200 Registration - 12 Parking - 10

Supervisors - 15–20

SPEND A BEAUTIFUL FALL AFTERNOON close to the Louisville Swamp area of the Minnesota Valley National Wildlife Refuge located 4.5 miles south of Shakopee. This 2,600-acre area has a unique mix of old fields, prairie remnants, oak savanna, floodplain forest and stone farmsteads. Help cut, haul and stack brush and buckthorn in the morning, have lunch on-site (provided by Greening) and spend the afternoon enjoying the beauty of the site's 13 miles of trails. Please be prepared to walk one easy mile from the parking lot to the work-site. A naturalist will lead the hike!

The river gorge and refuge projects have received funding from the Minnesota Environment and Natural Resources Trust Fund, as recommended by the Legislative Commission on Minnesota Resources.

Action Alert!

GREAT RIVER GREENING OCCASIONALLY NEEDS last-minute volunteer help. The work might involve an office project such as a mailing, database updates or donor calling and research. Or it could involve field-work such as preparing for a planting or hauling brush.

To reach volunteers quickly, we are creating an e-mail list of supervisors and volunteers who are willing and available to volunteer on short notice. Most projects will involve daytime hours during the work week, but others may be done from home during the evening. If you would like to be added to Greening's "Action Alert" list, please contact Jane Stubblefield at jstubblefield@greatrivergreening.org. Thanks for your flexibility!

Supervisors Needed!

Swede Hollow and Harriet Island

Greening needs supervisors *only* to help volunteers from local schools and colleges replace native plants installed at previous Greening events that did not survive because of on-site construction or natural forces. We need four supervisors to lead Macalester students at Swede Hollow on Tuesday, September 3, from 11:30 a.m. to 3:30 p.m. The Harriet Island event will take place on Thursday, October 10, in two shifts—9 a.m. to 11:30 a.m. or 12 noon to 2:30 p.m. Supervisors can sign up for one shift or both.

To volunteer at a fall event, call our Volunteer Hotline at 651-665-9500 x2, or register online at www.greatrivergreening.org. Advance registration required.

Supervisors Field Guide

Opportunities for Greening's volunteer supervisors

Fall Trainings to Highlight Restoration

MORE THAN 400 VOLUNTEERS have completed Greening's Basic Training course—the first step to becoming a supervisor. With so many supervisors eager to lead our events, Greening will *not* be offering the Basic Training this fall to certify new supervisors. Instead, we will focus on providing opportunities for our current supervisors, and we will offer three new Advanced Trainings sessions, all with a restoration theme.

The Fall Line-Up

TOPIC: Prairie Plant Restoration

TIME/DATE: 9 A.M.—1:30 P.M., SATURDAY, SEPTEMBER 14

LOCATION: Gentian Farms in Osceola and Somerset, Wisc.

DR. SHAWN SCHOTTLER, ASSOCIATE SCIENTIST at the Science Museum of Minnesota, will lead a tour of his Osceola farm, where he will talk about the harvesting, processing and combining of prairie seeds, seedbed preparation and use of equipment. He will also discuss different types of prairie restorations and what makes them successful. After a brown-bag lunch, we'll drive a short distance to Somerset to see examples of different types of restorations.

Tree Care Advisor Trainings

Greening supervisors have been invited to attend several Tree Care Advisor (TCA) training sessions. "Fall Planting," "Selecting Trees and Shrubs for Fall Interest" and "Specialized Pruning Practices" will be held from 9 a.m. to 12 noon, Saturday, October 12, at the Minnesota Landscape Arboretum. TCA's Introductory Training sessions will be held from 8:30 a.m. to 3:30 p.m. on February 22, March 1, 8 and 22, at Green Hall at the U of M, Saint Paul Campus. For more information, please contact Dave Hanson at 612-624-1226 or dlhanson@umn.edu. To sign up, call Greening's Volunteer Hotline: 651-665-9500 x2.



Supervisors Darlene Charboneau and Susan Tracy take a break during Greening's Native Shrub Identification training at the Minnesota Landscape Arboretum in June.

TOPIC: The Process of Restoration

TIME/DATE: 9 A.M.—NOON, SATURDAY, JAN. 25, 2003

LOCATION: Green Hall, U of M, Saint Paul Campus

HOW DO WE BEGIN? What do we do? Why do we do it? Restore to what? Is there only one right way to restore the land? A restoration expert will help us think "green" on a frosty winter morning and discover the answers to these and many more questions about the history and current thinking regarding the overall process of restoration.

TOPIC: Native Landscape Design for Homeowners

TIME/DATE: 9 A.M.—12 NOON, SATURDAY APRIL 12, 2003

LOCATION: Minnesota Valley National Wildlife Refuge, Bloomington

ARE YOU INTERESTED IN applying the concepts of restoration in your own backyard? A panel of experts, including Greening staff, will offer information on transforming your plot of bluegrass into a landscape of native trees, shrubs, grasses and wildflowers. Bring your questions to the experts and gather tips on selecting plants, creating storm-water gardens and working within the limitations of your property.

Controlled Burns Help Woodlands and Prairies Flourish

FIRES CONTINUE TO PERFORM VITAL FUNCTIONS in natural communities, as they have throughout history. They release nutrients from burned plants, help seeds grow by breaking their protective coats and expose the soil surface to sunlight in woodlands and prairies so native wildflowers can flourish. Oak woodlands and prairies are not just adapted to fire; they depend on it to survive.

Burned areas also attract wildlife, including deer and many kinds of birds. Fire maintains prairies that provide nesting cover for waterfowl and upland birds such as prairie chickens, upland plovers and many songbirds. In some oak and pine forests, burning can encourage oak and pine regeneration and boost acorn and pinecone production, benefiting deer, squirrels, wild turkeys and other wildlife.

In an effort to restore the natural fire cycle, natural resource managers often set fires, called *controlled* or *prescribed* burns, which are carefully watched and kept under control. Trained personnel burn parts of the woods and prairie at specified intervals to clear out weedy plants that choke out bur oaks and native wildflowers.

Controlled burns have specific objectives, are done according to carefully prepared plans and are conducted only during certain weather conditions. Trained crews use specialized equipment and are supervised by experienced fire professionals called burn bosses. Local fire departments are notified in advance of a burn. Before a fire is started, wind speed and direction, temperature, relative humidity and fuel conditions are calculated. The burn boss also takes into account smoke drift, nearby buildings, livestock and other safety factors. The burn

will be conducted only if the conditions meet the approved "prescription" for the project.

When fire is reintroduced to grassland as part of a

management program, the burn is often conducted in the spring (in early May for much of Minnesota).

Burning at this time effectively controls unwanted, early-growing European weeds that have invaded the prairie. Fall burning can also be effective. Native grassland should be burned at least once every three or four years. Although mowing can be used to manage prairie, regular mowing tends to reduce diversity, deplete the soil of nutrients and compact the earth.

Mowing and removal of encroaching brush by cutting are often used with burning to restore native prairie. Fire is the most natural, efficient and effective method of maintaining prairie and savanna.



Great River Greening's field crew conducted a controlled burn at a high-quality dry prairie that Greening is helping to restore at Belle Plaine Elementary School. Students later broadcast prairie seed they had collected before the burn.

Controlled Burns Conducted by Great River Greening

Great River Greening's field crew has received training and certification from the National Wildfire Coordinating Group and The Nature Conservancy. Over the last 18 months, Great River Greening has conducted controlled burns in Bloomington, Belle Plaine and the Pine Bend area. We have also executed burns in downtown Saint Paul, at the American Red Cross building at Robert Street and Plato Boulevard, and on the levee by the Saint Paul Downtown Airport at Holman Field.

For more information about controlled burns or upcoming burns by Great River Greening, please contact lead ecologist Ellen Fuge, 651-665-9500 x19; efuge@greatrivergreening.org.

Legislators Champion Habitat Restoration

IN A UNIQUE PROGRAM STARTING NEXT SUMMER, Great River Greening will work with the Minnesota Deer Hunters Association to engage young hunters in habitat restoration in the St. Croix River valley, thanks to a \$400,000 funding recommendation by the Legislative Commission on Minnesota Resources (LCMR). The project, called Bucks and Buckthorn, was part of a total \$49.3 million package recommended by the LCMR to the 2003 Legislature for special natural resource projects.

In addition to the Bucks and Buckthorn project, Greening has been recommended to receive other monies through two significant partnerships. One, the Metro Wildlife Corridors project, is a \$5 million multi-agency collaboration organized by the Trust for Public Land and the Department of Natural Resources for land acquisition and habitat restoration along major greenways in the metropolitan area. At least \$150,000 is earmarked for Great River Greening projects in the Mississippi River gorge and the Minnesota Valley National Wildlife Refuge. Greening will also partner with the Saint Paul Port Authority in a \$300,000 project to install native plants to enhance the Port Authority's Barge Terminal 1 facility along the Mississippi River.

Funding for these projects will not be assured until the LCMR recommendations are approved by the Legislature next session. Final legislative actions on the proposals must also be signed by the governor in May.



Volunteers Chad and Nancy Nelson helped Great River Greening and the Saint Paul Port Authority plant native trees and shrubs in June at the main entrance to Barge Terminal 1. The multi-phase project to promote environmental improvements while supporting local river industry will continue with funding recommended by the Legislative Commission on Minnesota Resources. The initial project was made possible by a Five-Star Restoration Partnership grant funded by the Environmental Protection Agency and administered through the Wildlife Habitat Council.

Natural Areas Gain Ground with Open Space Campaign

EVERY DAY IN THE TWIN CITIES about 60 acres—a space nearly the size of the Mall of America—are paved over. Yet Minnesota ranks second in the nation as a state of wildlife watchers, and 84 percent of us agree that we must act now to preserve our last remaining natural areas.

How can the Twin Cities balance regional economic and population growth with the preservation of open, green space close to where people live? By engaging citizens more deeply in public land-use decisions, says The McKnight Foundation and a dozen other organizations, including Great River Greening, that have been working on a grassroots public service campaign that will be launched in late September.

The open space campaign, which will involve advertising, direct mail and a website, will encourage Twin Cities residents to become involved in civic, municipal

and state decisions about how land is used. The website will feature kits that include facts and figures; resources and partner organizations; ways to spread the word among friends, family, neighbors and other community members; tips for contacting public officials and the media, and a list of 20 other things you can do to speak out to protect open spaces.

A keystone of the campaign is a list of 10 Twin Cities land treasures that need protection. These treasures will be announced at the kickoff, and, throughout the campaign, citizens will be invited to nominate favorite open spaces they feel are threatened.

"The loss of irreplaceable forests, wetlands, vistas and farmland will continue until we Minnesotans get personally involved in land-use decisions that are being made every day in our neighborhoods and cities," says Rip Rapson, McKnight's president. "If we cherish these special places, we have to speak up to save them."

The River Steward Review

Tips for managing your investment in native plantings

Jabbin' Joe-Pye vs. Knock 'em Down Knapweed

IT'S THAT SEASON when we have to pull on the gardening gloves and go after the weeds. Weed control helps prevent unwanted plants from threatening desirable trees, shrubs and prairie species. Weeds, whether native species or non-native exotics that have invasive qualities, compete with desirable species by shading them or by stealing valuable soil moisture and nutrients. While weeds do not necessarily kill the trees and shrubs, they can reduce growth or stress plants, making them more vulnerable to insects and diseases. The most important weeds to control are those that are non-native to this area and that are particularly invasive. These plants threaten the ecological value of the planting and may even degrade adjacent natural areas.

Hand-pulling is the preferred method of weed control because it is less damaging to the environment and human health and is effective on a wide variety of weeds. Key to fighting the spread of weeds is controlling them before they grow too large or set seed. Fast-growing herbaceous weeds can easily overwhelm your planting beds. We recommend weeding once a month. At a minimum, you should monitor weeds regularly and conduct one thorough weeding in the spring and fall.

Herbicide should not be needed in planting beds if you're diligent about hand-pulling the weeds. Occasionally, herbicides may be the best option for controlling particularly invasive weeds. Only certified and licensed personnel should select and apply the herbicide.

Two common herbaceous weeds that are likely to take up residence in your plantings are spotted knapweed and common burdock. Here are tips for knocking these weeds out of your plantings.

Spotted Knapweed (*Centaurea maculosa*)

HABIT: Biennial or short-lived upright perennial forb.

SIZE: 3-4 feet in height.

LEAVES: Alternate, pale, rough 1-3 inches in length. Leaf margins on lower leaves are divided about halfway to the midrib. Upper leaves are more linear in shape.

STEM: Slender, hairy, erect, growing in a branched pattern, two feet in height on drier sites and up to four



feet in height on moister sites.

SEEDS: 1/4 inch and brownish. Notched on one side of the base with a short tuft of bristles at the tip.

FLOWER: Lavender flower head has stiff bracts marked with fine, vertical streaks and tipped in with dark, comb-like fringes.

ROOT: Stout, elongated root.

ORIGIN: Eurasia.

Spotted knapweed attains high densities on sunny sites, reducing the presence of native species. Infestation can also contribute to poor water quality and erosion by increasing run-off and sedimentation. Plants average 1,000 seeds per plant. Seeds are viable for seven years and germinate throughout the growing season.

The most effective method for controlling spotted knapweed is to dig up or pull the entire root. (Remember to wear your gloves.)

Source: Wisconsin Department of Natural Resources, 1997. Minnesota Department of Natural Resources, 1995. United States Department of Agriculture, 1971.

Burdock (*Arctium minor*)

SIZE: 1-5 feet in height.

HABIT: Large leaves with flower heads in spike-like clusters.

LEAVES: Up to 12-14 inches across, dark green, dull, somewhat heart-shaped.

FLOWER: Small, lavender or pink, and similar in shape to thistle blossoms.

ORIGIN: Europe.

Burdock is an opportunistic species native to the United States. Extremely prolific, it will inhabit many environments disturbed by humans. Burdock produces burs about 1 inch in diameter in the fall. Burdock is aggressively opportunistic on disturbed soil and tends to shade out smaller, herbaceous flora.

Burdock can be easy to control because it reproduces only by seed and takes two years to mature. The best method for controlling burdock is to hand-pull, making sure to sever the roots below ground to kill the plant.

Source: Wisconsin Department of Natural Resources, 1997.



Volunteer Donates Time and Resources

PATSY HUBERTY'S SMILING FACE under her wide-brimmed sun hat is familiar to Greening staffers and volunteers. She has volunteered at 25 Greening events over the past five years. The first thing she did with Greening was attend a supervisor training session so she could lead volunteers at our events.

"You were doing what I believe in," says Patsy, "helping to restore the earth."

Because she believes in our work, Patsy also supports Greening with charitable gifts.

"I realized that you depended on the general public for financial support," she says.

Although Patsy's love of plants and the outdoors initially brought her to Greening, it's the people she's worked with that are most memorable. She says she likes the diversity

of groups that come to our events, and she especially enjoys the children. Ask her about the 4-year-old who cried because there were no more trees to plant or about teaching inner-city kids about wood ticks. She had so much fun supervising one group of teenagers that she wrote the group leader a thank-you note.

Patsy says that educating people about the natural landscape and building community are the most important things Greening does.

"People who work on a park are going to be protectors of that area. It becomes much more 'their' park or 'their' planting. I bring out-of-town visitors to see 'my' plantings," she says.

That sense of ownership makes Patsy want to protect what she's planted. A few weeks after



Greening supervisor Patsy Huberty (without her trademark sun hat) at an advanced training in June.

Greening's May event on the Mississippi River bluff in Saint Paul, Patsy found herself at the site again. "At the Memorial Day fireworks, I was at the High Bridge. I kept an eye on people walking along the bluff to make sure they stayed on the right side of the fence. I was worried, but they all stayed off our plantings."



Juanita the Warbler

Thank you for planting trees and wildflowers along the Mississippi and Minnesota rivers. They provide food and safe places to hide during my spring and fall migrations. You can help in other ways as well. Even from your own neighborhood!

Have you ever noticed the holes next to your curb in the street? These *storm sewers* all lead into a nearby river or lake. When it rains, water running into them carries a variety of pollutants directly into the water. There are no filters under ground to take out these poisons.

Garbage, fertilizers and other chemicals from our lawns, driveways and streets wash into our lakes and rivers and poison many of the

plants and animals that live along or in these waters. Even grass clippings and leaves can cause pollution; if too many enter the water all at once, they break down and take too much oxygen out of the water.

Here's what you can do to help!

- Don't litter, and keep the curbs along your street clean.
- Never pour anything into the storm sewer drains.
- Rake and bag your leaves and grass clippings.
- Tell your family and friends to keep their cars and lawn mowers tuned up so they don't leak gas and oil.

—As told to Thomas Ibsen, Mississippi National River and Recreation Area, National Park Service.

Illustration by Nora Wildgen.

Can You Help?

If you would like to contribute to Great River Greening, please check the appropriate box below and fill in the requested information. Please make your check payable to Great River Greening.

Enclosed is my tax-deductible contribution of:

☐ \$125 ☐ \$100 ☐ \$50 ☐ \$25 ☐ \$_____

Name _____

Organization _____

Phone (h) _____

Phone (w) _____

Address _____

City _____

State/Zip _____

E-mail address _____

☐ Please do not include my name in the annual list of donors.

THANK YOU FOR YOUR SUPPORT!

8/02

STAFF DIRECTORY

651-665-9500 • fax 651-665-9409
www.greatrivergreening.org

Laura Bates, *Restoration Technician, x25*
Rob Buffer, *Executive Director, x15*
David L. Cathcart, *Director of Operations, x16*
Ellen Fuge, *Lead Ecologist, x19*
Cade Hammerschmidt, *Restoration Technician, x29*
Deborah Karasov, *Assistant Director, x14*
Sara Halverson, *Intern*
Marjorie Hundtoft, *Intern*
Sean Jergens, *Intern*
Meredith McNab, *Communications/Education Manager, x23*
Veronika Phillips, *Landscape Designer, x28*
Adam Robbins, *Field Manager, x20*
Dan Shaw, *Restoration Ecologist, x12*
Jane Stubblefield, *Events and Volunteer Manager, x11*
Susan Troha, *Development Director, x18*
Gregory Wenz, *Business Director, x17*
Adam Zielie, *Administrative Assistant, x10*

All e-mail addresses for staff are first initial last name
@greatrivergreening.org

BOARD OF DIRECTORS

David Boyce	Joe Kingman
Charles Casperson	Lorrie Louder
Gary Gardner	Deborah L. Osgood
Steven J. Holmstoen , <i>Chair</i>	Jonathan Wilmshurst

Spring Season Initiates New Staff

A BUSY SPRING PLANTING SEASON has made Greening veterans of Jane Stubblefield, event and volunteer manager, and Laura Bates, restoration technician. Both started at Greening on March 25. Jane, who oversees all aspects of Greening's event and volunteer program, says the job combines two of her favorite things—nature and volunteers. Already she has worked with more than 400 Greening volunteers and supervisors at six trainings and events! "I love watching volunteers' expressions of accomplishment through the mud, sweat and cheers at the end of an event," she says.

As a member of Greening's field crew, Laura conducts exotic species removal, plantings and restoration site preparation. Before joining Greening, she worked for the Department of Natural Resources as a member of the Minnesota Conservation Corps State Parks Crew. "The job is rewarding," she says of Greening. "I like seeing the tangible results of our work in the field every day."

And the Award Goes to ... Greening

GREAT RIVER GREENING RECEIVED the Minnesota Shade Tree Advisory Committee's 2001 Innovation Award in March for our "innovative and exemplary model of a community-based program involving thousands of community volunteers in restoring and preserving our great river valleys."

In recognition of the groups' longstanding collaboration, the Minnesota Conservation Corps presented Greening with an Exemplary Partnership Award at its 20th anniversary celebration at the Minnesota History Center on June 24. For several years, MCC members conducted Greening's fieldwork, and our current field crew are all MCC alums.

Great River Greening is a
nonprofit, community-based
organization that exists
to help communities
restore, manage and
learn about their
natural environment through
volunteer involvement.

OUR MISSION

G
R
E
A
T
R
I
V
E
R
G
R
E
E
N
I
N
G
35 West Water Street, #201
Saint Paul, MN 55107



Nonprofit Org.
U.S. Postage Paid
Saint Paul, MN
Permit #6864

Susan Thornton

From: Susan Thornton
Sent: Tuesday, July 27, 2004 1:28 PM
To: 'Greg Wenz'; 'Dana Jensen'; 'dkarasov@greatrivergreening.org'
Cc: Sandy Smith; Susan Von Mosch; 'gail.fox@dnr.state.mn.us'; John Velin; Susan Thornton
Subject: BRP final 2001 Report

Dear Greg, Deborah and Dana,
Thank you for your final work program report for Big Rivers Partnership: Helping Communities to Restore Habitat from ML 2001, First Special Session, Chapter 2, Sec. 14, Subd. 4(i). It is my understanding that copies were sent via U.S. mail about a week ago along with several deliverables. As we discussed today, the mail has not yet arrived. Please let this email serve as acceptance of your work program final report and our request that DNR reimburse your final payment request if the billing is all in order. While, we would still like copies of the products you produced under this project we do not want to hold up your final reimbursement. We will work with you to obtain additional copies if nothing is received in the mail in the next week or so. It is our hope, however, that the package will find its way to Rm. 65 of the SOB. We wish to thank everyone for their hard work on making this project a success over the last three years. Please submit a copy of the final report, Attachment A and the cost share summary to Gail Fox at the DNR.

On behalf of LCMR staff,
Susan Thornton

Susan Thornton
Asst. Director
LCMR
Rm. 65 State Office Building
St. Paul, MN 55155
651-296-6264

FINAL

2001
4(i)

2001 Project Abstract

For the Period Ending June 30, 2004

TITLE: Big Rivers Partnership: Helping Communities to Restore Habitat

PROJECT MANAGER: Deborah Karasov

ORGANIZATION: Great River Greening

ADDRESS: 35 West Water Street, Suite 201, Saint Paul, MN 55107-2106

WEB SITE ADDRESS: www.greatrivergreening.org

FUND: Environment and Natural Resources Trust Fund (TF)

LEGAL CITATION: ML 2001, 1st Special Session, Ch. 2, Sec. 14, Subd. 4(i)

APPROPRIATION AMOUNT: \$910,000

JUL 27 2004

Overall Project Outcome and Results

Led by Great River Greening, the Big Rivers Partnership is one of the first important restoration collaborations in the state, bringing together nonprofit, government and private landowners to restore river valley habitat in the Twin Cities. Guided by ecological and resource criteria, projects were located within the important and beautiful Mississippi River Gorge running through Minneapolis and Saint Paul; the Pine Bend Bluff Natural Area, a regionally significant ecological resource on the urban Mississippi; the Minnesota River Valley, a critical and unmatched urban corridor of wetland and associated upland habitat; and numerous native plant community remnants. Projects consisted of plant and animal surveys and restoration activities that regularly engaged volunteers. More than 3,600 volunteers participated in habitat projects, triple the goal. The partnership also leveraged over \$1.3 million in non-state funds, almost double the goal, and implemented restoration on over 1,500 acres, 150% of the goal.

Project Results Use and Dissemination

Great River Greening lists projects and surveys on its website, promoting them to partners, cooperators, and landowners. Where possible, we work with cooperators to continue stewardship beyond state funding with volunteers or other community members. The Partnership also completed an ecological ranking of sites within the river valleys to complement the regionally significant areas identified by the Minnesota Department of Natural Resources and disseminated the ranking information through various conservation forums. The success of the project proves that multiple organizations can work together to achieve conservation goals.

Date of Report: July 13, 2004
Title: LCMR Final Work Program Report

Date of Next Status Report:
Date Work Program Approved: December 3, 2001
Project Completion Date: June 30, 2004
#CH34

I. PROJECT TITLE: Big Rivers Partnership: Helping Communities to Restore Habitat

Project Manager: Deborah Karasov
Affiliation: Great River Greening
Mailing Address: 35 West Water Street, Suite 201, Saint Paul, MN 55107-2016
Telephone Number: 651-665-9500 **e-mail:** dkarasov@greatrivergreening.org **Fax:** 651-665-9409

Web Address: www.greatrivergreening.org

Total Biennial Project Budget:
\$910,000 LCMR Appropriation - \$ 909,140 Amount Spent = \$ 860 Balance

Legal Citation: ML 2001, 1st Special Session, Ch. 2, Sec. 14, Subd. 4(i)

Appropriation Language: (i) Big Rivers Partnership: Helping Communities to Restore Habitat \$455,000 the first year and \$455,000 the second year are from the trust fund to the commissioner of natural resources for an agreement with Great River Greening to implement private and public habitat projects on cost-share basis in the Mississippi and Minnesota river valleys. This appropriation is available until June 30, 2004, at which time the project must be completed and final products delivered, unless an earlier date is specified in the work program.

Status of Cost-Share Requirement: Cost-share of \$691,771 will be provided for this project. Cost-share is defined as an expense that is directly related to the approved activities of the project (Big Rivers Partnership - BRP), that is not paid for with state funds, and is not an expense that is ineligible for LCMR reimbursement. Cost-share may be incurred by the project's recipient (GRG), partners (see Section VI-C), or landowners. Cost-share may include volunteer work, which would be valued using a rate of \$11.00/hr.

Documentation of cost-share is the following:

- Brief itemized description of the costs incurred (or the in-kind services provided) and their value
- Statement that certifies that the cost-share expenses reported are directly related to the approved activities of the Big Rivers Partnership.
- Statement that certifies that these expenses are NOT paid for with state funds.
- Signature by private individual landowner, or person authorized to represent the organization.

II. and III. FINAL PROJECT SUMMARY:

Overall Project Outcome and Results

Led by Great River Greening, the Big Rivers Partnership is one of the first important restoration collaborations in the state, bringing together nonprofit, government and private landowners to restore river valley habitat in the Twin Cities. Guided by ecological and resource criteria, projects were located within the important and beautiful Mississippi River Gorge running through Minneapolis and Saint Paul; the Pine Bend Bluff Natural Area, a regionally significant ecological resource on the urban Mississippi; the Minnesota River Valley, a critical and unmatched urban corridor of wetland and associated upland habitat; and numerous native plant community remnants. Projects consisted of plant and animal surveys and restoration activities that regularly engaged volunteers. More than 3,600 volunteers participated in habitat projects, triple the goal. The partnership also leveraged over \$1.3 million in non-state funds, almost double the goal, and implemented restoration on over 1,500 acres, 150% of the goal.

Project Results Use and Dissemination

Great River Greening lists projects and surveys on its website, promoting them to partners, cooperators, and landowners. Where possible, we work with cooperators to continue stewardship beyond state funding with volunteers or other community members. The Partnership also completed an ecological ranking of sites within the river valleys to complement the regionally significant areas identified by the Minnesota Department of Natural Resources and disseminated the ranking information through various conservation forums. The success of the project proves that multiple organizations can work together to achieve conservation goals.

IV. OUTLINE OF PROJECT RESULTS:

- | | | |
|--------------------|--------------------------------|--|
| • Result 1. | LCMR Budget: \$ 700,508 | Cost-Share Requirement: \$691,771 |
| | Balance: \$ 774 | Cost-Share Spent: \$ 1,339,100 |

Implement habitat projects on 1000 acres with community volunteers

This project will build on the already burgeoning partnership for community restoration in the river corridors, spearheaded by Great River Greening. The partnership will implement habitat projects evaluated through GRG's ecological inventory and analysis, using GRG's successful collaborative process of soliciting landowners, recruiting and training volunteers and volunteer supervisors, training property owners, and providing restoration and management plans. Habitat projects will include activities such as plantings of native trees, shrubs, grasses and wildflowers, removal of exotic species and prescribed burning. Project sites that we intend to implement beginning in the spring of 2002 are Koch Refinery's Pine Bend Bluff property in Dakota County, the Minneapolis Longfellow River Gorge, and Belle Plaine Prairie. A nomination process to be completed early in 2002 will identify additional projects for 2002 and 2003.

Great River Greening has distinguished itself by the success and quality of its volunteer restorations and plantings largely because of its volunteer training program. Specialized training provides volunteer supervisors with opportunities to learn many skills, including native plant identification, principles of ecological restoration, and how to organize workers in carrying out activities for habitat restoration projects. Other volunteer field workers learn about ecological restoration by participating in habitat restoration activities. This project will recruit and train 1000 field workers and 75 supervisors, and will strengthen this training by focusing on two key areas: working with diverse groups of volunteers, and teaching principles and methods for restoring native plant

communities. This project will also strengthen our collaborations with community partners and landowners to develop restoration and adaptive management plans.

Budget:	Budgeted	Current	Balance
Personnel	\$ 390,756	390,489	267
Contractual Fees	139,442	139,060	382
Supplies/Equipment	160,748	160,623	125
Operations	5,737	5,737	0
Other	3,825	3,825	0
Result 1 - LCMR Subtotal	\$ 700,508	699,734	774

See attached worksheet for detail of expenses.

Completion Date:

- a) Implement habitat projects on 1000 acres
 - 1) 500 acres; December 31, 2002
 - 2) 500 acres; June 30, 2004
- b) Recruit and train 1000 volunteer field workers
 - 1) 500 volunteers; Dec. 31, 2002
 - 2) 500 volunteers; June 30, 2004
- c) Recruit and train 75 volunteer supervisors
 - 1) 45 supervisors; Dec. 31, 2002
 - 2) 30 supervisors; June 30, 2004

Result 1 Status:

June 30, 2004 Final Report: The Partnership completed all of its quantitative goals in this result. We implemented restoration on more than 1,500 acres. More than 3,600 volunteers participated in habitat projects, including over 300 volunteer supervisors. The partnership also leveraged over \$1.3 million in non-state funds.

Geographically, our work stretched from Belle Plaine and the west end of the Minnesota Valley National Wildlife Refuge to the Mississippi River Gorge in Minneapolis to Pine Bend Bluffs near Hastings. Site selection considerations encompassed both ecological and social factors. Ecological factors included the physical characteristics of site, the condition of the surrounding landscape, the plant community goal, and long-term maintenance and monitoring needs. Social considerations included screening/visibility, aesthetic needs, environmental needs, and sufficient management resources. Project sites included: East Mississippi Bluff-Desnoyer unit, Pine Bend Bluffs in Dakota County, Hastings Riverfront, Mississippi River Gorge-South Park in Minneapolis, West Side Bluff-Cherokee Park in Saint Paul, Minnesota Valley National Wildlife Refuge, River Park-Brooklyn Park, South Saint Paul Levee, and Prospect Crest in Saint Paul, to name a few.

The involvement of thousands of volunteers in our habitat projects helps to ensure that restoration will continue to occur far beyond the end of the BRP program funding period. For example, the Belle Plaine Elementary School Prairie is an integral part of the school. Our recent volunteer planting event at River Park, Brooklyn Park drew a tremendous response from the community, tapping into their volunteer spirit and providing educational opportunities for the volunteers, as well as direct habitat improvement on the Mississippi River bank.

As noted above, the Partnership leveraged over \$1.3 million in non-state funds. These dollars demonstrate the commitment from the community for such important work. Large contributors of cost-share included the Mississippi National River and Recreation Area (MNRRA), the National Fish and Wildlife Foundation (NFWF), and the City of Saint Paul.

- **Result 2. LCMR Budget: \$ 128,762**
Balance: \$ 56

Prioritize sites for ecological restoration and management

In partnership with the Minnesota Department of Natural Resources (DNR), National Park Service (NPS), and others, Great River Greening (GRG) will evaluate ecological inventory data of portions of the urban/suburban stretches of the Mississippi and Minnesota River valleys and habitat connections not covered by the Minnesota County Biological Survey (MCBS). MCBS looks only at the highest quality natural areas, not at those areas that could or should be restored. Using this GIS-based, landscape-scale ecological inventory data and social and economic criteria (e.g. landowner participation, BRP partner involvement, local community planning and interest), high-priority restoration project areas will be identified and mapped. Potential users of this mapped information will be municipal parks programs, DNR, NPS – Mississippi National River and Recreation Area, Trust for Public Lands, Friends of the Mississippi River, Friends of the Minnesota Valley and others.

In addition, we will conduct detailed site surveys at high-priority sites to collect information necessary to guide ecological restoration. These surveys will characterize infestations of exotic species, abundance of native species, social uses of sites, and other important factors.

Budget:	Budgeted	Current	Balance
Personnel	\$ 117,839	117,839	0
Contractual Fees	5,000	4,980	20
Supplies/Equipment	3,911	3,875	36
Operations	2,012	2,012	0
Other	0	0	0
Result 2 - LCMR Subtotal	128,762	128,706	56

See attached worksheet for detail of expenses.

Completion Date:

- a) Combine ecological inventory data with social and economic criteria to establish a prioritized map of restoration project areas
 - 1) Identify and map high-priority restoration project areas along the Mississippi and Minnesota River corridors; December 31, 2002 (mailed under separate cover 12/31/03)
- b) Conduct at least 4 detailed site surveys
 - 1) Conduct 2 surveys; December 31, 2002 (mailed under separate cover 12/31/03)
 - 2) Conduct 2 surveys; June 30, 2004 (attached)

Result 2 Status:

June 30, 2004 Final Report:

Great River Greening completed a land cover inventory using the Minnesota Land Cover Classification System for the portion of the Mississippi National River and Recreation Area (MNRRA) upstream of the Lower Saint Anthony Lock and Dam. In addition, Greening developed an ecological ranking protocol for MNRRA. The National Park Service, in conjunction with Greening, produced a CD-ROM which contains information on the ecological value of undeveloped land within MNRRA. The information provided on the CD, when used in combination with other regional or local data, can help identify open space protection opportunities (OSPO) within the river corridor.

Greening, in conjunction with the City of Saint Paul, also completed an ecological inventory (included in the larger restoration management plan) at Cherokee Park Prairie in Saint Paul. The plan makes recommendations, based upon the inventory, for the ongoing management of the vegetation to meet ecological goals and social needs. Recommendations include plantings of native plant species, actions to reduce/re-direct trails and prevent erosion, and the removal and control of invasive plant species that are degrading the ecological health of the bluff.

Great River Greening ecologists conducted surveys for four Mississippi River islands owned by the National Park Service in the BRP prioritized areas. Following an initial assessment by a Department of Natural Resources ecologist, Greening focused on these islands due to the immediate threat to their ecological quality from the invasion of exotic species, severe erosion, and/or human use. Two islands were located adjacent to Inver Grove Heights and two were adjacent to Anoka. For each island, Greening completed species lists and documented concentrations of invasive species, hydrologic conditions, soils, and structural conditions.

Greening ecologists also conducted a survey of plant communities in 68 acres of Cherokee Park (aka West Side Bluff Sector 1) that will guide restoration and management activities in the area adjacent to the oak forest and prairie remnant. This will further assist in the restoration, management and protection of these critical bluffland remnants.

In conjunction with Saint Paul Parks and Recreation, Great River Greening conducted a site survey of current trail conditions at the Bluff Trail at Crosby Park and created a design plan to guide improvements. The plan will help the City of Saint Paul manage the problematic Bluff Trail and will also act as a model for similar projects in the Twin Cities area.

• Result 3. LCMR Budget: \$ 80,730

Balance: \$ 29

Create a landowner stewardship support program.

Stewardship support and information sharing are critical to the efficiency and effectiveness of restoration. Technical information, on-going encouragement, and key stewardship services are essential to sustain current and future restoration efforts. Through a GRG River Steward Program, we will interpret and distribute relevant information through an information network (e.g. GRG website) to participating landowners, project cooperators, and natural resource managers. The River Steward Program will also provide additional services to selected participating landowners, including monitoring the condition of project sites and recommendations for volunteer stewardship activities.

Budget:	Budgeted	Current	Balance
Personnel	\$ 72,573	72,573	0
Contractual Fees	6,000	5,971	29
Supplies/Equipment	1,021	1,021	0
Operations	1,136	1,136	0
Other	0	0	0
Result 3 - LCMR Subtotal	\$ 80,730	80,701	29

See attached worksheet for detail of expenses.

Completion Date:

a) Compile information for distribution and install on website.

- 1) Compile technical information and install on website; June 30, 2002
- 2) Update information and website; December 31, 2002
- 3) Update information and website; June 30, 2003
- 4) Update information and website; June 30, 2004

b) Select and monitor at least 4 sites, and provide recommendations for volunteer stewardship activities.

- 1) Complete for 2 sites; Dec. 31, 2002. (mailed under separate cover 12/31/03)
- 2) Complete for 2 sites; June 30, 2004 (attached)

Result 3 Status:

June 30, 2004 Final Report:

We have updated and added information to our website. We developed a new and updated interactive project map with all our projects categorized by type of work. We added three management plans to the resources page: Bloomington Bluffs, Eagle Creek and West Side Bluff Action Plan Phase I, as well as fact sheets on storm water management. We also added the East-Central Minnesota Species list and a news page with the past year's project news releases and monthly e-postcards to update our constituents on Greening activity. We also made sign-up for Greening volunteer events easier with on-line registration. Technical information on Greening's website is now comprehensive and we have established a process for updating and expanding it. We have already notified our partners and other interested organizations of the website resource.

In 2002, Greening ecologists completed recommendations for stewardship activities for two important sites. Belle Plaine prairie is one of eleven remaining small native prairie remnants identified by the Minnesota County Biological Survey's 1995 inventory of Scott County. Pine Bend Bluff Natural Area is a unique natural area in the metropolitan area because of its size and quality, and a large portion is owned by Flint Hills Resources (FHR). For both of these sites, specific volunteer activities were identified within a larger stewardship plan, and volunteer groups were identified (school and company employees, respectively). Tasks for volunteers included cutting invasive species, replanting with native species, periodic weeding, and girdling trees.

In 2003, Greening finished an inventory and survey, included within a larger management plan, for the remnant savanna and forest in Cherokee Park (aka West Side Bluff Sector 1). This document provides guidance on how to incorporate volunteers in future planting/restoration events with West Side Citizens Organization and Saint Paul Parks and Recreation.

Working with employee volunteers at Flint Hills Resources, a Greening ecologist outlined and recommended 2004 activities for monitoring exotic plant population numbers and flowering, prairie seed

ripening, and acorn mast ripening, collection and storage. All of the Greening ecologist's recommendations have been accepted and are an integral part of the on-going restoration.

V. TOTAL PROJECT BUDGET:

All Results: Personnel	\$581,168
All Results: Contractual Fees	150,442
All Results: Supplies/Equipment	165,680
All Results: Operations	8,885
All Results: Other	3,825
Total Project Budget	\$910,000

See attached worksheet for detail of expenses.

In order to meet the cash flow needs of GRG, LCMR will advance \$60,000 to GRG at the beginning of the Big Rivers Partnership contract with the State of Minnesota. This amount is the most that GRG estimates it will need in any given month to cover expenses that are eligible for reimbursement by the State (see attached sheet detailing GRG's estimate of cash flow needs). In order to mitigate the interest lost by the State through this advance, GRG will hold any unused portions of the advance in an interest-bearing account that is transferable on demand to their checking account, and will deduct any interest earned on the balance each month from their requests for reimbursement. As the contract comes to a close, GRG will perform reconciliation in order to use the \$60,000 to pay the final reimbursements of the \$910,000 contract.

Final Report Status: We are submitting the final bill for \$130,003.06, of which \$60,000 has been advanced to us, and we received \$5.43 in interest on the advance in the month of June, 2004. Therefore we are requesting a final payment of \$130,003.06, less \$60,000, less \$5.43, which comes to \$69,997.63.

VI. PAST, PRESENT, AND FUTURE SPENDING:

A. Past Spending

GRG will serve as project manager for the proposed collaboration, which builds on the past work and present organization of the GRG project and its many partnerships. The budget numbers below relate to LCMR funding of the GRG project since its inception. In particular, this project builds on the successful Big Rivers Partnership, funded by an \$800,000 appropriation from the LCMR last biennium and matched by at least \$374,000 through public and private sources. The LCMR appropriation also leveraged \$50,000 of funds from the National Park Service for inventory work. In addition to the work sponsored by LCMR, this partnership of eleven government and community collaborators has stimulated hundreds of thousands of dollars of future projects. (LCMR History: \$1.1 million RIM Critical Habitat Match & LCMR appropriation; Non-LCMR History: \$2.2 million foundation, corporate, and private contributions.)

B. Current and Future Spending

In terms of future spending, this project will result in a database of prioritized future projects for restoration and management, which could not be included in the project period. The private landowners will provide matching resources. As well, the National Park service, which is one of our partners, awards approximately \$150,000 in planning grants to communities within the Mississippi Natural Resources and Recreational Area corridor; communities may leverage these funds with GRG resources. The National Park service also works with the MN DNR to award nearly half a million dollars in land use planning funds for the Mississippi Critical Areas Program; once communities complete planning they will be in a position to implement perhaps with GRG's assistance.

This proposal is also a complementary effort to the nearly \$6 million of proposed money for the Metro Greenways program. While that program focuses on planning, acquisition, and land protection, this project will stimulate on-the-ground stewardship expertise and work. For example, while the Greenways program may provide grants to local communities to complete resource inventories, this project may assist communities in implementing habitat improvements.

C. Project Partners

Great River Greening

MN Department of Natural Resources, Metro Region (in-kind contribution of ecological services; also technical support and housing for GIS ecological inventory)

Nelson French, Lori Nelson, Holly Buchanan, Kevin Bigalke, Friends of the Minnesota Valley

Whitney Clark, Tom Lewanski, Friends of the Mississippi River

Arne Stefferud, Metropolitan Council (in-kind contribution to coordinate implementing agencies)

Kate Hanson, Nancy Duncan, National Park Service (in-kind contribution of technical support for GIS)

Judy Barr, Rebecca Stenberg, City of Saint Paul (recipient of restoration services)

Cordelia Pierson, Trust for Public Land (in-kind contribution of landowner outreach)

Greg Mack, John Moriarty, Ramsey County Parks and Recreation (in-kind contribution of ecological services; also recipient of restoration services)

Rick Schultz, U. S. Fish and Wildlife Service (in-kind contribution for partnership development and planning; also recipient of restoration services)

Private landowners

D. Time

Restoration and planning require a minimum of two entire field seasons. We will begin the project January 1, 2002 and end it June 30, 2004.

VII. DISSEMINATION:

Volunteer events and supervisor training sessions are announced and reported on through the GRG newsletter and website. The map of the project area with high-priority project areas identified will be provided to the members of the Big Rivers Partnership. The Great River Greening website will be the primary tool for dissemination of landowner stewardship support and other information resulting from this grant.

VIII. LOCATION:

The Mississippi-Minnesota river corridors and habitat connections in the seven-county metropolitan region.

IX. REPORTING REQUIREMENTS:

Periodic work program reports will be submitted not later than June 30 and December 31, 2002, and June 30 and December 31, 2003. A final work program report and associated products will be submitted by June 30, 2004, or by the completion date as set in the appropriation.

Project Number: #CH34

2001 Special Session Minnesota Laws Chapter 2, Section 14, subd. 4(i)

Funding Appropriated: \$910,000

	2004 Jun	Result 1: Implement habitat projects on 1000 acres with community volunteers				Result 2: Prioritize sites for ecological restoration and management				Result 3: Create a landowner stewardship support program				Project Total		
		Budget	Current Invoice	Previously Submitted	Current Balance	Budget	Current Invoice	Previously Submitted	Current Balance	Budget	Current Invoice	Previously Submitted	Current Balance	Budget	Current Invoice	Current Balance
Personnel																
Project Manager	-	14,166.00		14,166.00	-	14,167.00		14,167.00	-	14,167.00		14,167.00	-	42,500.00	-	-
Project Assistant	2,227.58	41,230.00	774.16	40,455.84	-	27,485.00	726.72	26,758.28	-	27,485.00	726.70	26,758.30	-	96,200.00	2,227.58	-
Field Coordinator	1,068.00	58,147.63	1,068.00	57,079.63	-	-		-	-	-		-	-	58,147.63	1,068.00	-
Ecologist	4,955.00	41,978.00		41,978.00	-	76,186.64	3,463.46	72,723.18	-	30,921.00	1,491.54	29,429.46	-	149,085.64	4,955.00	-
Volunteer Program Coordinator	1,010.00	88,592.41	1,010.00	87,582.41	-	-		-	-	-		-	-	88,592.41	1,010.00	-
Landscape Designer	1,571.00	13,256.91	1,571.00	11,418.91	267.00	-		-	-	-		-	-	13,256.91	1,571.00	267.00
Restoration Technician	2,899.94	133,385.54	2,899.94	130,485.60	-	-		-	-	-		-	-	133,385.54	2,899.94	-
Subtotal		390,756.49	7,323.10	383,166.39	267.00	117,838.64	4,190.18	113,648.46	-	72,573.00	2,218.24	70,354.76	-	581,168.13	13,731.52	267.00
Contractual Fees																
Field Crews	40,181.90	93,641.59	40,181.90	53,077.06	382.63	-		-	-	-		-	-	93,641.59	40,181.90	382.63
Consulting Ecologist	4,980.00	-		-	-	5,000.00	4,980.00		20.00	-		-	-	5,000.00	4,980.00	20.00
Web Page Consultant	5,449.50	-		-	-	-		-	-	6,000.00	5,449.50	522.00	28.50	6,000.00	5,449.50	28.50
Community Liaison (Friends of Miss)	-	27,500.00	-	27,500.00	-	-		-	-	-		-	-	27,500.00	-	-
Community Liaison (Friends of Minn)	-	18,300.00	-	18,300.00	-	-		-	-	-		-	-	18,300.00	-	-
Subtotal		139,441.59	40,181.90	98,877.06	382.63	5,000.00	4,980.00	-	20.00	6,000.00	5,449.50	522.00	28.50	150,441.59	50,611.40	431.13
Supplies & Equipment																
Field Supplies	2,409.27	11,037.48	2,409.27	8,628.21	-	-		-	-	-		-	-	11,037.48	2,409.27	-
Volunteer Event Supplies	1,570.37	13,166.45	1,570.37	11,596.08	-	-		-	-	-		-	-	13,166.45	1,570.37	-
Office Supplies	3.67	1,853.00	0.73	1,845.19	7.08	736.00	1.84	697.82	36.34	353.00	1.10	351.43	0.47	2,942.00	3.67	43.89
Field Equip/Tools	4,416.41	15,505.38	4,416.41	11,088.97	-	1,971.16		1,971.16	-	-		-	-	17,476.54	4,416.41	-
Volunteer Events Equip/Tools	3,116.02	4,500.10	3,116.02	1,384.08	-	-		-	-	-		-	-	4,500.10	3,116.02	-
Office Equip/Tools	-	78.67	-	78.67	-	-		-	-	-		-	-	78.67	-	-
Field Equip Rental	-	1,754.59	-	1,754.59	-	-		-	-	-		-	-	1,754.59	-	-
Volunteer Event Equip Rental	1,142.16	7,615.76	1,142.16	6,473.60	-	-		-	-	-		-	-	7,615.76	1,142.16	-
Field Equip Maint/Repair	-	246.33	-	246.33	-	-		-	-	-		-	-	246.33	-	-
Vehicle Rental	-	11,068.30	-	11,068.30	-	-		-	-	-		-	-	11,068.30	-	-
Vehicle Upkeep	175.76	1,142.75	175.76	966.99	-	-		-	-	-		-	-	1,142.75	175.76	-
Plants/Seeds	47,997.77	79,491.92	47,997.77	31,494.15	-	-		-	-	-		-	-	79,491.92	47,997.77	-
Chemicals	4,646.88	9,615.83	4,646.88	4,851.34	117.61	-		-	-	-		-	-	9,615.83	4,646.88	117.61
Printing	-	2,085.00	-	2,085.00	-	828.00	-	828.00	-	397.00	-	397.00	-	3,310.00	-	-
Copying	-	387.90	-	387.90	-	191.50	-	191.50	-	183.00	-	183.00	-	762.40	-	-
Computer Software	-	368.00	-	368.00	-	-		-	-	-		-	-	368.00	-	-
Educational Materials	-	463.00	-	463.00	-	184.00	-	184.00	-	88.00	-	88.00	-	735.00	-	-
Photography	-	368.00	-	368.00	-	-		-	-	-		-	-	368.00	-	-
Subtotal		160,748.46	65,475.37	95,148.40	124.69	3,910.66	1.84	3,872.48	36.34	1,021.00	1.10	1,019.43	0.47	165,680.12	65,478.31	161.50
Operations																
Telephone	37.06	3,585.00	17.86	3,567.14	-	1,187.00	0.10	1,186.90	-	728.00	19.10	708.90	-	5,500.00	37.06	-
Postage	-	1,688.68	-	1,688.68	-	641.80		641.80	-	319.80		319.80	-	2,650.28	-	-
Garage Space Rental	-	463.00	-	463.00	-	184.00	-	184.00	-	88.00	-	88.00	-	735.00	-	-
Subtotal		5,736.68	17.86	5,718.82	-	2,012.80	0.10	2,012.70	-	1,135.80	19.10	1,116.70	-	8,885.28	37.06	-
Other																
Automobile mileage	144.77	2,758.46	144.77	2,613.69	-	-		-	-	-		-	-	2,758.46	144.77	-
Subscriptions/Dues	-	-	-	-	-	-		-	-	-		-	-	-	-	-
Permits/Fees	-	669.50	-	669.50	-	-		-	-	-		-	-	669.50	-	-
Advertising	-	396.92	-	396.92	-	-		-	-	-		-	-	396.92	-	-
Subtotal		3,824.88	144.77	3,680.11	-	-	-	-	-	-	-	-	-	3,824.88	144.77	-
Total	130,003.06	700,508.10	113,143.00	586,590.78	774.32	128,762.10	9,172.12	119,533.64	56.34	80,729.80	7,687.94	73,012.89	28.97	910,000.00	130,003.06	859.63

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - Totals

EVENT	VOLUNTEERS	SUPERVISORS	ACRES	COST-SHARE
YEARLY 2002 TOTAL	1836	145	773	\$ 456,698.26
YEARLY 2003 TOTAL	1262	105	713.19	\$ 729,744.47
YEARLY 2004 TOTAL	545	67	60.9	\$ 152,657.41
Grand Total:	3,643	317	1,547	\$1,339,100.14
REQUIRED, Result 1 total	1,000	75	1,000	\$ 691,772.00

GRG Volunteer Documentation

EVENT	DATE	VOLUNTEERS	SUPERVISORS	HOURS	COST-SHARE	COMMENTS
2002						
Spr 02 Invasive Species Training	3/30/2002		28	84	\$ 924.00	GRG vol time @ \$11/hr
Spr 02 Basic Training	4/20/2002		35	105	\$ 1,155.00	GRG vol time @ \$11/hr
East Mississippi Bluff-Desnoyer unit	5/4/2002	30	6	114	\$ 1,254.00	GRG vol time @ \$11/hr
Westside Blufflands-Smith Avenue	5/11/2002	125	21	459	\$ 5,049.00	GRG vol time @ \$11/hr
Native Shrub Identif. Training	6/5/2002		32	96	\$ 1,056.00	GRG vol time @ \$11/hr
Barge Terminal No. 1, phase 1	6/8/2002	35	3	117	\$ 1,287.00	GRG vol time @ \$11/hr
Harriet Island entrance planting	10/10/2002	134	5	426	\$ 4,686.00	GRG vol time @ \$11/hr
Training for public parks	8/20/2002			48	\$ 528.00	GRG vol time @ \$11/hr
Mississippi River Gorge--South Park	9/21/2002	150	15	510	\$ 5,610.00	GRG vol time @ \$11/hr
2002 Total					\$ 21,549.00	
2003						
Advanced Training	2/22/2003		19	57	\$ 627.00	GRG vol time @ \$11/hr
Advanced Training	4/12/2003		27	108	\$ 1,188.00	GRG vol time @ \$11/hr
Ford	4/26/2003	82		287	\$ 3,157.00	GRG vol time @ \$11/hr
Cherokee Prairie Implem	6/7/2003	10		30	\$ 330.00	GRG vol time @ \$11/hr
Mounds Park East Overlook	4/3/2003	55	6	185	\$ 2,035.00	GRG vol time @ \$11/hr
Westside Blufflands-Cherokee Park	4/3/2003	33		128	\$ 1,408.00	GRG vol time @ \$11/hr
Eagle Creek Invasive removal	5/17/2003	103	10	349	\$ 3,839.00	GRG vol time @ \$11/hr
Minnesota National Wildlife Refuge	10/11/2003	71	6	243	\$ 2,673.00	GRG vol time @ \$11/hr
Land O' Lakes	9/27/2003	50	5	170	\$ 1,870.00	GRG vol time @ \$11/hr
Rexam	8/20/2003	10		30	\$ 330.00	GRG vol time @ \$11/hr
Training St Paul Parks & Rec	9/9/2003	20		60	\$ 660.00	GRG vol time @ \$11/hr
Advanced Training	11/1/2003		16	64	\$ 704.00	GRG vol time @ \$11/hr
2003 Total					\$ 18,821.00	
2004						
Advanced Training	4/3/2004		13	26	\$ 286.00	GRG vol time @ \$11/hr
Crosby Event	2/21/2004	46	3	150	\$ 1,650.00	GRG vol time @ \$11/hr
Hastings River Flats	3/20/2004	87	8	293	\$ 3,223.00	GRG vol time @ \$11/hr
Brooklyn Park	5/1/2004	216	20	728	\$ 8,008.00	GRG vol time @ \$11/hr
Prospect Crest	5/22/2004	70	5	230	\$ 2,530.00	GRG vol time @ \$11/hr
St. St. Paul Levee	6/12/2004	126	18	450	\$ 4,950.00	GRG vol time @ \$11/hr
2004 Total					\$ 20,647.00	
GRAND TOTAL					\$ 61,017.00	

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2002

EVENT	DATE (if event)	VOLUNTEERS	SUPERVISORS	ACRES	COST-SHARE	COMMENTS
SPRING-SUMMER 2002						
Spr 02 Invasive Species Training	3/30/2002		28		\$ 924.00	GRG vol time @ \$11/hr
Spr 02 Basic Training	4/20/2002		35		\$ 1,155.00	GRG vol time @ \$11/hr
East Mississippi Bluff-Desnoyer unit	5/4/2002	30	6	20	\$ 3,806.00	City of St. Paul
					\$ 1,254.00	GRG vol time @ \$11/hr
Westside Blufflands-Smith Avenue	5/11/2002	125	21	8	\$ 40,805.00	City of St. Paul
					\$ 5,049.00	GRG vol time @ \$11/hr
Native Shrub Identif. Training	6/5/2002		32		\$ 1,056.00	GRG vol time @ \$11/hr
Barge Terminal No. 1, phase 1	6/8/2002	35	3	1	\$ 8,140.00	Wildlife Habitat Council cash
					\$ 1,287.00	GRG vol time @ \$11/hr
Carpenter Nature Center (FMR)		108		40	\$ 7,224.00	FMR
Pine Bend Bluffs (FMR)		15		54	\$ 50,000.00	FHR
					\$ 418.00	FHR volunteers
Hastings Riverfront plan (FMR)					\$ 7,300.00	FMR
SUBTOTAL Spring-Summer 2002		313	125	123	\$ 128,418.00	
FALL 2002						
Harriet Island entrance planting	10/10/2002	134	5	5	\$ 4,686.00	GRG vol time @ \$11/hr
Training for public parks	8/20/2002				\$ 528.00	GRG vol time @ \$11/hr
Mississippi River Gorge--South Park	9/21/2002	150	15	12	\$ 2,883.00	LCC
					\$ 5,610.00	GRG vol time @ \$11/hr
Mississippi River Gorge--other					\$ 5,170.00	MNRRA volunteers
					\$ 18,152.00	MNRRA inkind
					\$ 1,294.00	MNRRA tool purchases
Ford property crew buckthorn-cutting				20	\$ 89,466.00	Ford
Westside Blufflands-Cherokee Park		31	0	35	\$ 28,400.00	NFWF, federal
					\$ 6,868.00	WSCO inkind
					\$ 6,210.00	City of St. Paul inkind
Ordway Nature Area (FMR)	10/12/2002	68		7	\$ 17,074.00	Macalester inkind
					\$ 494.00	FMR
					\$ 2,244.00	FMR volunteers
Battle Creek north				35	\$ 45,000.00	MNRRA
Prospect Blvd.					\$ 4,709.00	City of St. Paul staff time
River Corrditor					\$ 2,000.00	survey staff/City of St. Paul
FMV 2002						
event totals:		1140		536	\$ 52,002.50	
non-state employee contributions:					\$ 24,979.02	
incurred expenses:					\$ 10,343.84	
mileage:					\$ 166.90	
FMV Projects:						
MN Valley Refuge--Louisville Swamp						
MN Valley Refuge--Long Meadow						
Belle Plaine						
Black Dog						
Visitor Center						
Pond Dakota Mission Park						
Minnegasco Dakota Station						
Fort Snelling						
SUBTOTAL Fall 2002		1523	20	650	\$ 328,280.26	
YEARLY 2002 TOTAL		1836	145	773	\$ 456,698.26	
REQUIRED, Result 1 2002		500	45	500	\$ 345,886.00	

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2003

EVENT	DATE(if event)	VOLUNTEER	SUPERVISOR	ACRES	COST-SHARE	COMMENTS
SPRING-SUMMER 2003						
Advanced Training	2/22/2003		19		\$ 627.00	GRG vol time @ \$11/hr
Advanced Training	4/12/2003		27		\$ 1,188.00	GRG vol time @ \$11/hr
Ford	4/26/2003	82		8.1	\$ 3,157.00	GRG vol time @ \$11/hr
					\$ 163,967.00	Ford
					\$ 7,954.59	Ford inkind
Minnesota National Wildlife Refuge:						
Louisville Oak Savanna Restoration	3/1/2003				\$3,653	USFWS Staff time
Ordway (FMR)	5/3/2003	62	4	1	\$ 12,670.00	Macalester inkind
					\$ 2,079.00	FMR vol time @ \$11/hr
Cherokee Prairie Implem	6/7/2003	10		13.3	\$ 330.00	GRG vol time @ \$11/hr
Mounds Park East Overlook	4/3/2003	55	6	0.125	\$ 6,433.00	City of St. Paul
					\$ 825.00	City vol time @ \$11/hr
					\$ 2,035.00	GRG vol time @ \$11/hr
Westside Blufflands-Cherokee Park	4/3/2003	33		35	\$ 1,408.00	GRG vol time @ \$11/hr
					\$ 17,000.00	NFWF 2003 Grant
Eagle Creek Invasive removal	5/17/2003	103	10	6	\$ 3,839.00	GRG vol time @ \$11/hr
Brooklyn Park River Park				12.8	\$46,000	MNRRA
35 West Water Street	6/10/2003			0.1	\$6,732	Mitigation
Crosby Exotic Removal					\$40,000	MNRRA
FMV 2003						
event totals:		448		314	\$ 15,697.00	
non-state employee contributions:					\$ 28,566.26	
incurred expenses:					\$ 2,363.34	
mileage:					\$ 83.96	
FMV Projects:						
Savage Fen						
Pond Dakota Mission Park						
Minnegasco Dakota Station						
Fort Snelling/Pike Island						
Long Meadow						
SUBTOTAL Spring-Summer 2003						
		793	66	390.425	\$ 366,608.15	
FALL 2003						
Pine Bend Bluffs (FMR) event	10/25/2003	100	10	7	\$ 50,000.00	FHR cash
					\$3,630	vol time @ \$11/hr
		8			\$88	FHR vol time @ \$11/hr
					\$3,456	FHR food, t-shirts, table tent & chair rental, biffs
South St. Paul Levee				11	\$7,000	City cash
					\$49,000	MNRRA cost share
Hastings Riverfront Park (FMR)				250	\$ 24,531.57	NAWCA
					\$ 1,168.00	Hastings city staff time
					\$ 1,800.00	Wetland plants
		35			\$ 990.00	vol time @ \$11/hr
3M (FMR)					\$ 9,460.00	3M cash (FMR)
Sand Coulee (FMR)				4.26	\$ 3,000.00	Landowner contrib.
		10			\$ 275.00	vol time @ \$11/hr
Minnesota National Wildlife Refuge	10/11/2003	71	6	30	\$ 2,673.00	GRG vol time @ \$11/hr
					\$ 2,000.00	plant material Refuge
					\$ 3,087.00	MN Refuge staff time
Land O' Lakes	9/26/2003	50	5	1	\$ 167,000.00	DCSWCD
					\$ 1,870.00	GRG vol time @ \$11/hr
Battle Creek	9/2/2003	165	2	19	\$ 5,533.00	cash and inkind
Rexam	8/19/2003	10		0.5	\$14,250	Mitigation/Cash
					\$330	GRG vol time @ \$11/hr
Training St Paul Parks & Rec	9/9/2003	20			\$660	GRG vol time @ \$11/hr
Advanced Training	11/1/2003		16		\$704	GRG vol time @ \$11/hr
Saint Paul Park & Rec. Projects:						
Cherokee					\$5,038	Staff time
Crosby					\$948	Staff time
Desnoyer					\$1,566	Staff time
Mounds					\$1,385	Staff time
Prospect Crest					\$1,484	Staff time
Lilydale Wetland					\$210	Staff time
SUBTOTAL Fall 2003						
		469	39	322.76	\$ 363,136.32	
YEARLY 2003 TOTAL						
		1262	105	713.185	\$ 729,744.47	
REQUIRED, Result 1 2003						
		500	30	500	\$ 345,886.00	

Great River Greening
Big Rivers Partnership
LCMR Costshare Summary - 2004

EVENT	DATE(if event)	VOLUNTEER	SUPERVISO	ACRES	COST-SHARE	COMMENTS
2004						
Crosby	2/21/2004				\$ 10,000	Carolyn Foundation
		46	3	4.5	\$ 1,650	GRG vol time @ \$11/hr
Advanced Training	4/3/2004		13		\$ 286	GRG vol time @ \$11/hr
Hastings River Flats	3/20/2004	87	8	11	\$ 3,223	GRG vol time @ \$11/hr
					\$ 1,459	FMR
Brooklyn Park River Park	5/1/2004	216	20	3	\$ 8,008	GRG vol time @ \$11/hr
					\$9,848	MNRRRA staff time
					\$81,000	MNRRRA
Prospect Crest	5/22/2004	70	5	0.25	\$ 2,530.00	GRG vol time @ \$11/hr
South St. Paul Levee	6/12/2004	126	18	0.25	\$ 4,950.00	GRG vol time @ \$11/hr
Flint Hills	5/1/2004			38	\$ 21,514	FHR
Dayton Install	5/1/2004			3.9	\$ 8,190	City of Dayton
SUBTOTAL, 2004		545	67	60.9	\$ 152,657	
YEARLY 2004 TOTAL		545	67	60.9	\$ 152,657	